be explained by a mild hypovitaminosis-K in certain individuals. Kark and Lozner (1939) showed that this hypovitaminosis could in fact occur in adults owing to causes other than liver disease. In any case it would seem worth while giving vitamin K a further trial in this widespread and irritating condition.

Summary

The treatment of eight cases of chilblains, ranging from the mildest manifestation to severe and ulcerated forms, is described. In four there was complete alleviation of symptoms and signs, the response in Case 1 being exceptionally good. In the other four there was an improvement. Two of these patients defaulted before treatment was completed, so the final result was not known. In the remaining two there was improvement, especially of the actual chilblains, although the pernicious condition of the digits remained; and in one there was a recrudescence of symptoms in spite of continued treatment.

As regards the mode of administration it was considered that intramuscular injections were superior to the oral route. They were, however, accompanied by considerable pain, so they had to be abandoned, as the patients were not prepared to put up with this for the possible cure of their relatively minor ailment.

REFERENCES

- Black, J. H. (1945). *J. Allergy*, **16**, 83.
 Howell, W. H. (1941). *J. Amer. med. Ass.*, **117**, 1059.
 Kark, R., and Lozner, E. L. (1939). *Lancet*, **2**, 1162.
 MacKenna, R. W. (1937). *Diseases of the Skin*, 4th ed., p. 291, London.
 Scarborough H., and Macfarlane, R. G. (1942). *Proc. roy. Soc. Med.*, **35**, 407.
 Stewart, J. D., and Rourke, G. M. (1939). *New Engl. J. Med.*, **221**, 403.

CALCIUM DEPOSITS IN THE IRIS IN A CASE OF SECONDARY HYPERPARATHYROIDISM

BY

C. PONIEDEL, M.B., Ch.B.

Late Senior Physician, Ministry of Pensions Childwall Hospital,
Liverpool

The following case is of interest for two reasons: (a) a review of the literature does not reveal any previous reference to such deposits in these cases; (b) renal calculi and metastatic calcification were absent from the kidneys.

Oliver (1939) refers to three cases of hyperparathyroidism, as follows. (i) Dawson and Struthers (1923). A man aged 49 with a long history of multiple fractures, ending in syncope and death; necropsy showed a parathyroid tumour, calcification of vessels (heart, lungs, kidney, liver, spleen). (ii) Lowenburg and Ginsberg (1932) describe the case of a boy aged 5 who received an overdose of parathormone, 100 units daily for 6 days, with resultant vomiting and listlessness after two days, followed by mental depression and high fever. Blood examination on the sixth day showed serum calcium 19.6 and blood phosphorus 4.4 mg. per 100 ml. On withdrawal of the drug the patient became normal. (iii) Hanes (1939) mentions a woman aged 49 who complained of weakness, pains in the legs, and loss of weight. Previous radiography of the kidneys, when in hospital for pyelitis, had shown diffuse fine mottling. Serum calcium was 12 mg. per 100 ml. and renal functions were markedly impaired. Five years later, on final admission to hospital, she had a tumour of the lower pole of the thyroid. Blood calcium was 20 mg., phosphorus 4.7 mg., phosphates 23 mg., and non-protein nitrogen 58 mg. per 100 ml. Skiagrams showed metastatic calcification in the kidneys and decalcification of bones. The patient became weak, depressed, and febrile, and she soon died. Necropsy showed a necrotic partly calcified parathyroid tumour.

Moore (1943) states that calcification had been reported in many organs in hyperparathyroidism, but not in bone-marrow. He mentions calcification in kidney, stomach, heart, liver, and marrow found in experiments on animals. Werner (1942) describes metastatic deposits in the kidney, and says they can occur in other organs and tissues (none mentioned specifically). He also quotes Allbright, Baird, Cope, and Bloomberg for renal calculi; and Dawson and Struthers as finding calcium infiltration in practically every organ of the body. Jaffe (1940) states that secondary parathyroid hyperplasia is found more regularly in connexion with chronic renal insufficiency than in any other condition, and that the hyperfunction becomes severe enough to induce pronounced skeletal and other tissue changes, the chief lesions being skeletal and renal (calculi). Nephrosclerosis is less common. Severe renal insufficiency is followed by calcification in soft parts (subcutaneous tissues and arteries).

Case History

The patient had a history of having been taken ill in October, 1944, while serving in the East. At that time he complained of loss of weight, loss of appetite, frequency of micturition, and looseness of the teeth. He was admitted to hospital, where the urine was found to be of low specific gravity and to contain albumin and hyaline and granular casts. Skiagrams of the kidneys, after intravenous "uroselectan," failed to reveal the outline of any renal tissue. The blood urea was 140 mg. per 100 ml.

In November, 1944, he became drowsy and the skin dry. At this period he developed a left-sided Bell's palsy; and the ear-drums were found to contain chalky deposits. Soon afterwards vision became blurred, and examination revealed a chalky deposit in the anterior chamber of the right eye and retina—choroidal atrophy. By this time the rough skin showed nodularity, which appeared to be due to chalk-like deposits. Blood examination revealed a serum calcium of 17 mg. and inorganic phosphorus of 4.3 mg. per 100 ml.; the phosphate index was normal. Radiology of the long bones showed widespread osteoporosis.

By February, 1945, the serum phosphate was 22 units (normal 3–13), and a diagnosis of hyperparathyroidism was made. Blood calcium and blood urea could be controlled temporarily by reducing intake of phosphates. The blood calcium fell to 11 mg. and the blood urea to 82 mg. per 100 ml., only to rise again in two weeks. Primary hyperparathyroidism was considered to be the diagnosis.

In June, 1945, the urine volume was 3½ pints (2 litres) daily, S.G. 1012 or less. Urea clearance was 30% of normal, and the blood urea 80 mg. per 100 ml. X-ray investigation showed well-marked calcification of the peripheral vascular tree in places. His general condition had improved, although wasting of the legs was extreme, and the diagnosis of primary hyperparathyroidism was thought to be incorrect because of the normal serum phosphates, absence of urinary calculi, and absence of skeletal weakness (neither fractures nor bowing of long bones).

In February, 1946, the patient was admitted to Childwall Hospital for review. On admission he complained of loss of balance when walking. He was emaciated and there were calcium deposits beneath the nails. Disks and ear-drums were as on previous occasions. In addition the irises showed fine deposits of calcium. Urea concentration was 1.3%, and faeces calcium 60% (normal reading 80% to 90%). Skiagrams of arteries showed slight calcification in the walls of the arteries of both hands, and those of the skull, spine, and long bones revealed generalized slight osteoporosis. Blood calcium amounted to 14 mg., blood urea 52 mg., serum albumin 4.5 mg., serum globulin 1.5 mg., and inorganic phosphate 3.5 mg. per 100 ml. The phosphate index was normal; there was a secondary anaemia with moderate eosinophilia. Examination of the urine showed protein 1%; sodium chloride, 6.5 mg. in 24 hours; calcium oxalate ++, hyaline casts +, and a few granular casts; red cells were occasionally present.

Progress of the Case.—Owing to the emaciation present it was considered advisable to put the patient on a fuller diet about

mid-March. His blood urea kept about 60 mg. per 100 ml. and his progress was satisfactory. However, on April 3 he was found to be strange in manner and unresponsive when questioned, and further probing showed that he was suffering from delusions. A diagnosis of uraemia was made and he was given glucose and fruit drinks up to 3 pints (1.7 litres) a day and a diaphoretic mixture; the bowels were attended to, and he also received 1/2 gr. (32 mg.) phenobarbitone s.o.s. He slept well that night and was rational the next day. The blood urea reading was 86 mg. per 100 ml. on April 3, and had fallen to 66 mg. the following day. The blood calcium was 11 mg. per 100 ml. (normal). He usually passed 20 oz. (568 ml.) of urine by day and 26 oz. (740 ml.) at night when receiving normal diet and fluids. This indicated a lack of concentrating power of the kidneys. Shortly afterwards he took his discharge from hospital against advice.

On March 8 the patient had developed auricular fibrillation, which responded to digitalis treatment, the drug being discontinued a month later. His pulse remained regular and the heart steady.

Comment

The evidence was in favour of a diagnosis of primary chronic renal disease with secondary increase of calcium in the blood. The fact that the blood calcium had become normal was against a primary hyperparathyroidism, as it is usually stated that without operation a primary condition of the parathyroid would not improve. Price mentions that hyperparathyroidism occasionally occurs as a result of chronic nephritis.

Interesting points were (1) the presence of deposits in the irises; (2) the absence of renal calculi; (3) a temporary attack of auricular fibrillation associated with uraemia.

I wish to thank the Director-General of Medical Services, Ministry of Pensions, for permission to publish this article.

REFERENCES

- Dawson, J. W., and Struthers, J. W. (1923). *Edinb. med. J.*, **30**, 421.
 Hanes, F. M. (1939). *Amer. J. med. Sci.*, **197**, 85.
 Jaffe, H. L. (1940). *Bull. N.Y. Acad. Med.*, **16**, 291.
 Lowenburg, H., and Ginsberg, T. M. (1932). *J. Amer. med. Ass.*, **99**, 1166.
 Moore, R. D. (1943). *Arch. Path.*, **36**, 51.
 Oliver, W. A. (1939). *St. Bart's Hosp. Rep.*, **72**, 225.
 Werner, A. A. (1942). *Sth. med. J.*, Nashville, **35**, 671.

BABCOCK'S OPERATION FOR THORACIC ANEURYSM

BY

F. T. RANSON, M.B., F.R.C.S.Ed.
 Shanghai

The operation described by Babcock (1926; 1932) consists of an anastomosis between the common carotid artery and the internal jugular vein of one side. Former operations for these aneurysms had used the principles of reduction of flow of blood through the sac or else the introduction of foreign bodies into the sac. Babcock's operation is based on the principle of speeding up the flow of blood through the sac and thus reducing the pressure of blood inside the aneurysm.

I have had the opportunity of performing this operation in eight cases. The patients were all male Chinese with aneurysms eroding the anterior chest wall; all complained of severe pain and all had a positive Wassermann reaction.

Method

The operation is performed as follows. Under local anaesthesia a transverse incision is made in the neck (usually right side) about 1½ in. (3.75 cm.) above the clavicle. The sterno-mastoid muscle is cut across and the common carotid artery and internal jugular vein are dissected out and

cleaned. Both vessels are ligated high up, and umbilical tapes are placed around them low down so as to control bleeding. The vessels are divided just below the ligatures and the heart ends are anastomosed. After slowly releasing the tapes and seeing that the anastomosis is watertight the wound is closed.

McCarthy (1930) in an excellent detailed article on the subject discusses the hydromechanics of the operation. Half the circulation to the head is short-circuited back into the heart, and this removes the resistance of all vessels above the anastomosis. Added to the removal of this resistance is the aspiration effect exerted by venous suction. This results in lowered pressure and increased blood velocity inside the sac.

McCarthy uses the analogy of the garden hose with a weak place in its wall. If the water pressure is high and if the spigot of the hose is partially obstructed, the hose will tend to bulge at the weak spot. If, however, the spigot is opened widely, the pressure inside the hose will be reduced, the velocity of the water increased, and the bulge will tend to subside.

Results

Whatever happens inside the aneurysm, there is one striking immediate result of the operation, and that is relief of pain. This disappearance of pain took place in all but one of my cases and was remarked on by the patients immediately after release of the controlling tapes.

Of the eight patients seven left the hospital improved and relieved of pain. The eighth, who had a swelling as big as a fist on the chest wall, died of external rupture of the aneurysm five days after operation. He had not experienced any relief of pain.

In a personal communication in 1946 Babcock wrote: "I have now performed between 26 and 30 of these operations. As some have felt that the relief is due to the division of fibres connected with the carotid plexus, I have had opportunity to note the effect upon pain in four cases of large abdominal aneurysms in which an anastomosis was made between the heart ends of the external iliac artery and vein. In the abdominal cases it was possible to see the effect of the operation upon the aneurysm characterized by a prompt partial collapse of the sac immediately after the systolic wave, showing that prolonged systolic tension was not possible with the large leak back into the veins that had been established. The relief from pain was also striking, particularly in one patient who had required large doses of morphine at frequent intervals before the operation and had complete relief after operation." The operation is an easy one to those familiar with the technique of vessel suture.

At a surgical meeting in Chicago I heard the procedure described by an eminent and thoughtful surgeon as unjustified. It cannot be expected to cure the patient, but surely it is something to be able to relieve his pain, and this is often a distressful feature of these cases.

The only other operation for thoracic aneurysm of which I have had experience is wiring, but I have not observed any relief of pain with this method.

REFERENCES

- Babcock, W. W. (1926). *Ann. clin. Med.*, **4**, 933.
 — (1932). *Amer. J. Surg.*, **16**, 401.
 McCarthy, P. A. (1930). *Ann. Surg.*, **91**, 161.

The Central Office of Information has produced two films on child psychology for parents. The first, "Children Learning by Experience," is a study of children learning in their own way and was designed for teachers in training to provide material for discussion. The second, "Children Growing up with Other People," depicts children's development towards a balance between individualism and co-operativeness.