Many Russians in my care during the war, even those who had been wounded before capture, awaited their return home with extreme terror. Why?

I agree that social security, whatever it may mean, has tended to reduce effort for the reasons advanced by Dr. Crichton-Miller. Many economists suggest that the economic basis for this social security is crumbling, so we may soon see the next phase developing.—I am, etc.,

Barnet, Herts.

G. C. PETHER.

Prognosis of Pleural Effusion

S.R.—Dr. Brian C. Thompson's reply to Dr. J. D. L. Reinhold's letter (Aug. 30, p. 348) prompts me to add my views on the point at issue in this important correspondence. A study of my cases in private and hospital practice during the last eight years goes to support Dr. Thompson's statement that "there was no evidence that differences in treatment of the original pleurisy affected the subsequent incidence of pulmonary tuberculosis." The study brings out two points conspicuously:

(1) The subsequent development of pulmonary tuberculosis is more common in those who show a "tuberculous diathesis." The latter can be recognized clinically to some extent from the history of tuberculosis in the family, relatively low weight, inelastic skin, dry hair, pallor, easy fatigue, "phthisical" configuration of the chest. Such persons are often styled by laymen as of "delicate constitution."

(2) Rest and sanatorium line of treatment have no effect on the prognosis. Under Indian conditions patients of pleural effusion, when afebrile and free from respiratory embarrassment, have to work with fluid still in the pleural cavity. They cannot afford to take rest till, as Thompson writes, "their blood sedimentation rate becomes normal." Under these conditions many have not developed pulmonary tuberculosis for over five years. Two female patients had one and two deliveries during this period, in spite of medical advice, without untoward effect. One patient who had effusion on one side and subsequently on the other is quite well and working now three years after.

This has been my impression based on clinical experience but unsupported by statistical data. I have come to believe that development of pulmonary tuberculosis subsequent to pleural effusion depends mainly on the "soil" and the resistance of the patient, to less extent on the environment and conditions of life of the patient, and to a negligible extent, if at all, on the rest and sanatorium treatment as held by Drs. Reinhold and Coope.—I am, etc.

Poona City, India.

P. L. DESHMUKH.

Early Diagnosis of Tuberculous Meningitis

SIR,—A patient was recently admitted for streptomycin treatment of a tuberculous meningitis only after he had been retained one week in another hospital because the chlorides in the cerebrospinal fluid (C.S.F.) had been found to be 0.72% and the diagnosis of tuberculous meningitis, to which other facts pointed, had consequently been rejected. A clear answer seems to be needed to this question: Given a patient with a meningitic picture whose C.S.F. shows a moderate pleocytosis, does a knowledge of the C.S.F. chloride level materially help to distinguish those cases suffering from tuberculous meningitis?

The chloride levels of the first specimen of C.S.F. examined in 52 cases of tuberculous meningitis, 8 cases of benign lymphocytic meningitis, and 10 cases of anterior poliomyelitis (all children) were as follows:

C.S.F. Chlorides	Tuberculous Meningitis Cases	Benign Lymphocytic Meningitis Cases	Acute Poliomyelitis Cases
0.70% or over ..	11	5	10
0.65–0.69% ..	23	3	—
0.60–0.64% ..	18	—	—
Total ..	52	8	10

It is seen that in 1 in 5 cases of tuberculous meningitis the C.S.F. chloride level was not lower than normal, while in nearly half the cases the chloride level fell within the zone of slight reduction (0.65–0.69%) in common with several of the cases of lymphocytic meningitis. In only a third of the cases of tuberculous meningitis was the chloride level substantially reduced

(0.60–0.64%), and in none did it reach the level of 0.59%, which is accepted as diagnostic of the condition. Thus, while a substantial lowering of the C.S.F. chloride level to 0.64% or less would, under the circumstances postulated, favour a diagnosis of tuberculous meningitis, the finding of a normal C.S.F. chloride level *should in no wise argue against tuberculous meningitis.*

All in all, I suggest that the time devoted by a clinical pathologist to an estimation of the C.S.F. chlorides would be more profitably spent in a search for acid-fast bacilli.—I am, etc.,

Newcastle-upon-Tyne.

DOUGLAS GAIRDNER.

Reiter's Syndrome

SIR,—As Reiter's syndrome occurs apparently mainly in young adults, I may be permitted to put the following case on record.

On July 23, 1945, an engineer, 55 years of age, consulted me for a purulent urethritis. No gonococci were found in the discharge. On Aug. 3, after 20 g. of sulphathiazole had been given, the discharge had practically ceased, and the smear contained only epithelial cells; but by that time the patient had a bilateral purulent conjunctivitis, again without gonococci in the smears.

On Aug. 10 the eyes were nearly normal, but he then had a painful swelling of an interphalangeal joint of the right foot, which took about 10 days to disappear. On Aug. 15 he started to complain of pain across the shoulders and in the lumbar region, without any obvious pathology, possibly due to a spinal arthritis.

He then went to his home in the North of England, and told me on his return that those pains did not leave him completely for more than two months. In the North a film of his spine was taken in which no changes were seen. He had no diarrhoea either before or during his illness.—I am, etc.,

Harrow Weald, Middlesex.

ALFRED LETCHNER.

Safer Administration of Intravenous Anaesthetics

SIR,—A method for increasing the safety in administering intravenous anaesthetics consists in giving the patient oxygen to inhale for two or three minutes before and during the administration of the intravenous anaesthetic. It is, in my opinion, particularly indicated in patients over 60 or ones who are sub-normal risks.

The rationale of this method is as follows. With most intravenous drugs given for anaesthetic purposes, and particularly with thiopentone, one is struck by the initial and severe depression of respiration that occurs. It is, I think, inevitable that some degree of anoxaemia must ensue, and it is this factor that endangers the patient. In many patients, particularly the elderly, in whom alarming symptoms are most likely to occur with these agents, there is the factor that their lungs have lost their full efficiency and that oxygenation of the blood is neither so quick nor so easy to bring about by the usual methods. If, however, one makes certain that the respiratory system is flooded with a high percentage of oxygen before this period of respiratory depression, then the period will do little harm.

In actual practice I have found that to all appearances my hopes have been justified; the patients have kept an excellent colour and there has been no cause for worry. Occasionally the patient has been told that "he won't have anything put over his face to put him to sleep"; then I assure him that what I am giving him to inhale will not put him to sleep but will help.—I am, etc.,

London, N.W.7.

D. C. CLARK.

Morphine in a Patent Medicine

SIR,—I should like to comment on the medico-legal note which appeared in your issue of Sept. 27 (p. 511), and referred to the death of a 2-year-old child after taking a patent medicine containing morphine. I gave evidence to the effect that the bottle when full contained just under 1 gr. (65 mg.) of morphine, but according to the *Daily Mail* (June 25) the manager of the firm which manufactures the medicine denied this, alleging that each bottle contained 0.432 gr. (28 mg.). In point of fact the incriminated bottle when full would contain 6 oz. (170 ml.) of fluid and as the concentration of morphine was 0.033% (as stated on