SIR,—In all the literature on this matter that I have read the general explanation of reduced efficiency of penicillin seems to indicate that the bacteria have acquired a tolerance to penicillin by reason of small doses given in mild conditions, usually by local application. It ill becomes a general practitioner with no special knowledge of pharmacology to dispute this explanation, but it has occurred to me that the penicillin itself may have lost some of its potency by reason of long-continued cultivation of *Penicillium notatum* in the laboratory. The analogy of a bacterial culture of, say, *B. pestis* losing some of its virulence by culture *in vitro* might apply to penicillin.

After all, penicillin is a biological product and not a synthesized substance, and its rapid production under artificial conditions might adversely affect its potency. *P. notatum* in its natural state feeds on a large variety of organic substances, but commercially it is bred presumably on a very limited number of media. Can it be that *P. notatum* (like some of us) is getting tired of its monotonous diet and is losing some of its virility, if the latter quality can be attributed to a vegetable? Whatever the real explanation may be, it is certainly time that the experts investigated the matter. I am beginning to feel guilty whenever I prescribe a bottle of troch. penicillin.—I am, etc.,

Holcombe Rogus, Somerset.

J. V. MAINPRISE.

Staphylococcal Infection by Penicillin-resistant Strains

SIR,—An important cause of the multiplication of penicillin-resistant strains of *Staphylococcus pyogenes* is the indiscriminate local use of penicillin in the treatment of skin diseases. Penicillin, at any rate in cream or ointment form, has little or no therapeutic value even in conditions generally regarded as superficial infections by pus cocci. Moreover, penicillin is responsible more or less directly for the conversion of many trivial and benign dermatoses into severe and troublesome cases of acute dermatitis.

Penicillin, originally puffed as a talisman of immunity on the battlefield, has been the subject of such intense advertisement that it is demanded by patients who, thanks to 50 years of compulsory education, can more or less recognizably lisp its name. Our figures at the Skin Hospital correspond with those of Dr. Mary Barber (Nov. 29, p. 863). We have also seen several penicillin-resistant β -haemolytic streptococci.—I am, etc.,

Manchester.

J. H. TWISTON DAVIES.

Penicillin in Scarlet Fever

SIR,—I was interested to read your annotation in the *Journal* of Dec. 6 (p. 915) on "Penicillin in Scarlet Fever," as I was the writer (French) of the paper in the *Journal of Hygiene* to which you make reference. I have to point out, however, that there is a printer's error in the reference you quote, as my paper on scarlet fever was published not in 1945 as stated, but in 1939 (Sept. 16). I think this is important, inasmuch as if my conclusions on the value of sulphonamides in scarlet fever stand unrefuted after a period of eight years more weight may be given to them than if they were only unrefuted for two years.

I may add that I have had a considerable experience of the treatment of scarlet fever since 1939, and I am still quite convinced that the sulphonamides are of little or no value—even the newer and less toxic varieties that have appeared since 1939. I am still equally far from knowing why they should be so useless. Perhaps someone with more opportunity than I now have will be able to elucidate this. I would also like to know the results of a properly controlled therapeutic trial of penicillin in scarlet fever. If you know of any such I would be much obliged if you would tell me about it.—I am, etc.,

Darwen, Lancs.

JANE O. MILLAR.

Measles

SIR,—May I support Dr. Harwood Stevenson's advocacy (Dec. 6, p. 928) of the use of parental whole blood as a routine measure in the attenuation of measles in contacts, and particularly in the dangerous period up to the age of 5? It is a method particularly applicable to general practice, where as a rule attenuation and not prevention is the objective. There

can be little doubt that if this procedure were widely used there would be an appreciable drop in the death rate from measles.

In your leading article of June 29, 1946 (p. 991), there occurs this startling statement: "Despite the success which has been achieved in lowering the mortality from this disease it remains one of the largest causes of death among children." This fact alone would justify any safe method which could reasonably be expected to attenuate the disease, and in the *Journal* of June 30, 1945 (p. 923), I described a series of cases of 25 contacts so treated with most satisfactory results. An earlier series of cases of successful attenuation was described by Dr. T. D. Culbert (*British Medical Journal*, 1938, 2, 705). As regards dosage, I feel that if disappointment is to be avoided not less than 8 ml. of whole blood should be injected, and that it should not be reduced on account of the age of the child, for it is after all in the youngest contacts that attenuation is most important. (I do not think one would reduce the dose of diphtheria antitoxin because of the age of the patient.)

Finally, I would venture to suggest that Dr. Stevenson is a little optimistic if he expects any material attenuation when the blood is given "at any time up to the sixth or seventh day of incubation." This would probably succeed if one were using convalescent serum, but it would almost certainly fail to produce satisfactory attenuation in the case of whole blood and might well bring the method into disrepute. In my experience if it is to be really effective it should be given forthwith—that is to say, on the day when the fact of contact has been established or as soon after as possible.—I am, etc.,

London, N.W.7.

A. H. MORLEY.

Dogs and Poliomyelitis

SIR,—Dr. Norman Macfadyen's letter (Dec. 6, p. 930) is interesting because a paragraph in an evening paper a few days ago stated there has been an epidemic of paralysis in dogs in the eastern counties and that 300 cases had been reported. I have two "pekes," and last June one of them, after a day's malaise, became partially paralysed in her hind legs and tail. Recovery ensued in 14 days. Then a month afterwards the other, after 24 hours of lassitude, became totally paralysed in both hind legs and tail.

The lesion must have been in the dorsal cord, for both hind legs and the tail, and also the abdominal muscles, were spastic. The bladder function was unimpaired but the bowel was difficult from the loss of the muscles accessory to defaecation. The dog was very ill for a week and then got better, but the spastic paralysis remained unchanged for over two months and then recovery ensued, though the hind legs are still weak. Injury can be absolutely ruled out; and both dogs are beyond the age for distemper, and, moreover, there were no symptoms of it. It seems pretty certain that the condition was due to a virus, and the occurrence of such cases at the same time that an epidemic of anterior poliomyelitis is affecting the country is surely suggestive.—I am, etc.,

London, W.1.

VICTOR BONNEY.

Chronic Sinusitis in Children

SIR,—Mr. F. M. Walker (Dec. 6, p. 908) raises the question of the treatment of chronic nasal sinusitis in children. This matter has been receiving attention at Warwick Hospital for some years, and the following observations may prove of interest.

In order to obviate the necessity of giving repeated general anaesthetics for antral washouts in cases that have not been cured by the use of nasal drops and attention to the general health, we have been placing a ureteric catheter into each infected antrum, some inches of the tube being strapped on to the opposite cheek, the extra-nasal end being wrapped in sterile gauze, following the technique described by R. H. Hunt Williams (*Proc. roy. Soc. Med.*, 1946, 39, 280). The antra have then been washed out from one to several times daily. Solutions we have been using are: "soluthiazole" 5%; ephedrine 0.5%. "flavazole" 1 in 2,500, with or without penicillin, 500 units per ml., in normal saline; isotonic or slightly hypertonic sodium chloride; proflavine 1 in 2,000 in normal saline.

In a recent American paper (the exact reference to which I am unable to trace at present) evidence was presented indicating that isotonic or somewhat hypertonic saline solutions were as effective therapeutically as solutions containing chemotherapeutic agents. On