A.I.H. will be successful in most, or in hardly any case, and the same applies to A.I.D. A few illustrations are given below:

1. *A.I.H.*—In a recent series of 30 successive cases A.I.H. was carried out because of low invasive power of the semen. In addition, spermatozoal density was low (less than 10 millions per c.cm.) in six cases; and more than 15% abnormal head forms were present in the semen of five cases. Female infertility factors—e.g., impaired tubal function, infrequent ovulation, etc.—were detected or strongly suspected in six cases. Insemination was carried out in at least four successive cycles where necessary and, as a rule, more than once in each cycle. One case is counted twice, for the patient miscarried after the first series of inseminations, and then made a second attempt. Nine conceptions occurred—in one case during the first cycle, in the others in the third or fourth cycle. Four pregnancies terminated in miscarriage in the fourth month of pregnancy or earlier; four others resulted in the birth of a viable infant. In the last case the pregnancy continues at the time of writing. In 21 cases no conception was recorded with certainty, but two conceptions may have occurred (? early miscarriage). The negative cases are characterized by a higher average age of the wives, but not by a higher frequency or greater severity of manifest infertility factors in either husband or wife.

2. *A.I.D.*—A recent series of 15 successive cases in which donor M. was employed consisted of women in whom no serious infertility factor was discovered, though four of them were 40 years of age or over. Insemination was repeated where necessary in at least six cycles. Conception resulted in 10 cases. The negative cases were represented by one woman in whom insemination was repeated during eight cycles and then abandoned, and the women aged 40 years or over. The fate of the 10 pregnancies varied. One was tubal and had to be terminated; one ended in miscarriage (end of the second month); five went to term uneventfully or are nearing term at present; and three terminated successfully at 45 weeks. The number of cycles during which inseminations were carried out in the positive cases varied from one to seven. It may be said that, in our experience, repeated A.I.D. is nearly always successful in reasonably young women in whom infertility factors have not been detected. The infants produced by donor M. in this as in another series show no abnormality and are progressing favourably. Their physical types vary widely, and the parents state in every case that they are glad to have availed themselves of this service.

Summary

In some cases of male impotence, dyspareunia, ejaculatory failure, and defective, cervical invasion, artificial insemination with the husband's semen may be indicated. In the first three conditions named, self-insemination by the wife is possible.

Artificial insemination with donated semen has been carried out in cases of incurable male sterility and for genetic reasons.

The procedure and technique of artificial insemination with the husband's semen and artificial insemination with donated semen (choice of cycle day, suitability of semen, method of cervical instillation) are described.

The selection of semen donors and the organization of artificial insemination with donated semen are discussed.

Some potential dangers of artificial insemination—e.g., uterine infection, dysgenic conception from defective or damaged sperms—are enumerated.

Two small series of cases are reviewed.

REFERENCES

- Barton, M., and Wiesner, B. P. (1944). *Lancet*, **2**, 563.
 Bickers, W. (1944). *Amer. J. Obstet. Gynec.*, **47**, 425.
 Cary, W. H. (1940). *J. Amer. med. Ass.*, **114**, 2183.
 DeLee, B. (1924). *Principles and Practice of Obstetrics*, 4th ed., Philadelphia.
 Green-Armytage, V. B. (1943). *Proc. roy. Soc. Med.*, **36**, 105.
 Guttmacher, A. F. (1942). *J. Amer. med. Ass.*, **120**, 442.
 Hamilton (1925). *Industrial Poisons in the United States*, New York.
 Lane-Roberts, C., Sharman, A., Walker, K., and Wiesner, B. P. (1939). *Sterility and Impaired Fertility*, London.
 Seguy, J., and Vimeux, J. (1933). *Gynec. Obstet.*, **27**, 346.
 Sims, J. M. (1868). *British Medical Journal*, **2**, 465, 492.

A search for methods of controlling Scotland's "plague of midges" will be the task of a small committee of Scottish scientists headed by Brig. F. A. E. Crew, professor of public health and social medicine at Edinburgh University, and director of biological research at the War Office. Mr. Thomas Johnston, Secretary of State for Scotland, invited the committee to consider whether measures similar to those used in the Army against mosquitoes and other insect pests dangerous to health could usefully be employed against midges and other biting flies. The committee's first step will be to review what is known about methods of attack on insects and their breeding grounds and to advise whether these methods can be used or can be modified for use against midges.

METHYL BROMIDE POISONING EFFECTS ON THE NERVOUS SYSTEM

BY

A. BARHAM CARTER, M.D., M.R.C.P., D.P.M.

Physician to Staines County Hospital

Methyl bromide (CH_3Br), a non-inflammable colourless gas at ordinary temperatures and pressures, fairly easily compressible into liquid form, is used extensively as an insecticide, fire extinguisher, fumigant, and refrigerant. It is known to be toxic to man in various ways, and as it is without smell in low concentrations and is 3.3 times as heavy as air it constitutes a very real industrial hazard. The toxic effects described in the literature are: (1) on the skin, giving rise to a vesicular irritative dermatitis and second-degree burns; (2) on the respiratory system, producing bronchitis, broncho-pneumonia, pulmonary congestion and oedema; (3) on the alimentary system, with gastro-intestinal upset and hepatic dysfunction; (4) on the nervous system, with both an acute and a chronic symptomatology. It is with the last that this paper deals, and although references show a familiarity with other manifestations, no attempt has been made to evaluate or define the more chronic effects on the central nervous system.

Review of Literature

Heiman (1944) has given a full review of the literature (chiefly Continental) concerning the general toxic effects of methyl bromide on man. An analysis of cases describing the acute neuropathies produced by short exposure to high concentrations shows in the majority a prodromal stage of headache, dizziness, nausea, vomiting, and generalized weakness with transient diplopia and amblyopia, followed by a period of mental excitement with focal or generalized epileptiform convulsions, muscular tremors and spasms, mental confusion, psychomotor acceleration, and restlessness, amounting at times to acute mania. This stage is accompanied by fever and occasionally cyanosis with methaemoglobinemia, and in fatal cases is followed by coma, increase in muscular spasms, and convulsive seizures with death in hyperpyrexia.

If the patient survives, there follows a long period of convalescence, lasting up to 18 months in a few cases, in which the epileptiform attacks become less frequent, the mental confusion and excitement gradually abate, and visual disturbances slowly improve, although nystagmus and sluggishly reacting dilated pupils persist, with mental depression and insomnia. Morbid hunger has also been described in this connexion.

Pathological material confirms the involvement of the brain and meninges in the acute type of case, but no reference is made to the spinal cord or peripheral nerves. Glaser (1928) described oedema of the brain with no further details. Goldschmid and Kuhn (1920) found degeneration of the cortical cells in the cerebral hemispheres, particularly in the frontal and parietal lobes. Duvoir (*et al.*, 1937; 1939) found acute congestion of the brain and meninges, and Heiman decrease of cerebrospinal fluid, swelling of the brain with engorgement of pial veins but no haemorrhages. The brain was found to contain traces of methyl alcohol and formaldehyde, and 62 mg. of bromide in 300 g. of tissue. No microscopical findings were mentioned.

The neurological effects of long-continued exposure to small concentrations have been described in only a few cases, and no pathological material is available, except from animal experiments. Jacquet reported one case with vertigo, weakness, transient diplopia, and dimness of vision following brief exposure, and two cases with similar symptoms plus flaccid paralysis of legs and psychic disturbances after more prolonged exposure to an unknown concentration. Watrous (1942) much later reported the occurrence of mild nervous symptoms in 31 persons out of a total of 90 working in a filling-and-sealing room of a methyl bromide plant who were exposed for two weeks to a concentration of less than 35 parts per million. Headache, vertigo, difficulty in focusing, muscular pains, and dimness of vision were the symptoms observed, but no neurological abnormalities were found on examination. All these cleared up within two months of the patients' leaving the factory.

De Jong (1944) has reported three cases with involvement of the central nervous system who were exposed to methyl bromide vapour in low concentrations for an average of six months before the onset of symptoms. One of these patients had a history of numerous methyl bromide blisters on his hands. The nervous symptoms in these cases were vertigo, unsteadiness, and blurring of vision, followed by diplopia, weakness, numbness and tingling of the legs, unsteadiness of gait, and headache. No psychic disturbances were observed. Examination revealed moderate dilatation of pupils, with nystagmus, some cerebellar ataxia of the limbs, exaggerated tendon reflexes, and Babinski response in one case; diminished ankle-jerks in one case; normal abdominal and cremasteric reflexes, diminution of vibration sense in both legs with retention of position sense, and diminished cutaneous sense peripherally in one case. The C.S.F. was normal in each instance, not under pressure. Blood counts were normal, and the blood bromide varied from 9 to 13 mg. of sodium bromide per 100 c.cm. All the cases showed slow improvement over three months.

Meadows, in a personal communication, describes the case of a man aged 36—employed in the same factory as the case described below in detail—who had been working with methyl bromide for two months when he began to complain of numbness and tingling of both hands and feet, with progressive weakness of the legs. This was followed by dimness of vision, with inability to read near print, and peeling of hands and feet. On examination visual acuity was impaired, nystagmus was present, there was weakness but no wasting in the lower limbs, ankle-jerks were absent, and the left plantar was doubtful. There was loss of vibration sense in the legs, but no other sensory changes. Investigations of blood and C.S.F. were negative. The condition improved, and the patient was advised not to return to his work. It is only fair to add that at this time (a year ago) investigations failed to reveal any leakage of the gas, but when a further case arose a leakage was found in the department dealing with insufficiently filled extinguishers requiring reopening and refilling.

Animal experiments are interesting. Four investigators in America exposed rats, rabbits, guinea-pigs, and monkeys to varying concentrations of methyl bromide for varying times. The particular experiments appertaining to chronic neurological effects showed that 14 to 46 exposures to concentrations of 0.25 mg. per litre produced paralysis of a flaccid type in the limbs of rabbits and rhesus monkeys: these paralysees were bilateral, and affected equally the front and hind legs of the rabbits but chiefly the legs of the monkeys. On discontinuing the exposures when these paralysees had developed, the monkeys showed loss of equilibrium with a general ataxia of movement suggesting to the experimenters a cerebellar involvement. The reflexes became normal when the function of the limbs returned. Exposures to concentrations of 0.13 mg. per litre produced paralysees in rabbits, but had no effect on monkeys. Careful histological examination of the brain and spinal cord in the affected animals showed no abnormalities, but no mention is made of examination of the peripheral nerves.

Illustrative Case History

A male aged 17 was admitted to Staines County Hospital with the history that he had worked at a methyl-bromide-producing factory for 18 months, but had changed his type of employment a year later. During the six months before admission he had worked in the refilling department, where fire extinguishers which had been insufficiently filled were reopened and refilled with methyl bromide until they had attained the standard weight. He and the other employees in this department realized that a certain amount of gas escaped, but there appeared usually to be sufficient extraction by ventilation. The more efficient the workman the less the escape of gas. The patient was always rather clumsy, and the employees remarked on the increase of escaping gas in his work. On questioning, it was stated that the gas had a peculiar stale odour, "as though the air had not been changed for a fortnight" (this is contrary to the general belief that the gas is odourless).

Two months before admission the patient began to notice weakness of his legs, interfering with his ability to jump on chairs, buses, etc.; at the same time he felt his toes tingling and found difficulty in forming his words. His mates remarked that "he was more stupid than usual," and he developed a methyl bromide dermatitis and burns on his hands. At no time did he complain

of his symptoms to the factory doctor, who examined the employees carefully each week. One month later he found that his hands were clumsy and that he was much slower in his work and unable to run owing to weakness of the legs. He had difficulty in going up and down stairs, and began to notice he was seeing double. There were no further pains or any dysaesthesiae. Three weeks later—a week before admission—he was unable to read or see things close to him, and his speech was becoming worse, although his unsteadiness of hands was stationary. On the day of admission he was hardly able to walk to work; he complained of constant double vision, giddiness, and weakness, and when he got to his job he fell down and was quite unable to get up. He said his head ached and "he couldn't see his feet." Many of his symptoms had been noticed by his fellow employees, although they had experienced none of them, and these were ascribed to his unhappy home life with a cruel stepfather. On inquiry the home life of the boy was found to be unsatisfactory, but no overt cruelty seemed likely and his diet appeared to be adequate. He had no history of sore throat.

On admission he was mildly confused, with a considerable slurring ataxic dysarthria, and was unable to stand or walk, owing to weakness of his legs. His confusion was chiefly a disorientation in time, and it was impossible to elicit any chronological history from him. He was somewhat dull and retarded, but no other psychic abnormalities were found. Pulse and temperature were normal, and the only abnormal signs were found in the nervous system. His pupils were dilated, with good reaction to light but less to convergence-accommodation: he was unable to read, but had 6/9 vision in both eyes at 6 metres. Visual fields were full to confrontation, and his disks were normal. He had lateral nystagmus, and a fine rotatory nystagmus in both eyes on looking up, with diplopia at extremes of lateral movement. The upper limbs were of normal tone, without weakness, and showed diminished tendon reflexes. Slight motor incoordination was present and he was unable to perform rapid repetitive movements. There was no sensory loss. The abdominal reflexes were greatly increased and cremasterics normal. In the lower limbs undue flaccidity with marked motor weakness at all joints was observed, and the knee-jerks were just present, although the ankle-jerks could not be obtained. Motor ataxia in both legs was very obvious, and the left plantar response was Babinski. Vibration sense was absent in both legs up to the knees, and there was poor appreciation of position on the right. No superficial sensory changes were found. There was precipitancy of micturition. Lumbar puncture was performed and a clear freely communicating fluid obtained at a pressure of 100 mm. of water.

Two days after admission he began to be able to read and could move his legs about in the bed without trouble; his hands had lost their ataxia. A week after admission he showed nystagmus, ataxic dysarthria, and a broad-based staggering gait, with generalized reduction of tendon reflexes and diminished vibration sense in both legs. His plantars were now flexor, and he could focus normally without diplopia. The picture was very reminiscent of hydantoinate (epanutin) intoxication. The patient's condition slowly improved and he was fit for physiotherapeutic re-educational exercises a month after admission, by which time he had a slightly slurred speech, some difficulty in turning and walking fast, and nystagmus. All other abnormalities had cleared.

Seven weeks after admission he was discharged with no residual sequelae, and was advised not to return to employment where methyl bromide was used.

Investigations.—Cerebrospinal fluid:—Protein, 40 mg. per 100 c.cm.; 4 lymphocytes per c.mm.; chlorides, 640 mg. per 100 c.cm.; bromide (sodium), 27 mg. per 100 c.cm.; W.R. negative; Lange gold curve, 0000000000. Blood count:—Red cells, 4,850,000; haemoglobin, 90%; white cells, 9,000—polymorphs 70%, lymphocytes 26%, monocytes 2%, eosinophils 2%. The urine showed no abnormality.

Discussion

In exactly half the cases of methyl bromide poisoning described in the literature (39 out of 78) subacute or chronic effects on the nervous system have appeared, and in addition animal experiments confirm this type of damage from prolonged exposure to small concentrations of the gas. The neurological signs divide themselves into central and peripheral effects, and the latter seem more important in the chronic cases. These appear to be the result of a toxic peripheral neuritis with numbness, tinglings, flaccid pareses, loss of reflexes, and few sensory changes except loss to vibration. In some cases there was muscular wasting with complete paralysis, although no increase in protein or cells was found at any time in the cerebrospinal fluid.

The central effects described comprise headache, transient diplopia, difficulty in accommodation, dimness of vision, dysarthria, generalized incoordination, and vertigo. In four cases only, with prolonged exposure, psychic disturbances appeared,

showing retardation of thought, drowsiness, mild disorientation, and slow response to questioning. Objective signs were present in 10 cases of this type, and included nystagmus, pupillary dilatation with sluggish reflexes to light, diminution of vision, cerebellar ataxia and dysarthria, and loss of vibration and position sense. Centrally, therefore, the main incidence seems to be on the optic nerves, oculomotor nerves, cerebellar connexions, and posterior columns, with occasional involvement of the higher centres producing psychic disturbances.

Treatment up to the present has been only symptomatic, as these cases improve with rest in bed and removal from exposure to the gas, and no great increase in blood or cerebrospinal fluid bromide has been found (up to 27 mg. of sodium bromide per 100 c.cm.). It is presumably the methyl combination that does the damage, as bromide poisoning is not clinically apparent until blood concentrations of above 100 mg. per 100 c.cm. are reached.

Convalescence is prolonged, with persistence of some ataxia, undue fatigue, and nystagmus long after the disappearance of the peripheral neuritis, but most of the cases seem to have recovered within a year. Two patients were returned to employment where the risk of exposure to methyl bromide was still present, and both relapsed within six months and had lost time from work owing to residual and recrudescing symptoms.

Summary

The effects of chronic methyl bromide poisoning on the nervous system are described, with a review of the literature and an illustrative case history.

These effects, on analysis, form a pattern of dysfunction mainly affecting the peripheral and optic nerves, cerebellar connexions, and certain spinal cord tracts, although the material available is not sufficient to postulate a definite syndrome associated with methyl bromide poisoning.

On removal of the patient from the toxic hazard the symptoms slowly clear up without any specific treatment and apparently without permanent sequelae, but it is inadvisable to allow return to further exposure.

BIBLIOGRAPHY

- Barbour, R. F., Pilkington, F., and Sargent, W. (1936). *British Medical Journal*, **2**, 957.
De Jong, R. N. (1944). *J. Amer. med. Ass.*, **125**, 702.
Duvoir, M., Fabre, R., and Layani, F. (1937). *Bull. Mém. Soc. méd. Hôp. Paris*, **53**, 1540.
— (1939). *Bull. Sci. pharmacol.*, **14**, 15.
Floret (1915). *Zbl. Gewerbehyg.*, **3**, 146.
Frieman (1937). *Samml. Vergift.*, **8**, 31.
Glaister, J. (1942). *Medical Jurisprudence and Toxicology*, Livingstone, Edinburgh.
Glaser, E. (1928). *Dtsch. Z. ges. gerichtl. Med.*, **12**, 470.
Goldschmid, E., and Kuhn, E. (1920). *Zbl. Gewerbehyg.*, **8**, 28.
Heiman, H. (1944). *Industr. Hyg. Bull.*, **23**, 3.
Irish, D. D., Adams, E. M., Spencer, H. C., and Rowe, V. K. (1940). *J. industr. Hyg.*, **22**, 218.
Schifferli, E. (1942). *Rev. méd. Suisse rom.*, **62**, 244.
Watrous, R. M. (1942). *Industr. Med.*, **2**, 575.

HYPOPROTEINAEMIA DURING RECOVERY FROM SEVERE ANAEMIA

BY

TREVOR DAVIES, M.R.C.S., L.R.C.P.

Resident Medical Officer, Wrexham and East Denbighshire War Memorial Hospital

Generalized oedema due to diminution of the serum proteins is not a common complication of the treatment of anaemia and does not occur unless the ingestion of protein is low during the period of regeneration of haemoglobin. It is not generally recognized that the formation of haemoglobin makes great demands upon the protein reserves of the body. Heath and Taylor (1936) state that for a gain of each 10% of haemoglobin about 80 g. of protein is required and that haemoglobin formation is accompanied by a positive nitrogen balance only when the diet contains more than 6.2 g. of nitrogen a day. They say, however, that haemoglobin formation takes place at a normal rate even in the face of a negative nitrogen balance, and that in these circumstances the demand for nitrogen for haemoglobin formation appears to be supplied by tissue and plasma proteins.

The case described below is one of severe nutritional oedema following the treatment of pernicious anaemia. Faulty dietetic

habits had been present for years, and they were not fully corrected during the period in hospital preceding the onset of the oedema.

Case Report

The patient, a married woman aged 49, was admitted to hospital on June 13, 1944. She complained of increasing lassitude and loss of weight over a period of a few years. She stated that she had lost about 4 st. in the past 18 months. She had suffered from dyspepsia most of her life, and often vomited if her food "disagreed" with her. Her appetite had become very poor, and she existed mainly on a soft carbohydrate diet, rarely eating meat. She had had occasional diarrhoea during the past four years. The climacteric occurred early in 1944. There was no history of any previous severe illness.

On examination the patient was very emaciated and pale. The skin had a faint yellowish tint, but she had a malar flush. Her hair was thin and lustreless, and her general appearance suggested a mild degree of pituitary cachexia rather than pernicious anaemia. The tongue showed slight atrophic glossitis, but there was no koilonychia. She had a tachycardia of 100 a minute, but the pulse was regular and of good volume. B.P. 120/75. The heart was slightly enlarged, but there were no murmurs. Examination of the chest revealed no abnormality, and there were no abnormal neurological signs.

A blood count revealed a macrocytic anaemia. Red cells, 1,600,000 per c.mm.; Hb, 35%; C.I., 1.13; white cells, 7,500 per c.mm. A fractional test meal showed complete achlorhydria. A glucose-tolerance test gave normal figures, and the Wassermann reaction was negative. Radiological examination of the alimentary tract by barium meal and enema revealed no abnormality.

Treatment with liver extract, ferrous sulphate, and vitamin B complex was begun, and the haemoglobin and red cells increased steadily, with a maximum reticulocyte response of 10%. On July 7 a painful thrombophlebitis occurred in the left saphenous vein, and a week later a similar condition occurred in the right saphenous, both accompanied by oedema of the leg. The inflammation and oedema subsided quickly, however. On Aug. 9 the haemoglobin was 76% and the red cells 3,700,000 per c.mm. The patient felt very much better and lost her feeling of lassitude, but she could not be induced to eat the full hospital diet. She constantly refused meat, but ate moderate quantities of carbohydrate foods. On Aug. 11 gross oedema of the legs, abdominal wall, and trunk appeared, almost overnight. Ascites was present; but the oedema of the trunk finished abruptly at the mid-dorsal level—none was seen in the arms or face. There was no pain or pyrexia and the urine was normal. No pleural effusions were present, but moist sounds were heard at both lung bases. The urinary output was small (15 oz. in 24 hours). The serum proteins were low: total protein, 4.16 g. per 100 c.cm. (albumin 1.8%, globulin 2.36%); serum sodium, 346 mg. per 100 c.cm.

On Aug. 16, 80 g. of protein in the form of reconstituted plasma was given intravenously by slow drip and the patient was persuaded to eat a high-protein diet. Almost immediately a marked diuresis resulted, the urinary output often exceeding 80 oz. a day during the next three weeks, during which time the oedema almost completely disappeared. On Aug. 20 the total serum proteins had risen to 5.25%, the diuresis was still being maintained, and the oedema was steadily diminishing. On Sept. 8 the patient was out of bed and felt quite well. Very slight oedema of the feet was present, but this was probably attributable to the thrombophlebitis. The serum proteins at this date were: total protein, 5.61% (albumin 3.36%, globulin 2.25%). The liver treatment was maintained and the haemoglobin had risen to 80%. She was discharged from hospital on Sept. 23.

Comment

The above case illustrates the importance, during the treatment of severe anaemias, of maintaining a diet rich in protein so as to provide the excess of protein necessary for haemoglobin regeneration. If protein intake is inadequate the depletion of the reserves of stored protein may not become evident until a considerable amount of haemoglobin has been formed. As stated before, in order to prevent this depletion a daily intake of more than 6.2 g. of nitrogen available for protein formation is necessary. Inadequate intake of nitrogen owing to a psychological distaste for food appears to have been the main factor in the causation of the protein deficiency in this case, but it is interesting to speculate that the fatty degeneration of the liver which occurs in pernicious anaemia may have impaired the hepatic synthesis of tissue protein and that the hypoproteinaemia may to some extent have been "hepatic" as well as "prehepatic."

The treatment of clinically severe states of hypoproteinaemia with gross oedema is a matter of some urgency. When the deficiency is of the "prehepatic" type, due to inadequate