RESUSCITATION METHODS FOR RESCUE BOATS

BY

F. C. EVE, M.D., F.R.C.P.

Resuscitation is in the melting-pot. We have lost our old confidence that if Schäfer's ritual is faithfully performed nothing better can be done. We now realize that resuscitation is a trinity—of ventilation of the lungs, warmth, and restoring circulation to the nerve cells of the brain, of which the last is often the most important. For the circulation, Silvester's method appears to be better than Schäfer's, and the rocking method (Eve, 1932) much the best on present evidence (Hemingway and Neil, 1944). Ventilation is adequate by the older methods in mild cases, but not in bad cases when chill, delay, and asphyxia have nearly killed the vital nerve cells. Then the diaphragm loses tone, stretches, and is pulled up by the elastic lungs into a contracted condition of extreme expiration. At that stage Schäfer's method is impotent: no air will enter or leave the lungs, which obviously need expansion, not compression. But the rocking method, in which the weight of the abdominal contents pushes and pulls the diaphragm up and down like a piston, still acts perfectly—being independent of muscular tone. These considerations led the Royal Navy (in which the results of the Schäfer method had been very disappointing) to adopt the rocking method when possible (Gibbens, 1942).

This, very briefly, is the present position, which rescuers should know. Evidently it is in the rescue-boat that lives can be won or lost: the victim may be too far gone for resuscitation by any method by the time that ship or shore is reached. This problem has never been systematically studied, and it is different for each type of boat. The men who will man the boat in an emergency must try out *beforehand* the method best suited to their boat and submit their plans to an appropriate doctor. For instance, the seats may be too short or crowded to permit Schäfer's method; the thwarts may be too deep or the floorboards too wet for the older manual methods. I am told that in a speed-boat the thumping on the waves makes Schäfer impossible, and hence the rocking method is being substituted by the R.A.F. Cmdr. Michelmores took me a trip in the smaller type of National Lifeboat. It was so cluttered with necessary gear that we agreed that to make a flat place with life-jackets for Schäfer's method was the only solution.

I now submit two methods which seem worthy of trial under actual rescue conditions: (1) manual rocking (Eve, 1943); (2) the Lap-Silvester method (Eve, 1944).

(1) The Manual Rocking Method, as applied in a Rescue-boat (Figs. 1-3)

Two men stand facing, gripping each other's hands (or, better, a bandage twisted into a small ring). The victim lies face downwards across the two pairs of linked forearms, which support his thighs and upper chest. His arms and head dangle limply. It is preferable for each bearer to place his forearm under both thighs

FIG. 1

FIG. 2

and grip the trousers, as photographed. The two men, by swaying from their hips, rock the casualty through about 45° each way, starting head down and pausing till all water has drained out before inducing inspiration by the feet-down tilt. This rocking is continued 10 or 12 times a minute till the patient breathes naturally, when he can be laid down, prone at first. This was done to me in

a very large ship's lifeboat. I did not have to breathe, so the ventilation must have been adequate. The stalwart pair said they could have rocked my 10½ st. easily for a quarter of an hour at least. A heavy casualty would be unsuitable: a child can be rocked by one person. I believe that with their legs astride a thwart the two operators (now linked into a quadruped) will have considerable stability. In a sea-way they could sit down if preferred (Fig. 3).

FIG. 3

FIG. 4

A Child Revived by Manual Rocking.—A little girl of 4 was immersed for some 5 minutes in the pit of a stationary mill-wheel near Truro last Easter. Pte. Jack Collings descended, and, also slipped into the water. At once he performed manual rocking through quite 45° a dozen times a minute. In the confined space this was the only prompt method. Respirations returned in 10 or 15 minutes, and recovery was complete. The first head-down tilt emptied at least half a teacup of water through the mouth. Fortunately, in his first-aid class, the operator had himself been rocked on the usual rocking stretcher. So that the credit of this first resuscitation by *manual* rocking must be shared with his instructor, Supt. Schoon, who supplied me with these particulars.

(2) The Lap-Silvester Method (Fig. 4)

This idea was originally suggested to me by the ladies of the Tunbridge Wells Red Cross (Dr. Taylor's class). The casualty lies face upwards on the laps of two bearers sitting on a thwart (oilskin trousers donned on the outward trip). It is better that they should face opposite ways so that the victim is oblique to the thwart. The operator stands astride the thwart and uses Silvester's method, grasping the forearms and alternately extending them horizontally, and compressing the chest with them, firmly but not violently. (There is no room for the orthodox outward sweep of the arms, which I fancy, after personal trial, is redundant.) The thigh of the first sitter gives room for the head to dangle right back, a posture in which one throat specialist assures me the tongue will not block the glottis. This is confirmed by the radiograph of the airway of a young adult in this posture, kindly taken by Dr. Bannen. The first sitter flexes the neck during expiration so that the brain is alternately congested and emptied of blood (important for its circulation). As necks vary, he should also watch the tongue in inspiration. The second sitter during expiration leans his weight forward so that his forearm compresses the abdomen (more effectual than the hand). This pushes up the diaphragm and also propels blood from the abdomen to the heart. This method is available if the victim is too heavy or the rescue-trip too long for manual rocking.

Dr. R. R. Macintosh, Nuffield Professor of Anaesthetics, investigated this method at Oxford and kindly allows me to quote the unpublished results. Mr. J. Roberts, a healthy athlete, submitted altruistically to anaesthesia with apnoea caused by hyperpnoea, in which the state of drowning was simulated. The ventilations produced by each rock (ten a minute) through 45° each way on a stretcher were: Schäfer, 530 c.cm.; Silvester, 650; rocking, 850 c.cm. These figures confirm the previous (Macintosh, 1943) historic experiment on Squad. Ldr. A. E. Pask, which are proportionally similar (340, 400, and 580 c.cm.), and were done by the same team of experts. Rocking Mr. Roberts face upwards was less effective than prone (570 c.cm. at 30°; 635 at 45°). If the piston action of the diaphragm was impeded by a pillow under the abdomen, rocking face downwards was only half as efficient (325 c.cm. con 725 at 30°). An interesting novelty was the excellent ventilation produced (630 and 670 c.cm.) by merely lifting the pelvis or coat when prone, thus allowing the abdomen to sag and pull down the diaphragm. (This should be useful to remember when doing the Nielsen modification of Schäfer.)

The *Lap-Silvester*, done on two laps with straightish spine (too much bent in Fig. 4), gave the adequate ventilation of 530 c.cm. improved to 600 c.cm. by abdominal pressure (cf. Schäfer, 530 c.cm.). With a Beverley team, I saw that the Lap-Silvester method is practicable in a 6-oared ship's lifeboat of 6-ft. beam (Fig. 4). Its merits are: warmth and dryness on laps; ventilation proved adequate; the dangling head should obviate traction on tongue or jaw; adequate abdominal compression for diaphragm and circulation; alternate congestion and emptying of cerebral blood vessels (probably important). Hence this method seems worthy

of trial in a rescue-boat, but neck or spine must never be forced backwards.

Summary of Conclusions

1. Resuscitation must be started *immediately* the victim is hauled aboard over the lee stern quarter: ship or shore may be too distant for revival. Water must first be drained out by a steep head-down tilt (Fig. 1).

2. If space allows, Schäfer may then be used provided that air can be felt or heard going in and out of the mouth.

3. If this is not satisfactory, or resilience feels absent in the lungs, do not persist in the Schäfer ritual, but change at once to the manual rocking method, especially in a child or light subject. Personally, I would start with this method, after draining out water.

4. For heavy subjects or longer boat-journeys, the Lap-Silvester method is suggested as being warmer and providing more circulatory aid to the brain and heart than plain Silvester or Schäfer.

5. Rescue-boats differ greatly; each type is a separate problem in methods of resuscitation, which need to be tried out *beforehand* by rescue teams who dwell beside deep waters.

Methods must be judged by records of successes and failures, which I should be glad to collect and collate until there is some official channel.

REFERENCES

- Eve, F. C. (1932). *Lancet*, 2, 995.
 — (1943). *British Medical Journal*, 1, 535.
 — (1944). *J. Amer. med. Ass.*, 124, 964.
 Gibbens, G. H. (1942). *British Medical Journal*, 2, 751.
 Hemingway, A., and Neil, E. (1944). *Ibid.*, 1, 833.
 Killick, E. M., and Eve, F. C. (1933). *Lancet*, 2, 740.
 Macintosh, R. R. (1943). *British Medical Journal*, 2, 493.

A BLAND-SUTTON MEMORIAL

At the instance of Sir Alfred Webb-Johnson, President of the Royal College of Surgeons, we are able to reproduce below a photograph of a mural tablet in memory of Sir John and Lady Bland-Sutton.

A Faculty has been granted by the diocesan authorities of Canterbury for the erection of this tablet in the Parish Church at Ringwould, Kent, where Sir John and Lady Bland-Sutton regularly worshipped when they were in residence at Villa Vita, Kingsdown, Deal. The memorial was designed by Ainer W. Hall, M.C., F.R.I.B.A.

KING EDWARD'S HOSPITAL FUND

The Speaker of the House of Commons presided, in the absence of the Duke of Gloucester, at the meeting on Dec. 11 of the general council of the King Edward's Hospital Fund for London. He referred in his opening address to the discussions which representatives of the Fund, "with their impartial and at the same time intimate knowledge of so many hospitals," had had with the Minister of Health, and to the help this would mean to the Minister in framing his proposals. Work on hospital diet was, he added, very much to the fore, and Sir Jack Drummond and his committee were now preparing a second memorandum offering more detailed advice based upon the experience of surveys at a good many hospitals.

Sir Edward Peacock as treasurer reported that in view of the heavy expenditure which would confront the hospitals after the war it had been thought wise to continue the slight reduction in the annual distribution, which was again £280,000. Subscriptions and donations, estimated at £38,000, were about the same as last year. Income from investments, at £213,000, showed an increase of £10,000 over the previous year, but this included certain arrears and other non-recurring items. These figures did not include the £300,000 received from the Nuffield Trust for the special areas and the transfer from this amount of £250,000 to Guy's Hospital.

Speaking of plans for a national health service, Sir Edward said that during the summer a statement of principles was prepared in close concert with the Voluntary Hospitals Committee and the British Hospitals Association, which underlined the meaning of partnership between voluntary and municipal hospitals, laid stress upon the need for a Central Hospitals Board and on regional and local machinery, dealt with the financial issues raised by the White Paper, and pointed out the importance of making it quite clear that payments from public moneys would be payment for services and not subsidies. The proposal of the White Paper, he continued, that all hospital services should be free of necessity cut at the roots of the contributory schemes which had become such a valuable feature of the life of this country. Everything must be done to preserve as large a part of the voluntary support of hospitals through such schemes as was possible. It would be almost criminal if, in our endeavour to secure another valuable thing, we should destroy that unique organized contribution of millions of pounds by millions of people to their hospitals.

The Public Instinct about Hospitals

Sir Alfred Webb-Johnson, in proposing a vote of thanks to the Speaker, said that the instinct of the people had been sound in regard to the type of hospital which the people had built for themselves—the great general and special hospitals, which acted as centres for consultation; the intermediate type of hospital, which served the immediate needs of a populous area; and the home or cottage hospital, which met the needs of the general practitioners in the smaller communities. This instinct had also been sound on the distribution of hospitals. The great consultant hospitals were based on university centres and their influence spread over large areas. People went to these consultation centres not because of any geographical arrangement, but because they served natural "catchment areas." National or county boundaries did not count; when a man was sick and his people were anxious about him they wanted to go where facilities were best and where the advice they sought was accessible and available. In the management of the voluntary hospitals the instincts of the people had again been sound. They had given absolute freedom to the professional men who work in the hospitals, had sought their advice in their corporate capacity through the medical committees, and had sought their aid in the management of the hospitals.

But in spite of all this there was need for more co-ordination and expansion, and it was to be hoped that the Government would follow the example of the King's Fund, and would guide and not control, would help the voluntary hospitals and not hinder them. It was important that the contributions under the National Health Insurance scheme, which were not really enough to meet the general practitioner service, should be devoted entirely to that service, and that the hospitals should be maintained by rates, taxes, and voluntary contributions. This would do something to preserve the voluntary contributions, because insured persons would not be told that they had already subscribed directly to the voluntary hospitals.

The people of this country must retain the advantages which they had gained for themselves, and the way to do this was for experts in hospital matters to have a responsible part in the planning and administration of the hospital services. Those experts—the representatives of the voluntary hospitals and representatives of the medical profession—must have their place in the council chamber.

In conclusion Sir Alfred spoke as follows: "When we talk about the freedom of the profession we are not talking about some vested interest. We are talking about one of the fundamental liberties of the subject—his right, within a free community, to obtain free and