problem of psychopathy and conduct disorder in women are noted, and an attempt is made to show the relation of this material to the general problem of the war neuroses.

We wish to thank Air Vice-Marshal C. P. Symonds, C.B., and Air Commodore R. D. Gillespie for encouragement and criticism in the preparation of this paper.

BIBLIOGRAPHY

- Ballard, S. I., and Miller, H. G. (1944). *British Medical Journal*, 2, 40.
 Fairbairn, W. R. D. (1943). *Ibid.*, 1, 183.
 Gillespie, R. D. (1942). *Psychological Effects of War on Citizen and Soldier*, New York and London.
 Henderson, D. K. (1939). *Psychopathic States*, London.

LUDWIG'S ANGINA

BY

JOHN FARR, F.R.C.S.Ed.

Major, R.A.M.C.

AND

E. D. STANHOPE, L.D.S.R.C.S.

Major, A.D. Corps

Ludwig's angina, if not promptly recognized as such, is often fatal. This symptom-complex of a floating tongue with submandibular cellulitis is not uncommon among East African native troops, and during the last 30 months we have had to treat 20 such cases. It is the purpose of this paper to direct attention once again to the cardinal signs which herald the onset of the syndrome and to outline a system of management which has proved satisfactory and, in our experience, has appreciably reduced the mortality.

The higher incidence among native troops may be due to several factors. Kekwick and Wright (1945) have recently demonstrated that low-grade vitamin C deficiency is prevalent, and it is possible that this leads to a decreased resistance to oral sepsis. The African also has a well-recognized liability to coccal infections, and submandibular cellulitis may be just another manifestation of this tendency. Further, the native shows a hesitancy in seeking dental treatment until he suffers from marked disability, and in some cases this neglect has initiated the condition.

Where the focus of infection in Ludwig's angina is dental the third lower molar is most commonly the tooth involved, for the following reasons. The inner alveolar plate is thinner than the outer, thus facilitating the spread of infection deep to the cervical fascia and myelohyoid muscles. Again, despite the fact that the African is more prognathous than the European, there is sometimes insufficient room for the complete eruption of the third lower molar. This partially erupted tooth is consequently more prone to pericoronal infection. Alternatively, the infection may be periapical in origin, arising from any of the lower teeth.

An intermediate stage in the pathology is an acute alveolar abscess in the case of periapical infection or a pericoronal abscess in the case of a partially erupted third lower molar. It is imperative to recognize the differential diagnosis between an uncomplicated alveolar or pericoronal abscess and one in which the sepsis has spread into the deep structures of the floor of the mouth, for in the latter stage intraoral drainage by extraction of a tooth or removal of a gum-flap will not provide adequate drainage once the pus has tracked forward in the slanting trough of the myelohyoids, and prompt external drainage is essential. For this reason, where such a spread is a possibility, it is important that the patient should be seen by both a surgeon and a dental surgeon, and that subsequent treatment should be carried out in the closest collaboration.

The cardinal signs (Fig. 1) are:

1. *A Floating Tongue*.—An inflammatory process with or without formation of pus in the floor of the mouth, firmly embraced by the myelohyoids and deep cervical fascia, causes the tongue to be elevated, and in severe cases extruded. Intraoral oedema is pronounced, and in some cases reaches up to the necks of the teeth. Dyspnoea and dysphagia may result from oedema or from pressure on the trachea and oesophagus.

2. *Submandibular Cellulitis*.—This is a diffuse and tender swelling beneath the jaw which may cross the midline of the neck. It must be emphasized that it is extremely dangerous to withhold treatment while awaiting fluctuation, for the patient may succumb to toxæmia

or respiratory obstruction before pus is formed, and it is essential that pressure on the deep structures be relieved by sectioning the deep cervical fascia and one or both myelohyoid muscles. Others have reiterated that when the tongue is elevated and there are signs of a submandibular cellulitis the patient is suffering from Ludwig's angina. The diagnosis of the lesion is the indication for operative treatment. Delay is dangerous, and in our opinion surgical intervention is urgent when these two signs are manifest.

FIG. 1

Surgical Preparation

It is of paramount importance that a tracheotomy set should be available in the theatre, and should accompany the patient on his return to the ward. It must remain there for 24 hours, or longer if indicated. This simple precaution is not superfluous, and must never be omitted from the surgical routine. The choice of anaesthesia is determined by the facilities available. We have found local analgesia to be unsatisfactory because of its inability to allow adequate dissection. Ethyl chloride is hazardous in the extreme because of the tendency to spasm of the laryngeal muscles. Where full hospital facilities are at hand the safest method of anaesthesia is by endotracheal nitrous oxide and oxygen, to which a small quantity of ether may be added. Where such facilities are not at hand we have found light chloroform analgesia to be the safest method, preceded by adequate premedication, but, as in any form of anaesthesia, the most important factor remains the skill and experience of the anaesthetist.

Operation

A curved incision is made about one inch below the mandible, extending from a point below the third molar to the midline, or across the midline if indicated. This is deepened to the level of the myelohyoid, which is sectioned. Frequently pus has spilled over the posterior border of this muscle and is found superficial to it, but provided the origin of the inflammatory process is not in the submental glands, section of the

FIG. 2

muscle is indicated. A drain is introduced, but instead of merely draining the space it is passed on through the floor of the mouth, emerging between the lingual border of the mandible and the tongue (Fry *et al.*, 1942). The part of the tube that is lying in the submandibular structures is fenestrated (Fig. 2).

The proximal end of the tube is brought out through the mouth and fixed with adhesive plaster to the side of the cheek. The distal end is anchored to the skin at the site of the incision. Vaseline packing is gently tucked into the incision, and the tube is occluded by a clamp at its distal end. If the condition has had a dental origin the extraction of the tooth involved completes the operation.

Post-operative Care

Chemotherapy in the form of sulphonamide drugs is used as a routine auxiliary treatment. Gas-forming organisms often predominate in the infection, and in at least two of our cases gas under tension has escaped on incising the deep cervical fascia. We have therefore used hydrogen peroxide 1/10 as an irrigant. It seems likely that zinc peroxide suspension would be more efficacious, but its availability is limited. During the patient's waking hours there is a continuous slow drip of irrigating solution introduced through the proximal end of the tube. This escapes from the openings in the tube and allows the region to be gently irrigated throughout its various layers (Shepherd, 1940). The fluid coming out of the incision around the sides of the tube presents some difficulties, as it tends to drip down the side of the neck, but with diligent and zealous nursing care it may be collected in a kidney dish hung on the patient's neck, or a close-fitting jaconet collar.

When continuous irrigation has not been convenient we have compromised with 100-c.cm. irrigations every four hours, after which the clamp is removed and the tube functions as an ordinary through-and-through drain. After 48 hours the general and local improvement is sufficient to consider shortening the tube. It is cut away flush with the buccal mucosa and shortened about half an inch. If further irrigations are indicated at this stage, they are most easily carried out by introducing a small catheter beside the drainage-tube and syringing hydrogen peroxide through it. The drainage-tube is removed on the fourth to sixth post-operative day, and a vaseline dressing, changed when necessary, completes the post-operative dressings.

Although all these incisions have healed by slow granulation and subsequent epithelization, a linear scar results, and no cases have required excision for cosmetic reasons.

Summary

Possible reasons are outlined for the relatively high incidence of Ludwig's angina among East African native troops.

The diagnostic signs that make surgical intervention imperative are stated.

A system of management is described effecting free drainage and submandibular decompression followed by a drip irrigation.

We wish to record our thanks to Brig. R. P. Cormack, O.B.E., D.D.M.S., East Africa Command, and Col. D. Bell, O.B.E., E.A.A.M.C., for permission to publish this paper; and to acknowledge the co-operation of S/Sgt. F. Finch, R.A.M.C., for the production of the accompanying diagrams.

REFERENCES

- Fry, W. Kelsey, Shepherd, P. R., McLeod, A. C., and Parfitt, G. J. (1942). *The Dental Treatment of Maxillo-Facial Injuries*, Blackwell Scientific Publications, Oxford.
Kekwick, A., and Wright, J. (1945). (Not yet published.)
Shepherd, P. R. (1940). *Brit. dent. J.*, **69**, 430.

INGUINAL HERNIA IN THE MERCHANT NAVY AN ANALYSIS OF 200 CONSECUTIVE CASES

BY

PERCIVAL P. COLE, F.R.C.S.

E.M.S. and Senior Surgeon, Dreadnought Hospital

The incidence and cure of hernia have attracted renewed and critical attention from the point of view particularly of the associated disability in men of the fighting Forces. Detailed observation and docketed conclusions have been made possible by the concentration of large numbers of men under an organized and centralized medical service. Inguinal hernia has been made the subject of a critical review by Brig. Harold Edwards (1943), who deals with it comprehensively but also with special reference to Army statistics and military considerations. In the Army, selection of cases for operation is determined by the answer to the question whether operation will make the man a more efficient soldier. In the Merchant Navy—not

ranked as one of the fighting Services—the question is the more vital one of going to sea or not.

Classification

To emphasize particularly the age at operation, the large number of cases requiring supplementary repair, the largest portion of direct hernias, and the early age at which direct hernia occurs, the 200 cases under review have been classified numerically into the following groups:

Type of Hernia.—Oblique, 97; direct, 103.

Age at Operation.—53 cases were over 45, 38 were over 50, 19 were over 55, and 9 were over 60.

Recurrent hernias dealt with in this series numbered 22. Of these, 18 were direct and 4 were oblique. Four of these recurrences followed a previous filigree operation, and occurred thereafter: 1 at 10 months, 1 at 6 years, 1 at 12 years, 1 at 25 years.

Primary direct hernia occurring at the age of 40 and under, 27 cases. Earliest age at which direct hernia was found, 23 years.

Number of cases dealt with by filigree repair, 130. The first operation was performed on Feb. 19, 1941, and the last on June 26, 1944.

For the purpose of comparison with procedure in the recognized fighting Services a brief analysis will be based upon that embodied in an Army Medical Department Bulletin on Inguinal Hernia, issued by the War Office in Feb., 1943. Therein it is stated that over 12% of inguinal hernias recur after operation, and this recurrence rate is regarded as representative of British surgery as a whole; the factors concerned are considered to be: (1) surgical skill and experience of the operator; (2) selection of cases for operation; (3) operative detail; (4) post-operative management.

Analysis of Cases

1. In connexion with this factor all the cases have been operated upon either by me or under my direct supervision.

2. *Selection of Cases for Operation.*—Under this heading the Memorandum excludes the following categories as being unsuitable: (a) small bulges in men of inferior abdominal musculature; (b) large scrotal hernias in pot-bellied subjects of poor physique; (c) large recurrent hernias in similar subjects; (d) large hernias in men of low category because of other disorders. Among the 200 cases on which this analysis is based will be found instances included in the first three categories, and, in fact, discrimination on these lines has not been exercised, for it is a cardinal principle that no case is rejected on account of the local condition. With regard to category *d*, I am in agreement with this reason for rejection, but actually no such case has been met with. As I have previously stated (Cole, 1941), the operability rate in my hands, taking all cases into consideration, is 90% at a modest estimate.

3. *Operative Detail.*—Under this heading it is stated that 75% of recurrences after operation for indirect hernia were indirect. Authorities such as Page (1934), Block (1933), Ogilvie (1937), and Gallie and Le Mesurier (1924) have agreed that by far the greatest number of so-called recurrences after operation for oblique hernia are direct in form. It is clear that in this mass reproduction of the original condition (75%) responsibility must be attributed to factors either non-operative or operative to a minimal extent in the collected series of the quoted authorities. Prior to the series under survey, and over a period of 20 years, recurrent hernias numbering approximately 400 have been observed, and the mode of recurrence has been so often direct that this predominance has never been questioned. The small number of recurrences in the present series bears this out, for it will be observed that of the 22 recurrences included 18 were direct and only 4 oblique. Recurrences, the Memorandum states, are commonest after the Bassini operation or one of its modifications. With this my own practice records agreement, for I have not performed a Bassini operation for many years. I further agree with the recommendation that "no type of operation which interferes with the normal muscular apparatus of the inguinal canal should be practised for uncomplicated indirect hernia in the young soldier"; and in this connexion it is the age of the patient that counts and not the nature of his employment.

Apart from this simple type of hernia occurring in the young man, some supplementary method must be employed to replace