PENICILLIN IN WOUNDS OF THE CHEST

BY

A. L. D'ABREU, Ch.M., F.R.C.S.
Lieut.-Col., R.A.M.C.

AND

J. W. LITCHFIELD, B.M., M.R.C.P.
Major, R.A.M.C.

(From a British General Hospital, C.M.F.)

In the mass of literature dealing with penicillin therapy it is difficult to obtain a clear picture of what is practical and what is academic. We hope this short paper may be of value.

Since penicillin became available to this centre more than a year ago, over 500 penetrating wounds of the thorax have been treated. We believe that the therapy is best employed as follows.

General Parenteral Therapy

This is used only occasionally, and has not supplanted the well-established sulphonamide drugs, which may often be used with less distress to the patient, as they do not involve three-hourly injections. The indications are: (i) When pulmonary suppuration or severe infection exists with a pyothorax that is being surgically drained. Traumatic lung abscesses—i.e., around foreign bodies—infected missile tracks, and the rare breaking-down lung haematoma have been greatly benefited by general penicillin, which in these instances is more helpful than the sulphonamides. (ii) In severe chest-wall lesions, associated with an open sucking septic pneumothorax, which used to be a grave problem. Their closure after re-excision and removal of rib or scapula fragments (if necessary) is readily effected if combined with general penicillin, assisted in the first three days by local penicillin given by instillation tubes; no such case has failed to remain closed. (iii) For a simple or infected haemothorax general penicillin is of little or no value: it has been established (d'Abreu, Litchfield, and Scott Thomson, 1944) that local intrapleural penicillin is far more efficient as a bacteriostat for Gram-positive infection in such effusions.

Prophylactic Use of Local Penicillin, for Haemothorax

(a) *Primary Treatment (e.g., in Forward Units).*—Repeated intrapleural injection of penicillin without bacteriological and radiological check is unwise. The intrapleural injection of 60,000 units in 6 to 10 c.cm. (no more) at the end of the first two or three aspirations is advised; this injection in no way replaces adequate aspiration. As soon as the patient's general condition has been improved by these early aspirations he is usually fit for transfer to a general hospital or chest centre, where further treatment can be more satisfactorily carried out.

(b) *At General Hospital Level.*—At this centre the uninfected haemothorax is treated by aspiration without intrapleural instillations except under the following circumstances: (1) Where there is coexistent chest-wall sepsis. (2) When there is recognized or suspected pulmonary infection—e.g., when the radiograph shows a peripherally placed foreign body in the lung tissue or in the pleural cavity. Our practice is to remove such bodies as soon as the patient's general condition is satisfactory; if operation is delayed because of poor general condition and there is a coexistent haemothorax, this is treated by aspiration and intrapleural penicillin. (3) At the conclusion of a thoracotomy for foreign-body removal penicillin, 120,000 units in 10 c.cm. of saline, is instilled after air has been drawn off by an artificial pneumothorax apparatus. Unless there is infection of the pleural cavity, post-operative drainage is not done, and the resultant effusions are aspirated and bacteriologically examined as a routine. The missile bed is insufflated with penicillin powder before the lung sutures are tied.

Local Penicillin in Treatment of Infection

(a) *Intrapleural.*—Penicillin is of great value in the treatment of infected haemothorax and empyema. It is obviously only of use in cases of infection by penicillin-sensitive organisms. Intrapleural therapy is useless for Gram-negative effusions and for penicillin-resistant staphylococci; it is therefore essential to

have most careful bacteriological checks. All syringes used in this work require the most careful sterilization by boiling: no chemical method is safe enough prophylaxis; the more careful the technique the less frequent is Gram-negative infection. In an infected haemothorax or empyema containing penicillin-sensitive organisms our procedure is to aspirate the effusion and inject 120,000 units of sodium penicillin for large cavities and 60,000 units for smaller ones in 10 to 20 c.cm. of solution, on alternate days. The instillation is repeated until three specimens of the fluid have been reported to contain no pyogenic organisms on smear or culture. Frequent radiological location of the cavity is essential: even after the cavity is no longer seen radiologically its disappearance must not be assumed, and aspiration must be continued until fluid can no longer be obtained. Towards the end of this treatment the fluid becomes serous and the interval between aspirations can be lengthened to once or twice a week. Active breathing exercises are essential throughout the treatment.

Despite sterilization of the cavity, operative treatment is necessary in certain cases for the purpose of removal of clot and the prevention of excessive fibrin deposition on the pleura (parietal and visceral). The indications for surgery are as follows. (i) The presence of a sucking chest-wall wound in association with a pyothorax demands surgical drainage (closed) and penicillin closure of the wound. (ii) The presence of a large broncho-pleural fistula demands pleural drainage. (iii) The existence of a very large infected effusion associated with a total lung collapse to the mediastinum calls for closed drainage because in these cases so much fibrin is deposited that lung re-expansion and chest-wall movements are hampered; a few intrapleural instillations may be of great value before drainage, because the Gram-positive organisms may be eliminated, with a great decrease in toxicity. (iv) Considerable fibrin-clot formation, as shown radiologically or by repeated needle blockage during aspiration, is an indication for "clot clearance." If at operation a localized pocket is found and the organisms are penicillin-sensitive, complete chest-wall closure is followed by post-operative aspirations and local penicillin therapy. The larger cavities are drained for the reasons given above in (iii). (v) Persistent Gram-negative infection is a source of trouble because it leads to further exudation and fibrin deposition, thereby causing mechanical hindrance to lung re-expansion and chest-wall movements and continued toxæmia. Clot clearance and drainage are done for these cases after an adequate trial of repeated aspiration. We are using succinyl-sulphathiazole suspension for these aspirations, but this work is entirely in the experimental stage.

(b) *In Chest-wall Wounds.*—There is little to add to our previous published experience (d'Abreu, Litchfield, and Scott Thomson, 1944). Wounds that have been treated by excision and muscle closure of the pleural "suck" in the forward area arrive with the skin left deliberately unclosed: suture (delayed primary) is done soon after arrival with the assistance of penicillin-sulphathiazole powder. We excise septic chest-wall wounds, including the removal of rib and scapula fragments if necessary, and close them by suture with the aid of penicillin solution introduced by tube instillation: this is given for five days if there is underlying bony damage. The results are very successful even in the presence of frank pus. This method does not seem to be known widely enough. It is the solution of the problem of the septic open sucking wound (d'Abreu, Litchfield, and Scott Thomson, 1944).

We are indebted to Col. William C. McKinnon, the officer commanding, for permission to publish this paper.

REFERENCE

d'Abreu, A. L., Litchfield, J. W., and Thomson, Scott (1944). *Brit. J. Surg.*, **32**, 679.

C. F. Dixon and A. L. Lichtman (*Surgery*, 1945, **17**, 11) record their observations on 50 cases of congenital absence of the gall-bladder that have been recorded since 1900 and on 10 cases that have been observed at the Mayo Clinic. Symptoms resembling those of cholecystic disease were present in 58%; 48 had jaundice and 27 stones in the common bile duct. Symptoms were present in 73% of cases in which the patients were more than 45 years of age. Prolonged drainage of the common bile duct with a T-tube is advocated in the cases in which symptoms are present.