Penicillin is effective against all the common causal organisms of ophthalmia neonatorum, including the presumed virus of inclusion blennorrhoea.

Local penicillin therapy is likely to replace general sulphonamide treatment as the method of choice in ophthalmia neonatorum.

Thanks are due to Dr. E. N. Young, of the Southern Group (L.C.C.) Laboratories, for her painstaking bacteriological examinations, and to Dr. Olga Blum, Dr. Rela Bartfeld, the Matron, and Sister Condon, of White Oak Hospital, for their collaboration. I am indebted to Prof. Sir Alexander Fleming for his interest and help.

REFERENCE

Sorsby, A., and Hoffa, E. (1945). *British Medical Journal*, 1, 114.

PENICILLIN TREATMENT OF HAND INFECTIONS

BY

G. J. GROSSMARK, M.R.C.S., L.R.C.P.

AND

L. W. PLEWES, F.R.C.S.Ed.

The local use of penicillin in powder and paste (lanette wax form) in the treatment of hand and finger infections has been shown by Florey and Williams (1944) to reduce the time required to clear up sepsis, to diminish pain, and to save an average of nine man-days for each case as compared with controls. In this series 100 cases of paronychia, pulp whitlow, web-space infections, and suppurative tenosynovitis, in all groups of the population, have been treated with penicillin liquid and cream locally, with results that mainly bear out Florey's findings, but with two main differences—namely, in the results of treatment of osteomyelitis and suppurative tenosynovitis.

Principles of Treatment

In all cases in which incision was necessary general anaesthesia was used, and a blood-pressure cuff to give a bloodless field. There was no soaking of the fingers or application of heat of any kind. Penicillin was applied either in liquid form or as a cream on a fine-mesh gauze wick. The cream consisted of penicillin mixed with a base—eucerin L.M. The penicillin in this cream was found by our pathologist (Dr. Seiler) to keep its potency for a longer time than the penicillin in lanette wax. The dressings were all done in a special dressing unit; the dresser used the "no-touch technique," and modern methods of prevention of cross-infection were employed. Of the total 100 cases (representing over 500 dressings) the number of times a second organism was grown was 19. Cultures were taken at the time of operation, and sensitivity to penicillin was tested. Cultures were also taken as a routine at the change of each dressing.

Dressings were done once daily if liquid penicillin was used, or every two days with penicillin cream, since it had previously been found that, although liquid penicillin in an open wound lost its activity in about 24 hours, when cream (penicillin-eucerin L.M.) was used penicillin activity could still be demonstrated at the end of 48 hours.

Surgical procedures in all groups followed orthodox lines, except in the treatment of acute suppurative tenosynovitis. In this condition previous experience with orthodox incisions was so discouraging that a new technique was used. It was based on the observation that acute pyogenic infection of joints responded dramatically to repeated aspiration of the infected fluid and injection of penicillin liquid. None of these fingers was incised.

Results

Pulp Infections.—Thirty-six pulp infections arrived at the hospital, having had throbbing fingers for an average of seven days. In every case the organism was a sensitive *Staph. aureus* (coagulase-positive). In three-quarters of these cases the operation was the usual bilateral incision, and the remaining quarter had either the "hockey-stick" or the "fish-mouth" incision. None of these cases was treated as an in-patient, nor was any splinting done, and the average number of dressings was six. In about half the cases sloughing was present to an extent that mechanical removal was required, and here the time taken

to heal was twice as long as when sloughing was not a feature. The occurrence of sloughing seemed to bear no relation to the length of time before operation. The average number of days to "no growth" was 9, and the period off work 17. All the end-results were good.

Paronychia.—There were 32 cases in this group. In all except four the causative organism was *Staph. aureus* (the others, *Str. pyogenes*). Fourteen of these cases had pus under the nail by the time they were first seen, and enough nail was removed at operation to allow access of the penicillin cream to all the infected area. The average time before being seen at hospital was 10 days. In just over half the operations it was found necessary to open the finger on both sides. In the remainder one lateral incision sufficed. On one occasion the pus could be aspirated through a needle, followed by penicillin injection through the same needle. Dressings were done at 48-hour intervals, and on an average four dressings were required. For all this group the average number of days off work was 12; the average number of days before a sterile culture was obtained was 5. The final result was satisfactory in every case.

Web-space Infections.—This group contained 23 cases, and the average time before treatment was 11 days. The operation consisted of an incision between the distal crease of the palm distally to the beginning of the web. Fourteen of these cases were infected by a sensitive *Staph. aureus*. The average time to "no growth" was 9 days and off-work 19 days; each case required on an average seven dressings. Sloughing as a prominent feature was present only once, and chemotherapy was used on two of four patients admitted (because of lymphangitis or adenitis).

Tendon-sheath Infections.—There were nine patients who had an acute suppurative tenosynovitis affecting one of the fingers for an average of four days, the organism in six cases being *Staph. aureus*, in three *Str. pyogenes*. None of these fingers was incised; instead, pus from the sheath was aspirated opposite the proximal phalanx, and after aspiration of all that was possible 1 or 2 c.cm. of penicillin was injected through the same needle. The puncture wound was then sealed off with collodion. No splinting was done, and all of these patients were treated as in-patients; the first five cases were given sulphathiazole for five or six days. Active movements were encouraged from the first, and the relief of pain was dramatic in all but one case. This case failed to respond to treatment, and amputation was necessary because of a fulminating infection of the whole finger. In seven cases the results were perfect after one, two, three, or four punctures of the sheath (at 48-hour intervals). One patient apparently had a perfect result after four treatments (aspiration of pus and injection of liquid penicillin), but a month later had developed a slight flexion deformity of the finger. This has remained stationary, and the finger can be used. The average stay in hospital for this group was seven days, and the time off work 13 days. There was no difference in the clinical course when sulphathiazole was used, and the last four cases (all perfect results) were not given systemic chemotherapy.

Osteomyelitis.—Another group, comprising cases of osteomyelitis, were tried with local penicillin, and although one could get negative cultures in a time not appreciably longer than with the above groups, the clinical course of the disease did not seem to be affected. The results of treatment were uniformly bad. It is difficult to report accurately on this group because, first, amputation was often carried out relatively early; secondly, some of the cases were sent elsewhere for x-ray therapy; and, finally, some are still unhealed.

Comments

Early on it was realized that removal of slough helped considerably, and the results show this to be true. After trying trypsin and pyruvic acid it was felt that simple excision with a pair of sharp nail scissors was not only quicker but usually more effective.

When this research was started penicillin at a strength of 50 units per c.cm. was used for a few weeks. Subsequently the strength varied from 50 to 2,000 units per c.cm. From our analysis the strength of the penicillin employed had no effect on the result.

Many hands were swollen as a result of prolonged soaking, and others (about 30%) had had operations (mostly without anaesthesia) before attending hospital. For all groups the average time before being seen was nine days—longest in the web-infection group, and shortest in the tenosynovitis group, probably because of the severity of the pain and disability in the latter.

The most striking effect of penicillin treatment as compared with other methods was the freedom from pain post-operatively, especially in the tenosynovitis series, and the absence of post-operative complications (spreading infection, lymphangitis, adenitis). The most gratifying finding was the spectacular results in acute suppurative tenosynovitis, a condition attended by distressing results in our hands prior to aspiration-injection therapy.

Conclusions

Penicillin used locally is effective in the treatment of infections of the fingers, such as pulp whitlow, paronychia, and web-space infections, when the organism is susceptible.

It is possible to get perfect results in the treatment of acute suppurative tenosynovitis of the fingers by aspiration of the sheath and injection of penicillin.

It is important to remove slough when using penicillin locally.

Local penicillin has no apparent beneficial effect on the end-result in acute osteomyelitis of the phalanges.

Penicillin cream (in a new base—eucerin L.M.—Dr. Seiler) is an improvement on other methods of applying penicillin locally.

REFERENCE

Florey, M. E., and Williams, R. E. O. (1944). *Lancet*, 1, 73.

INFECTED BRONCHIECTASIS TREATED WITH INTRATRACHEAL PENICILLIN

BY

H. B. MAY, M.D., M.R.C.P.

Assistant Director, Clinical Laboratory, London Hospital

AND

M. A. FLOYER, M.B., M.R.C.P.

Medical Registrar, London Hospital

In most cases of bronchiectasis there is a secondary infection of the dilated bronchi and bronchioles. It is rare for this to be caused by one organism, but the mixed flora in the sputum often includes bacteria susceptible to the action of penicillin, and it was thought that this antibacterial chemotherapeutic agent might have a beneficial effect in the treatment of the disease. It is clear that, in general, a reduction in the bacterial infection of the bronchiectatic cavities, or even their sterilization, although reducing the volume of sputum, would not affect the deformity of the air passages, and infection would recur, but the improvement in the patient's local and general condition might enable a lobectomy to be carried out with a considerable reduction in the operative risk.

Although penicillin would be fully active in the presence of pus, the coexistence of a mixture of organisms of varying susceptibility implies the desirability of obtaining a relatively high concentration of penicillin in the cavities; this cannot, in our experience, be obtained by systemic treatment. An aerosol containing penicillin, as used by Knott and Clark (1945), would probably be equally ineffective because of the deficient aeration of the collapsed area of lung connected with the dilated bronchi. It was therefore decided that the best method of getting adequate penicillin to the infected area was by injection of a solution into the trachea, using a technique similar to that employed in carrying out bronchography.

Technique

The patient practised postural drainage for 20 to 40 minutes before the injection. The skin was rendered analgesic with 2% procaine in the midline of the neck, between the cricoid cartilage and the first ring of the trachea. 2 c.cm. of 2% amethocaine hydrochloride (decicain) was then injected rapidly through a No. 14 needle into the trachea and the patient was asked to try to avoid coughing. The syringe was removed from the

needle, which was left *in situ*, and the patient was placed in such a position that fluid introduced into the trachea would run into the bronchiectatic cavities—e.g., well over to the left if the left lower lobe is affected. A solution of penicillin containing 10,000 units per c.cm. was prepared and 5 to 8 c.cm. injected rapidly through the needle into the trachea, the needle being immediately withdrawn and the patient encouraged not to cough. The patient was kept leaning in the appropriate position for 15 minutes after the injection. It was found that with an injection of 50,000 units the sputum coughed up 12 hours after the injection contained between 10 and 100 units of penicillin per c.cm., and with the larger doses of penicillin—i.e., 80,000 units—the sputum after 24 hours still contained up to 10 units per c.cm. Daily treatment was carried out for eight to ten days, and the only complaint from the patients was of some soreness of the skin of the neck. All solutions were made up in saline, and a very fine needle could therefore be used.

Case Records

Case 1.—The wife of a publican was admitted to hospital on Oct. 15, 1944, complaining of pleuritic pain in the right lower chest and general malaise. She had had a perforated duodenal ulcer three months before; this had been sutured under inhalation anaesthesia, but owing to war conditions no details of her immediate post-operative progress are now available. Since the operation she had had recurrent colds and cough, with sputum, and a general feeling of malaise and undue shortness of breath on exertion. On admission it was found that she had coarse crepitations at both bases, with a pleural friction rub at the right base. After two days a small pleural effusion appeared on the right side; this resolved in one week with rest in bed. However, the moist sounds and bronchial breathing persisted at the left base, and a radiograph indicated some fibrosis in this area. Bronchograms revealed an early cylindrical bronchiectasis at the left lower lobe. Her sputum, which was fairly copious, averaging 1 oz. to 2 oz. a day, was purulent and non-offensive, and culturally showed a predominant growth of a pneumococcus sensitive to penicillin; no tubercle bacilli were present. Her haemoglobin was 86% (100% = 14.9 g. of haemoglobin), and she had a normal white cell count. Her sedimentation rate was the upper limit of normal. She was treated with postural drainage in hospital with some improvement in symptoms, but she still produced copious sputum. Intratracheal penicillin was then started, consisting of 50,000 units of penicillin intratracheally for 3 days and 80,000 units for the next 5 days. Owing to her domestic responsibilities she was given the second part of the treatment as an out-patient. Her sputum became mucoid instead of purulent, and culture showed only a few colonies of diphtheroids; it decreased in quantity to less than 1/2 oz. a day. After discharge the patient was symptomless for two months and remains well. The sputum is still mucoid, and is coughed up only with difficulty. She is able to do a full day's work.

Case 2.—A housewife aged 32 was admitted to hospital on Jan. 5, 1945, when 7 months pregnant, complaining of undue shortness of breath on exertion and cough with purulent sputum. She gave a history of pneumonia at the age of 14, since which time she had always had a cough with sputum. For one year she had been breathless on even slight exertion; this became worse during her pregnancy. She had lost her appetite and complained of the foul taste of her sputum. On examination she was cyanosed, and in the chest there were signs of bronchial spasm and coarse rales at both bases, more marked on the left than on the right. Bronchograms revealed bronchiectasis in the left lower lobe. Her sputum contained a varied bacterial flora, with a penicillin-sensitive pneumococcus predominating. She was treated with intratracheal penicillin, 60,000 units a day for 10 days. Her sputum remained copious, but was less purulent and less offensive, and aerobic culture showed a steadily decreasing number of colonies, the final specimen being sterile. One unit of penicillin per c.cm. was present in the sputum 24 hours after injection. Her general condition considerably improved, and she had a normal labour without undue shortness of breath or exacerbation of her pulmonary symptoms.

Case 3.—A gatekeeper aged 59 was admitted to hospital on Oct. 20, 1944, complaining of increasing cough and sputum and loss of appetite for five months before admission. He had had osteomyelitis of the right femur and a right lung abscess in 1940, since which time he had always had a cough, with occasional pain in the right chest and copious foul-smelling sputum. The latter had increased in quantity to 8 to 10 oz. a day, and was so offensive that he had lost his appetite; he had also become very short of breath. On examination he was emaciated and utterly miserable; the only relevant physical signs were at the right base, where there were diminished movement, an impaired percussion note, distant bronchial breathing, and coarse rales. His sputum contained a very varied flora, including a non-haemolytic streptococcus, pneumococci, and species of *Neisseria*, fusiforms, and spirochaetes. Most