are near New York. If either of these antibodies do turn up again, as they are sure to do sooner or later, it may be possible to correlate the present work.

Comment

There are two points of general interest in this history. First, the patient must be extraordinarily sensitive to antigenic differences, and this supports the hypothesis that the symptoms of the disease lupus erythematosus are manifestations of some undetected antigen-antibody reaction. Secondly, given such a hypersensitive recipient, a glimpse is obtained of the multiplicity of red cell antigens which may be found in the future. One familiar and three "new" antibodies have been made to the blood of eight donors, although considerable care has been taken to use appropriate blood. Landsteiner was of the opinion that blood groups would some day be known to be as individual as finger-prints.

Our thanks are due to Prof. Cappell and Prof. Fisher for continued interest and encouragement; to the donors for permission to use their names; and to the patient, who has been very co-operative.

REFERENCE

Race, R. R. (1944). *Nature*, **153**, 771.

A CASE OF NOCTURNAL HAEMOGLOBINURIA

BY

JOHN A. MILNE, M.B., Ch.B.

House-physician, Western Infirmary, Glasgow

(From the Gardiner Institute of Medicine, University of Glasgow)

The syndrome of paroxysmal nocturnal haemoglobinuria with haemolytic anaemia was first recorded by Marchiafava and Nazari in 1911, and according to Wintrobe only 41 cases had been described in the literature up to 1943.

Case Report

The patient, a Royal Marine aged 27, was admitted from a naval hospital for special investigation to the wards of Prof. J. W. McNee in the Western Infirmary, Glasgow. He was complaining of anaemia, with frontal headaches, tiredness, and weakness. He entered the Navy 8 years ago and his only illness was in Oct., 1943, while in the Faroe Islands. It was characterized by diarrhoea, vomiting, and urticaria, which was diagnosed as food poisoning. He has never had malaria. He is married and has one healthy child aged 21 months. There is no history of anaemia or jaundice in the family.

In March, 1944, the patient began to suffer from frontal headaches while working. These persisted unless he sat down and rested for a short while. These headaches gradually increased in severity, and he noticed that he was becoming very easily tired and breathless on exertion. In May, 1944, he reported sick, and was admitted to a naval hospital, where he was found to be profoundly anaemic. Three weeks later he developed a painful tongue and tender bleeding gums. The condition yielded rapidly to large doses of ascorbic acid, and glycerin and borax applied locally. For the anaemia he received iron and a variety of liver preparations without any improvement. When he had been two months in hospital he was given one pint of blood, with slight improvement in haemoglobin level. A month later he was given another pint of blood of correct group, after which, for the first time, his urine was noticed to be very dark in colour for two days following the transfusion. He was admitted to the Western Infirmary on Dec. 9 for special investigation, as the exact diagnosis of the anaemia remained uncertain.

On admission he was seen to be a pale well-nourished man with a slight icteric tinge of the skin and sclerotics; a few discrete hard lymphatic glands were present in both axillae and groins. The heart, lungs, and nervous system showed no abnormality. The liver and spleen were not palpable.

Blood Picture.—The following investigations were made, using the technique described by Wintrobe. Haemoglobin was estimated by means of a photo-electric colorimeter. Erythrocytes, 3,060,000 per c.mm.; leucocytes, 2,400 per c.mm.; haemoglobin, 11.4 g. per 100 c.cm.; packed-cell volume, 32.5%; mean cell volume, 106 c. μ ; mean corpuscular haemoglobin concentration, 34.8%; reticulocytes, 12.5%. Differential leucocyte count: neutrophils, 44.5%; eosinophils, 2.5%; lymphocytes, 48%; monocytes, 4%. A film stained with Leishman's stain showed anisocytosis, slight poikilocytosis, and macrocytosis of the erythrocytes. There was no evidence of malaria parasites or pigmentation of leucocytes.

On sternal puncture marrow was easily obtained; it appeared to be hypoplastic, as the film made from it contained numerous fat spaces. A few primitive erythroblasts were seen, but the marrow was definitely normoblastic. The blood plasma gave a positive indirect van den Bergh reaction—0.7 mg. of bilirubin per 100 c.cm. The fragility of the patient's erythrocytes to saline was within normal limits, haemolysis starting at 0.45%. Wassermann reaction, negative. A fractional test meal showed a normal hydrochloric acid curve. The urine contained a quantity of urobilin. At this point a haemolytic anaemia was diagnosed, the cause, however, being unknown.

Progress.—One week after admission the patient complained of pain in one of his lower molar teeth. The tooth was extracted under local analgesia on Dec. 18. At 2 a.m. on Dec. 19 he awoke and complained of an ache across his lumbar region, and a specimen of urine passed at this time was noticed to be dark red. Another specimen of urine at 6 a.m. was still dark in colour, while one at 9 a.m. was perfectly clear. The van den Bergh reaction on Dec. 19 showed a positive indirect reaction—1.4 mg. of bilirubin per 100 c.cm. Early on the morning of Dec. 21 he again passed a dark-red specimen of urine, while at 9 a.m. the urine was clear. Examination of the urine showed the colour to be due to a haemoglobin derivative. No erythrocytes were seen. The haemoglobin derivative was identified spectroscopically as oxyhaemoglobin. Since then haemoglobinuria has recurred regularly, the specimen passed in the early hours of the morning being dark red, that passed about five hours later being perfectly clear.

Further Investigations.—(1) The Donath-Landsteiner test was performed in order to exclude haemoglobinuria due to cold; it was negative. (2) In order to determine whether the fall in the pH of the blood during sleep was a factor in the production of the haemolysis (Ham, 1939) the patient's erythrocytes and serum were set up against an isotonic system over a pH range of 6.5 to 7.0 by Dr. W. B. Kyles, clinical pathologist to the Infirmary. The result was quite negative. (3) Even the clear specimens of urine contained a constant deposit of haemosiderin, demonstrated by the hydrochloric acid-potassium ferrocyanide reaction (Wintrobe). (4) As necrosis of the liver is a feature of this disease (Scott, Robb-Smith, and Scowen, 1938) liver function was estimated by the laevulose-tolerance and Quick's sodium benzoate tests, but was found to be within normal limits. (5) A specimen of serum was examined by Prof. D. F. Cappell; this identified the patient as Group O and Genotype Rh,Rh₂. (6) In order to bring out the association of the haemoglobinuria with sleep, his sleep rhythm was altered and he slept through the day and remained awake at night. This caused the haemoglobinuria to appear through the day while the patient slept, and was absent at night when he remained awake.

Discussion

The pathology of this disease and the mechanism of production of the haemoglobinuria are fully discussed by Scott, Robb-Smith, and Scowen (1938). The hypothesis put forward for this explanation of the nocturnal haemoglobinuria is the alteration in the pH of the blood during sleep (Ham, 1939). While sleep undoubtedly is a factor, as seen by altering the sleep rhythm, the laboratory experiment carried out by altering the pH of a specimen of blood, while not conclusive, would tend to point against alteration of the pH being part of the mechanism, as the range of pH covered was much greater than would occur in the human body. It is known that transfusions and trauma are factors which increase or precipitate the haemolysis. This is well shown in this case.

Summary

A case of chronic haemolytic anaemia associated with nocturnal haemoglobinuria is presented. Blackwater fever, syphilis, and Lederer's anaemia were excluded. The patient was Rh-positive. The existence of the nocturnal periodicity of the haemoglobinuria was confirmed by altering the sleep rhythm, thus indicating that the patient suffered from the Marchiafava-Micheli syndrome.

I should like to thank Prof. J. W. McNee for his permission to publish this case.

REFERENCES

- Ham, T. H. (1939). *Arch. intern. Med.*, **64**, 1271.
Scott, R. B., Robb-Smith, A. H. T., and Scowen, E. F. (1938). *Quart. J. Med.*, n.s., **7**, 95.

The Mothers' Hospital of the Salvation Army in Lower Clapton Road, London, E.5, is entirely staffed by medical women. The report for 1944 is curtailed in the interests of economy, but space has been found for a foreword by Miss Margaret Basden, senior obstetric surgeon, who on the outbreak of war took up residence in the hospital. She writes in moving words of the fine spirit with which the nursing staff and patients and the fire-watchers endured the ordeal of air raids.