Discussion

The nine survivors were among the most appreciative and grateful patients I had in Iraq. They were able to return to their husbands and homes and begin a normal life again. It can be realized what that meant when one remembers that their average age was between 18 and 25 years. Nevertheless a far more efficient follow-up system than is possible in Iraq is necessary to compute the end-results of the operation. In Grey Turner's series some patients survived for 20 and 30 years. The mortality of 2 deaths in 11 cases makes the operation one of considerable hazard, certainly not to be undertaken lightly. But the lot of a patient with an incurable vesico-vaginal fistula is such a miserable one that I believe it to be justified. In more skilled hands the mortality is likely to be considerably lower.

Summary

Eleven cases of ureterocolostomy for incurable vesico-vaginal fistula are described. The operative modification whereby incision of the bowel is avoided is described.

There were two deaths in the series, both from peritonitis; the remaining nine patients returned to active life.

It is a pleasure to acknowledge the great help and advice given by Brig. C. Naunton Morgan in this series, the stimulation and practical assistance of the writings of my teacher and chief, Prof. Grey Turner, and the data of Prof. M. McDowall, of the department of gynaecology, Royal Hospital, Baghdad. Dr. Kadhim Shubbar, Dr. Najib-al-Yacubi, and Dr. Ghanim Akrowie were of the greatest assistance in pre- and post-operative care.

REFERENCES

- Lloyd, Seton (1943). *Twin Rivers: A Brief History of Iraq from the Earliest Times to the Present Day*, Oxford Univ. Press.
Turner, G. Grey (1943). *British Medical Journal*, 2, 535.

OTITIS EXTERNA

BY

J. F. BIRRELL, M.D., F.R.C.S.Ed.

T/Major, R.A.M.C.

Otitis externa is one of the most common conditions met with in any ear-nose-and-throat department. This is especially true in the Army over-seas, where it may account for one-quarter of all aural cases, or one-sixth of all E.N.T. patients examined. There are many problems connected with the disease that are worthy of study, and, indeed, many to which there is as yet no answer. This report is mainly based on nearly 1,000 cases of otitis externa seen in the Army. Furunculosis and dermatitis of the auricle are not discussed in any detail, but are mentioned briefly as coexisting affections.

Aetiology

The disease does not respect age or rank. It may occur in children and persist at intervals throughout life. The mechanism appears to be twofold. A meatal congestion is set up, causing irritation, which is attacked by the patient, who scratches the ear voluntarily or involuntarily, or by the doctor, who attempts to diagnose, treat, or remove the cause of the irritation. The result is a break in skin continuity, through which normally present saprophytic or invading pyogenic organisms may enter and cause trouble.

The meatal congestion may be caused by (i) irritants which enter the meatus from the middle ear, (ii) the natural secretions of the meatus, (iii) irritants entering the canal from the outside, (iv) general conditions predisposing to skin infections, or (v) local or general allergy. A brief consideration of the anatomy of the external meatus will show how easy it is for foreign material of all kinds to reach the canal, how difficult it may be for such matter to find its way out once it has reached the downward-sloping inner half of the meatus, and how simple it is to injure the meatal skin in an attempt to remove the irritant. Daggett (1942), in a valuable article, has drawn attention to the deep anterior meatal recess as a reservoir which may retain pus and debris.

The irritant entering the meatus from its inner end is pus, associated with an active otitis media. The meatal floor is constantly bathed in pus, the organisms of which welcome any

disruption in the epithelial surface through which they may enter and multiply.

Wax and sweat are the natural secretions of the external meatus. Wax is secreted by the ceruminous glands of the outer half of the meatus, lines the canal, separates from the walls apparently by the movements of the mandible, and works its way out, bringing with it desquamated epithelium. Unskilled attempts to remove the wax by a match, a Kirby grip, a knitting-needle, or some such similar article may abrade the skin. Similarly, the meatal epithelium may be lacerated by the tip of an aural syringe, especially if the wax is hard and difficult to remove. After syringing, the canal should be thoroughly dried with wool or gauze, and the walls painted with spirit, or the retained solution, if hypotonic, may macerate the skin. In the same way the epithelium may suffer during the removal by scoop, curette, or syringe of the tightly adherent mass of cerumen and epithelium called keratosis obturans, which in its milder forms is more common than is often realized. Sweat may be produced in excess in hot weather, and may, by the discomfort it causes, lead to scratching of the ear. It may cause a matted condition of the hair at the outer end of the meatus, and this in turn results in irritation.

Many things may be introduced into the meatus through its outer end. One of the most common, particularly in soldiers serving over-seas, is salt water, and in the "tideless" Mediterranean the water is permanently dirty in frequented beaches. Daggett, while discounting sea-bathing as an important aetiological factor in desquamative otitis externa, points out that in the heat of the day the water will be evaporated from the meatus, leaving a deposit of salt which may increase skin maceration by hygroscopic action. In Italy these swimming ears presented such a characteristic appearance that one came to regard them as a specific infection.

Cotton-wool or paper may be pushed into the meatus and forgotten—for example, by gunners who use these substances as ear-plugs against muzzle blast. After some weeks the meatal skin reacts to the breakdown of vegetable matter by producing an otitis externa which is often most acute. Dust and sand may be blown into the meatus and, mixing with wax and sweat, cause irritation. Insects, such as the bed-bug or the maggot, may find their way into the canal and, by their movement, scratching, or biting, may set up a most acute infection. Lastly, infection may be transmitted by the finger from some other part of the body.

Chemicals may act on the meatus by a direct burn, such as occurs with caustics, acids, hydrogen peroxide, vesicant gases, etc., or by a local allergic reaction, such as is seen with sulphanilamide powder in susceptible people. Insufflation of sulphanilamide powder was a common practice in the Tunisian campaign, but was discontinued in Italy as a result of adverse reports from otologists. The powder tends to form a hard caked mass, and otitis externa occurs in a significant number of cases.

General conditions which predispose the skin of the meatus to infection are seborrhoeic dermatitis, especially of the scalp, and relapsing furunculosis. It is possible that vitamin deficiency may produce the condition, although there is no positive evidence as yet. In recent years much thought has been given to allergic conditions of the ear, and otitis externa of purely allergic origin has been described.

Bacteriology

Friedman and Hinkel (1941), studying a series of 200 cases, found that in every instance more than one organism was present, and that the commonest were *Staph. albus*, diphtheroids, *Ps. pyocyanea*, and micrococci. Furthermore, they stated that in those cases which most resisted treatment no single organism predominated. A study of 25 normal ears showed the same group with the exception of *Ps. pyocyanea*. Daggett reported that diphtheroids, either alone or in combination with *B. proteus*, *Ps. pyocyanea*, or *Staph. albus*, were found most frequently in 49 cases. In this hospital *Ps. pyocyanea* was obtained as the predominant organism in nearly every case in which a swab was taken. An effort was made to swab the ears of patients showing different varieties of the condition in the hope of being able in future to diagnose the predominant

organism from the clinical appearances. This hope was not fulfilled. Apart from the constant presence of pure *Ps. pyocyanea* with green pus in the meatus, no association could be discovered. The subsidiary organisms were the same as those found by Daggett.

Varieties

The condition is usually described as being of two types—wet and dry—but this is inadequate for a proper understanding of the disease. The earliest manifestation is meatal injection. When infection supervenes there is desquamation of the meatal epithelium and pus formation. In some cases this is more acute, and is associated with a marked oedema of the meatal walls. Prolonged or recurrent infection leads to a hypertrophy of the epithelium, which is thrown into ridges. The various stages may be called congestive, exudative, oedematous, and hypertrophic; or the varieties named mild, acute, acute oedematous, and chronic hypertrophic otitis externa. Some cases, however, do not fit into this picture. There is the fungous infection, in which the debris resembles wet blotting-paper; the membranous variety, commonly found after sea-bathing, in which the membrane probably contains salt; and the granular type, in which granulations may form on the meatal wall and spread on to the intact drum.

Symptoms

The earliest complaint is of itching or irritation, which may be so severe as to be unbearable. In the congestive stage this is the only symptom. Then discharge appears—at first watery, but later becoming purulent and finally a cheesy debris as the pus mixes with desquamated epithelium. A purely external infection never results in a mucopurulent discharge, as there are no mucous glands in the meatus. Pain may be felt, depending on the amount of tension under the epithelium and the speed with which the tension is set up. Meatal skin is tightly bound down to perichondrium, and thus a furuncle or acute meatal oedema may cause extreme pain. The discomfort is referred, as a rule, to the region of the parotid gland and to the jaws, and is aggravated by any movement of the temporomandibular joint.

Where the middle ear is normal hearing is not affected, but if the meatal lumen is occluded by oedema or debris deafness may be complained of. Other symptoms are a feeling of swelling in the ear, associated with a slowly produced oedema; and dermatitis when the infection spreads to the auricle.

Appearances

It is important to examine the auricle before inspecting the meatus, as the diagnosis can often be made before a speculum is inserted. In a very large proportion of cases the skin of the anterior incisura, between the tragus and the lower anterior end of the helix, will show infection—a small crust, a fissure, or a small raw area. Where this is found the diagnosis of otitis externa is certain, although there may be a coexisting otitis media. Dermatitis of the auricle from an otitis media alone is found in the line of the pus—along the meatal floor, the floor of the concha, and via the incisura intertragica to the skin of the fissure between the lobule and the angle of the jaw.

In the external meatus congestion is the earliest sign of trouble. It often affects the floor about the junction of cartilaginous and bony meatus. In the exudative stage the meatus contains a varying amount of pus and desquamated epithelium. This must be carefully and gently mopped away with wool-tipped probes, particular attention being paid to the anterior meatal recess. A light wire probe and a light touch are the secrets of success in this manoeuvre. The probe should be held so loosely that the finger and thumb will slip along the shaft if the tip is impeded in any way. One must remember that the skin lining the inner half of the meatus is extremely sensitive to the touch. Thorough cleansing is essential for accurate diagnosis and successful treatment, and, while mopping under direct vision is ideal, the same result can be achieved by blind mopping if care and patience are exercised and frequent examinations with the auriscope made to determine how much secretion is left. In cases in which the epithelial debris is difficult to remove it is often helpful to moisten the tip of the mop with 20% sodium sulphate solution.

When the meatus has been cleaned the epithelium will be found to be rough, red, and desquamating. The outer surface of the drum may share in this process, leading one to suspect an otitis media. In the acute oedematous type the narrowed meatal lumen is obvious. Pain will be experienced on insertion of the speculum, and no attempt should be made to force its passage. Mopping—necessarily blind mopping—should be carried out as described. An estimate of the amount of oedema can be obtained by comparison with the other ear if it is healthy. The narrowing of the lumen in chronic otitis externa is much more resistant to the passage of a speculum than in the oedematous cases. This is due to hypertrophy rather than pain. Pus and debris will be found, and there is often a coexisting perichondritis of the auricle.

Regional adenitis may be present, the glands involved being situated in front of the tragus or over the tip of the mastoid process. Systemic upset varies a good deal. Some patients have no general disturbance, while others look ill, feel ill, have a raised temperature and pulse rate, show a furred tongue, complain of loss of appetite, and have difficulty in sleeping.

Differential Diagnosis

Diagnosis must be made from otitis media (see Fig.) and, if pain be present, from mastoiditis. The criterion in distinguishing between otitis externa and otitis media is the appearance of the drum, and, as the two conditions often coexist, one is not justified in ruling out otitis media until every part of the drum

1. Auricular spread in otitis externa. 2. Anterior auricular glands. 3. Auricular spread in otitis media. 4. Posterior auricular glands.

has been inspected for perforation. In many cases perforation in an oedematous congested membrane is found only after a diligent search. A purely external otitis is suggested if the hearing is good and there is no mucus in the discharge. The presence of mucus can be tested for by leaving a wick of ribbon gauze, soaked in 5% lead acetate solution, in the meatus for 24 hours. On removal, if mucus is present in the discharge the innermost end of the wick will be found to be black, while if the discharge is purely external in origin no colour change will be seen. This test has been used in doubtful cases for a year now, and has proved most reliable.

The diagnosis between otitis externa and mastoiditis is often difficult. Both conditions result in otorrhoea with pain, both may be associated with constitutional upset, and in both the mastoid region may be tender and oedematous. In mastoiditis tenderness is maximal on deep pressure, especially over the mastoid antrum, while in otitis externa tenderness is maximal on superficial pressure over the glands at the tip of the mastoid process. In external infection, also, there is tenderness on pressure over the tragus or on movement of the auricle. X-ray examination of the mastoids will show no abnormality if the middle ear is normal and if there is no post-auricular oedema.

Prognosis

The prognosis in any attack of otitis externa is good, a cure being effected in 2 to 14 days, depending on the severity of the infection: 60 to 70% of cases are healed within a week. Where chronic otitis media is present as well, healing takes longer, although the meatal infection should clear up in three weeks. Dermatitis of the auricle in association with otitis externa does not materially lengthen the time for healing in most cases, provided adequate treatment is given.

Unfortunately, relapses are common. The percentage of cases which recur cannot be measured accurately in the Army, as patients discharged from hospital do not necessarily stay in the locality, and thus an otologist cannot follow up his cases.

A relapse means either that treatment has been inadequate or that reinfection has occurred. No particular organisms appear to be responsible. Chronic otitis media is often the cause of a relapse. Frequent recurrences or lack of treatment lead to chronicity which, once established, is well-nigh impossible to cure.

Treatment

Brain (1942) stressed that "treatment should be governed by two important facts. The first is that our protection from external irritants and infections depends upon a dry and intact horny layer, and treatment should be designed to maintain it. . . . The second fact of therapeutic importance is that living cells are instantly damaged by water or by grossly hypotonic solutions, and this applies to the skin when the protective stratum corneum is missing, or its continuity is disrupted by exudation."

Treatment resolves itself into two approximately equal parts—the thorough cleansing from the meatus of all particles of discharge, and the application of lotions to the inflamed meatal walls. The mechanism of mopping has been described, and its importance cannot be too strongly emphasized. Drugs are applied to the infected skin on wicks of ribbon gauze or painted on by means of wool-tipped wire probes. The ribbon gauze wick must be inserted gently so that it maintains contact with the walls in the whole length of the meatus. It should be changed daily, and some authorities moisten the wick with the appropriate solution every 12 hours. Cotton-wool adheres to the raw skin surface and should never be used. The selected solution must be non-irritant and isotonic; it should not stain the meatal walls or leave a deposit thereon; it should not dry enough to cause pain on removal. Pastes and powders allow infection to spread under them.

An acute oedematous otitis externa must first have the oedema reduced to permit complete cleansing of the drum and the deep anterior recess. Then the inflamed epithelium must be healed and finally dried and hardened. The drugs used for reducing oedema are 20% sodium sulphate, 8% aluminium acetate, or 10% ichthyol in hypertonic saline. The first two both tend to dry off during the course of 24 hours and are thus painful to remove. The ichthyol preparation has been used by me for nearly three years, and has proved most efficacious. After 48 hours' treatment the case usually resembles one in the exudative stage.

In this stage numerous solutions may be used to heal the skin. Those already mentioned may be persevered with, or one may try 10% ichthyol in water, 10% ichthyol in glycerin, 5% lead acetate, 10% argyrol, or calamine lotion. Of these, ichthyol in water dries quickly and is painful to remove; calamine lotion leaves a deposit on the skin which obscures the view of the progress; lead acetate has the same fault to a lesser degree, but has considerable healing properties, and is, in my experience, more efficacious than argyrol, which is superior for inflamed mucous membranes. Ichthyol in glycerin is probably the best all-round treatment, being soothing and easy to insert and remove.

When no further exudation is found the case resembles one in the congestive phase. Here a paint of 1% silver nitrate or 1% alcoholic solution of gentian violet should be used to harden the meatal skin. The latter solution stains the meatal skin and should not be employed until all discharge has stopped. It is the best treatment for fungous infection and for the granular type. When pus is green, or if *Ps. pyocyanea* predominates without green pus, 1% acetic acid, or a mixture of 1% acetic acid and 10% ichthyol, will give good results, although some patients find them uncomfortable, and they tend to dry off.

Pain is best treated with heat—fomentations, kaolin poultices, or hot-water bottles—and sedatives. The patient under treatment should be instructed not to scratch the ear, however itchy, and not to let water enter the meatus during washing. He should be told specifically not to use cotton-wool as a plug while the ear is discharging, as this only increases skin infection. Sea-bathing is forbidden for several months.

If a dermatitis of the auricle is present, good results are obtained with 1% ammoniated mercury in Lassar's paste applied on spreads and moulded to the pinna. Five per cent. sulphathiazole paste was efficacious, but was discontinued when it

was found to produce local reactions and even generalized sensitivity to light.

The question of chemotherapy in otitis externa is a vexed one. Many patients had had a course of one of the sulphonamide drugs before reaching hospital, with no benefit to the local condition. One has seen cases in which penicillin has been given parenterally for wound infections, furunculosis, or the aural condition itself, and yet otitis externa persists. The reason is obvious. Otitis externa is a mixed infection, and most of the causal organisms are insensitive to penicillin and the sulphonamides.

Syringing the ear in a case of otitis externa is always liable to increase the infection; but it is necessary sometimes when the secretion in the anterior meatal recess is too hard for mopping. To leave this debris is to invite recurrence. I have been in the habit of syringing these cases on the first day on which the ear is dry, and treating with ichthyol in hypertonic saline or ichthyol in glycerin for a day or two before going on to the paint.

It is with pleasure that I record my thanks to Col. C. H. K. Smith, O.B.E., M.C., commanding officer of a general hospital, for permission to publish this article.

REFERENCES

- Brain, R. T. (1942). *Proc. roy. Soc. Med.*, **35**, 8, 519.
Daggett, W. I. (1942). *J. Laryng. Otol.*, **57**, 427.
Friedman, H. S., and Hinkel, C. L. (1941). *Arch. Otolaryng.*, Chicago, **33**, 749.

PENICILLIN GROWN FROM A NUTRIENT MEDIUM PREPARED FROM POTATO EXTRACT

BY

J. O. GAVRONSKY, M.D.

The production of penicillin is rapidly growing. This is likely to continue for some time to come. Other antibiotics in application to agriculture, horticulture, and veterinary surgery will soon have to be produced on a large scale, possibly on a scale exceeding the production of penicillin. Under these circumstances the question of preparation of nutrient media becomes of great importance.

Conditions for growing moulds, to obtain the needed product of their metabolism, vary with the mould one is dealing with. To a large extent the result depends on the nutrient medium used. This medium must be both satisfactory and cheap. With very few exceptions the composition of such media is kept secret. In making known the composition and mode of preparation of a nutrient medium used by me I cannot claim to get a higher content of penicillin in the metabolic solution than that obtained in some of the media already used. I can only say that the titre of penicillin compares favourably with that obtained by using Czapek-Dox medium with the addition of yeast or corn-steep liquor, or by using the medium introduced by Challinor and McNaughton. The preparation of the medium is as follows:

Method

Potatoes or potato peelings are hydrolysed in twice the quantity of tap-water with 2% of HCl on the weight of potatoes. The hydrolysis proceeds for 2½ hours at a pressure of 3 atmospheres. After filtering the hydrolysate through a linen cloth an extract is obtained with an average glucose content of 5.5%. The sugar content naturally varies with the starch content of the potatoes or potato peelings used. In the case of potato peelings, unless they are used soon after peeling, a longer time is needed for hydrolysis. In the case of potatoes, these are washed and cut into pieces of approximately 1/2 to 2 cm. thickness. The necessity for cutting is avoided by using potato peelings.

The quantity of extract thus obtained from 1 kg. of potatoes with 2 litres of water and 20 c.cm. of strong HCl, after disposing of the solid residue by wringing out the linen bag used for filtering, is approximately 2,450 c.cm. This extract is diluted to 4% glucose content. We thus obtain 3,360 c.cm. approximately. The pH is roughly adjusted to neutral to litmus. Corn-steep liquor is added, 370 c.cm.—that is, 10% of the total quantity of medium. NaNO₃ is added—3% per litre; and activated charcoal—1/2% on weight of potatoes. The pH is adjusted to 6, and the medium sterilized