best, even though only temporarily, to inhalations of CO₂ and oxygen. Cyanosis and respiratory distress responded well to continuous oxygen administration, though this treatment had to be continued as long as three days in some cases. Patients with meningitic symptoms were lumbar-punctured, principally from the diagnostic point of view. Removal of cerebrospinal fluid did not appear to alleviate the symptoms.

Treatment of Complications and Concurrent Diseases

Lobar and lobular pneumonia due to secondary pyogenic infection were treated with sulphathiazole in the usual dosage. Of 7 cases so treated, 3 developed empyema. Treatment adopted for this was repeated aspiration followed by instillation of penicillin into the pleural cavity, combined with intramuscular penicillin in dosage of 100,000 Oxford units daily. This treatment led to resolution in two cases, but the third became contaminated with *B. coli*, and rib resection was necessary to complete the cure.

The case of parotitis cleared up without resort to surgery. The two cases of femoral thrombosis were treated along the usual lines, but recovery was marred by a residual oedema of the leg. Retention of urine called for twice-daily catheterization until bladder control was regained. Bacillary and amoebic dysentery were controlled with sulphaguanidine and emetine respectively.

Summary

A series of 110 cases of scrub typhus in British patients is described.

The disease generally presented a clear-cut symptomatology, and clinical diagnosis was not difficult.

The Weil-Felix reaction becomes positive in significant titre too late in the illness to be of much assistance in the early diagnosis.

The differential diagnosis and prognosis are discussed and the pathological findings in the fatal cases recorded.

We wish to thank Col. P. H. Addison, I.M.S., for permission to present this account

BIBLIOGRAPHY

D.M.S., A.L.F.S.E.A. Technical Memo. (Medical) No. 6, 1944.
Ransome, G. A., Gupta, L. M., and Paterson, J. C. S. (1944). *British Medical Journal*, 2, 594.

THE EFFECTS OF MECHANICAL STRESSES ON MAN*

BY

B. H. C. MATTHEWS, C.B.E., Sc.D., F.R.S.

LECTURE II

In aircraft man moves at very high speeds, but high speed in itself does not produce any physiological effect if the pilot is protected from high wind-pressures in a closed cockpit. However, when motion changes rapidly, either in direction or in magnitude, the pilot is subjected to very great mechanical stresses not ordinarily met with on the ground. The practical effects of these may be separated into two classes of phenomena: (1) those associated with change of speed such as occurs in catapult-launching or in crash-landing, where high acceleration or deceleration acts for a short time; and (2) those associated with change of direction of motion, which may be of long duration but of less magnitude, as when an aircraft executes a tight spiral. The physiological effects produced depend on the value, duration, and line of action of the force acting on the body and also the area over which the force acts on the body. The magnitude of the force acting on a body, owing to its inertia during change of motion, depends on the rate of change of velocity and the mass of the body. The numerical factor to be related to physiological effects is the change of velocity in feet/second/second. The acceleration due to gravity giving a body its normal weight is 32.2 ft./sec./sec., abbreviated in the sign "g," and it is convenient in physiological work to consider the rate of change of motion in terms of multiples of "g." This makes it possible to give the forces acting a numerical value. It must be

emphasized that the acceleration communicated to one part of the body, if it acts only for a very short time, may not produce so great an acceleration on other parts of the body owing to transmission of the forces via parts of the body which yield. For example, if the chair in which a man is seated is given an upward acceleration of 5 g for only one-fiftieth second, the acceleration at the head will have a much lower value and act for a longer time owing to transmission of the force through yielding parts of the body.

If large forces act on a small area the commonest result is a penetrating injury. It is forces acting over a small area in this way which produce all common accident injuries, and to prevent these it is important in aircraft design to avoid projections in the cockpit with which the pilot may come into violent contact if the aircraft is suddenly decelerated, as in crash-landing. Injuries of this type are greatly reduced by the use of properly designed harness to prevent motion of the body relative to the cockpit. Here will be considered only those cases where a large force acts on the body over a considerable area and penetrating injury does not occur.

Physiological Disturbances

Fig. 1 gives in diagrammatic form the tolerance of the seated body to acceleration when it is adequately supported to meet the force. This is drawn as two separate lines, to distinguish between short-acting forces, which may result in mechanical failure of parts of the body, and long-acting forces of lesser

FIG. 1.—Acceleration tolerable to a seated man.

magnitude, which produce physiological disturbances, particularly of the circulation, without permanent structural damage. The upper part of the curve is only a rough indication, and requires further definition of the exact support of the body, which makes a very great difference to the tolerable acceleration. In other postures the tolerable acceleration varies widely. Thus in a standing man fracture of the lower leg is a possibility at 3 g, and is very likely to occur at 7 g. A seated man accelerated forward can tolerate very high acceleration, probably over 25 g. For acceleration in the reverse direction, such as the deceleration in crash-landing, the body must be restrained by a harness, and unless the head is restrained (which is not often practicable) neck injury may occur at less than 25 g. Above this, evisceration may occur. Very high acceleration can be tolerated if it acts on a man supported spread-eagle on his back—probably at least 250 g (de Haven, 1942).

These figures must be regarded as a rough indication only, for unless conditions of support are exactly defined, wide variations may occur in practical cases. They are based on evidence from several sources—viz., (1) From experiment. Squad. Ldr. Stewart has carried out many catapult experiments with various postures, experiencing the accelerations which can be tolerated without injury. (2) From analysis of accidents, where it is often possible to reconstruct probable forces that have been in action. (3) From an analysis of unsuccessful suicide attempts which has been made by de Haven (1942).

The problem of concussion is too extensive to be considered here. The forces necessary to produce concussion, if penetrating injury does not take place, appear from de Haven's figures to be very high indeed, perhaps 300 to 400 g. Experiments by Denny-Brown and Russel (1941) also show that very high acceleration is required to produce unconsciousness in animals, though in practical rather than theoretical cases the work of Holbourn (1943) suggests that injury may result to the brain when the head receives a much lower acceleration if the

* The second of two Oliver-Sharpey Lectures before the Royal College of Physicians of London, March, 1945.

force does not act through the centre of gravity of the head so that a sudden rotation is imparted to it. Damage to the brain may then result from its lagging behind the skull in rotation.

It is clear that these factors, and their proper application to aircraft construction and procedure, can do much to save life, especially in crash-landing, and much work has been carried out for the R.A.F., especially by Squad. Ldr. Pekarek, to give suitable protection to the body.

During prolonged acceleration physiological disturbances occur from the effects on the hydrostatic columns of blood in the body. Much physiological research has been carried out in the past on the effect of normal gravity on the circulation and the readjustments in the circulation which take place with change of posture. But in flight far greater forces, which easily produce unconsciousness, come into play.

"Blacking-out"

In the Schneider Trophy races pilots found that during turns at very high speed they suffered visual failure, which was given the name of "blacking-out." Experimental studies were carried out by R.A.F. medical officers Livingston (1939), Rook and Dawson (1938), Corner, and Wallace. Before the war intensive work on this subject was carried out in Germany, particularly by Ruff, by von Diringshofen (1934), and by Strughold. A centrifuge to reproduce the conditions of rapid turns was built in Berlin in 1935. By 1939 it was established that blacking-out was due to failure of the circulation above the heart owing to the great increase in effective weight of the blood while subjected to high acceleration. The subjective sensations under these conditions are those of a great increase in weight of all parts of the body. An increase to three times gravity makes it impossible to walk about, and at about four to five times gravity when seated it is impossible to raise the feet off the floor of the cockpit (Stewart, 1945). The pilot feels forced into his seat, and at 5 g, were his seat supported by a spring balance, it would indicate an apparent weight five times his own, so that a 200-lb. pilot would record about half a ton on the balance. This apparent increase in weight causes sagging of the soft tissues of the face, illustrated in the photographs (Fig. 2) of a

FIG. 2.—(i) Subject in straight and level flight. (ii) Subject at an acceleration of 4 g while turning. This illustrates the sagging of the soft tissues of the face owing to their virtual increase in weight.

subject in straight and level flight and during a tight turn. A number of R.A.F. aircraft have been equipped for research under these conditions, and a great deal of work has been

carried out, particularly by Squad. Ldrs. Stewart, Davidson, and Watt.

A centrifuge operating on the ground can approximately reproduce these conditions without the added hazards of flight, although, owing to the rapid rotation necessary, other effects are brought into play. These are of secondary importance to the large centrifugal effects. Centrifuges to carry men have been built in Australia, Canada, and the U.S.A. The R.C.A.F. centrifuge (Franks *et al.*, 1945) at Toronto is shown in Fig. 3.

FIG. 3.—R.C.A.F. centrifuge, Toronto, Canada.

The effects of centrifugal force are primarily due to disturbance of the circulation. The blood virtually becomes the weight of iron at 7 g, and, if the circulation in the brain is to continue, a pressure of over 200 mm. must be maintained in the aorta to force it upwards against these forces. Moreover, the venous return from the lower half of the body falls and, unless inflow is maintained, clearly arterial pressure must also fall. In addition the normal circulatory reflexes are greatly disturbed when subjected to these conditions. The first symptom is loss of vision, and this is due to failure of the retinal circulation, which occurs before failure of cerebral circulation because the intraocular pressure is opposed to that circulation. This does not occur immediately the force is applied, but lags behind it some 5 to 8 seconds. This lag presumably represents the oxidative reserve available in the retinal cells themselves. Experiments by Craik and Stewart in 1941 showed that increasing the intraocular pressure by pressure applied to the eyeball produces blacking-out after about the same time interval as occurs while flying; and applying pressure to the eyeball while under acceleration in flight causes blacking-out to occur at values of "g" at which vision is unimpaired with normal intraocular pressure. With increasing acceleration when a pilot progressively tightens a turn, peripheral vision is lost and central vision becomes hazy before complete black-out. When just blacked-out, other cortical functions are less obviously impaired. Squad. Ldr. Stewart showed that hearing and smell are still functional; if the acceleration is still further increased unconsciousness results, with loss of muscular tone; as this occurs other neuro-muscular disturbances are seen. When the acceleration is removed consciousness returns rapidly, though after severe blacking-out convulsive movements of the limbs may be experienced, and the subject suffers retrograde amnesia and is disorientated (Franks *et al.*, 1945). The factors determining the severity of blacking-out are: (1) magnitude of the acceleration; (2) duration of the acceleration; (3) posture of the subject; (4) individual susceptibility.

So far a normal seated posture has been considered, and the average tolerance of acceleration under these conditions is roughly 5 to 6 g for five seconds.

Protection against Centrifugal Force

There are three practical methods of protecting the circulation against centrifugal force: (1) "Self"-protection: voluntarily increasing the tone by contracting muscles of the lower half of the body is found to increase by about 1 g the threshold at which blacking-out occurs. (2) Posture: by reducing the length of the columns of blood involved. This can enable a man to increase his tolerance of acceleration. Fig. 4 gives a diagram-

matic summary of the effects of various postures on blacking-out thresholds. (3) Protection by the use of special garments.

In a normal sitting posture there are two processes leading to a failure of circulation in the head. The increasing weight of the blood requires increasing pressure in the aorta to maintain the circulation, and the venous side of the heart may be starved by pooling of the blood in vessels below that organ. This venous drainage may be reduced by surrounding the lower half of the body by a garment which applies equal pressure to the

sary to apply a similar gradient in the external pressure applied to the surface of the body. The Franks hydraulic suit effected this automatically, as the gradient of pressure in the water is similar to that in the columns of blood in the limbs. The first air suits applied greater pressures to the lower parts of the legs than to the abdomen and thighs; but the advantages of this have not been found to be worth the added complications, and a single pressure applied to the body surface below the heart is very much simpler to achieve and will effectively maintain the circulation. The value of protection of the circulation during tight turns is that not only does it increase the manœuvres the pilot can carry out, which may be of value in "dog-fighting," but frequent application of acceleration insufficient to produce blacking-out is found to cause very marked fatigue, and this is avoided by the use of these garments.

It is at present undesirable to give pilots unlimited protection against acceleration. If this could be done many mechanical failures of aircraft would occur, for aircraft designers in the past have accepted strength limits based on the human tolerance of acceleration, and by making the man stronger than the aircraft in resisting "g" a new danger arises. There is also a physiological cause to avoid very high "g" until more perfect protection can be achieved; for with artificial maintenance of the venous inflow to the heart the great pressures required in the aorta to maintain upward circulation constitute a very great abnormal load on the heart and also raise problems of the structural strength of blood vessels at or below heart level. It is clear that if venous return is kept up and the acceleration increased still further, a point must ultimately be reached where myocardial and other damage will result. Animal experiments by Greenfield (1945) in England and Jasper and Cipriani (1945) in Montreal suggest that this limit is at about 12 g, and may well be lower in man. Fortunately, up to the present there has been no evidence of cardiac damage in the application of up to about 9 g; but it is clear that this is a very serious aspect of the problem, and if danger exists here it may not become apparent for a number of years after the conditions initiating it. It is therefore very desirable that, for the present, the loads placed on the heart in this way should be restricted.

The vascular system of the legs is able to tolerate very considerable increase in pressure during positive acceleration, though petechial haemorrhages are on rare occasions produced locally in the lower leg. However, the factor of safety for abnormally high pressures is very much less in the head. When an aircraft executes an outside loop so that the pilot's head is away from the centre of the circle, the blood is forced in the reverse direction from that already considered, and the pressure in blood vessels of the head may rise to a dangerously high value. An acceleration of about 3 g can be tolerated without danger, but a "redding out" of vision resulting from congestion of the retina may occur. Higher values of acceleration occasionally arise by accident, and haemorrhage in the conjunctiva is to be expected at 4 to 5 g. With higher values of acceleration there is a great danger of haemorrhage in the cerebral vessels, with serious and possible permanent damage to the brain, and for this reason manœuvres which would produce large centrifugal forces acting towards the head are avoided in aircraft.

Conclusion

In these lectures I have tried to give an outline of the two stresses to which large numbers of flying personnel are exposed. Our present knowledge on these problems has resulted from team-work not only on a national scale but, in the last few years, on an international scale between all the Allies. The extensive research organization that the Director-General of Medical Services, Sir Harold Whittingham, has set up for the R.A.F. is kept very fully occupied, and but for the great resources of our Allies and the Empire many of the problems could not have been studied in detail. In the R.A.F. the performance of men and machines has been stretched to the limit to outfly the enemy, but in civil flying the first aim must be to reduce stress on the passenger. While the stresses considered above should never arise in civil flying, knowledge from war research is applied to make civil aviation safer.

Perhaps it is not too much to hope that these researches carried out in the interests of war may prove of value outside aviation in peace. Our knowledge of normal function may

FIG. 4.—The effect of posture on the blacking-out threshold of a human subject (redrawn after Ruff).

outside of the abdomen and lower limbs, to counter the outward pressure in the blood vessels while under acceleration.

The first scheme of this sort was that put forward by Marshall in 1933, using air bladders to compress the abdomen; the pressure being provided by the forward motion of the aircraft from a scoop which opened under the action of centrifugal force. An alternative scheme provided pressure automatically from a water-tank connected to the bladders at shoulder level. Flight tests showed that abdominal pressure alone was not entirely satisfactory, and much of the venous pooling occurred in the legs. A hydraulic suit to "pressurize" the thighs in addition to the abdomen, using water bladders within an inextensible suit, was worked out by Dr. Bourdillon in 1938, and submitted to the Air Ministry early in 1939.

In 1939 the late Sir Frederick Banting devoted much time and energy to organizing research in Canada on the development of anti-"g" garments, and Franks, working with him, constructed a number of suits on the hydraulic principle. After trials by the R.A.F. which established the effectiveness of this, a practical anti-"g" garment was produced from them by the research staff of the Dunlop Rubber Company, working with Wing Cmdr. Franks and Flying Officer Wilson. The Franks hydraulic suit was used in 1943 and 1944, but has now been discarded because, by the use of air pressure in place of water, effective protection can be achieved with a much less cumbersome garment.

Early in the war an air-pressure suit was developed in Australia by Dr. Cotton, working with the Australian Air Force, and using the centrifuge (McIntyre, 1945) which was set up there for the purpose. Since then the air principle has been very highly developed in America to give a comfortable and light garment which effectively pressurizes the lower limbs during turning in aircraft and enables the pilot to tolerate some 2 to 3 g beyond the point at which he normally blacks out. Development of air-pressure suits in the R.A.F. has followed on the principles worked out in America. The air-pressure suit is supplied from a pump on the aircraft engine via a valve operated by a weight acting against a spring, so that air is admitted only during positive acceleration in tight turns.

The blood below the heart is at a pressure which increases with the distance below the heart, and it was first thought neces-

often be increased by elucidation of function under abnormal stresses. One general truth is emphasized in this work—that is, the very great factor of safety in many physiological processes.

REFERENCES

- de Haven, H. (1942). *War Med.*, 2, 586.
Denny-Brown, D., and Russel, W. R. (1941). *Brain*, 64, 93.
Franks, W. R., Kerr, W. K., and Rose, B. (1945). *Proc. physiol. Soc.* (In press.)
Greenfield, A. D. M. (1945). *Ibid.* (In press.)
Holbourn, A. H. S. (1943). *Lancet*, 2, 438.
Jasper, H. H., and Cipriani, A. J. (1945). *Proc. physiol. Soc.* (In press.)
Livingston, P. C. (1939). *Brit. J. Surg.*, 26, 749.
McIntyre, A. K. (1945). *Proc. physiol. Soc.* (In press.)
Marshall, G. S. (1933). *J. roy. aero. Soc.*, 37, 389.
Rook, A. F., and Dawson, D. J. (1938). *Lancet*, 2, 1503.
Stewart, W. K. (1945). *Proc. physiol. Soc.* (In press.)
von Diringshofen, H. (1934). *Z. Biol.*, 95, 1, 551.

PERIPHERAL ARTERIAL EMBOLLECTOMY

BY

PHILIP HOPKINS, M.R.C.S.

Surgical Registrar, King George Hospital, Ilford

The purpose of this paper is to review the literature on embol-ectomy, with special reference to the results; to stress the urgency of early diagnosis and prompt surgical treatment; and to describe a case in which a popliteal embol-ectomy was attended by success even though the time lapse between the occurrence of the embolism and the operation was greater than any other previously recorded.

Aetiology

The emboli we are concerned with are mainly derived from ante-mortem clot formed in the left side of the heart when stasis is present—chiefly in mitral stenosis. Occasionally the clot may be formed in the pulmonary or systemic veins; in the latter case the so-called “paradoxical embolus” occurs, being due to a persistent foramen ovale. But, nevertheless, cardiac dysfunction is the commonest cause: according to Danzis (1933) 60% of cases are due to cardiac disease, while of the 47 peripheral emboli recorded in Great Britain (in the available literature) 40 were of cardiac origin. Three of these occurred in thyrotoxic subjects with auricular fibrillation; the rest were in patients with mitral stenosis—the commonest single cause of embolism. Of the remaining 7, 6 were post-operative and one followed an accident which resulted in a fractured rib.

History of Arterial Embol-ectomy

Suabanejew, in 1895 (cited by Key, 1923), was the first to attempt “arteriotomy in order to remove a clot.” He was unsuccessful, as were the next 11 surgeons who followed his example. Among these were Moynihan (1903; not recorded until 1907) and Handley (1907), in England; Stewart (1905; cited by Allen, 1929), Murphy (1909), and Carrel (1909; cited by Allen, 1929), in America; and Lejars (1902; Allen, 1929), in France.

The first recorded success was described by Mosney and Dumont (1911), when Georges Labey removed an embolus from the bifurcation of the left femoral artery six hours after its lodgment. The patient, a man of 38, had mitral stenosis, and made a good recovery. In the same year Gordon-Watson, in England, attempted to remove a post-operative embolus under similar conditions. The leg, however, had to be amputated later, but the patient survived.

Einar Key, of Stockholm, who was responsible for the second reported successful case, introduced the operation in Scandinavia, after Carrel's experimental work had made arterial surgery a practical proposition (Allen, 1929). By 1923 Key was able to report 12 further successes, mainly by Scandinavian surgeons. In 1923 Holst (Danzis, 1933) reported the first success in America, while it was not until 1925 that Jefferson described the first success in England. In 1928 Pemberton collected 216 cases of embol-ectomy, of which only 20 had been done in Canada and America, while 145 of them were Scandinavian. Twenty cases had been reported in England by Riddell (1935), but by this time there were 382 cases on record for Scandinavia alone. It is evident that the Scandinavian surgeons had accepted embol-ectomy as a useful operation, while the general practitioners in that country were co-operating by sending

patients to hospital as soon as the diagnosis of peripheral embolism was suspected.

At the present time only 41 cases have been recorded in England; 47 emboli occurred in these, and embol-ectomy was carried out for 42 of them. The Registrar-General has, for the years 1941–3 alone, details of 34 deaths in which peripheral embolism played a part (personal communication), so there is no doubt that more than the 41 cases recorded have occurred in England since the time of Moynihan's first attempt.

Results

The main factor in the recovery rate of the limb in embol-ectomy is time—the time allowed to lapse between the lodgment of the embolus and the attempt to remove it. Pearse (1933) and Danzis (1933) have shown this to be true in their reviews of relatively large numbers of cases, which are expressed diagrammatically in Figs. 1 and 2.

The total successes reported in the past tend to give rather a gloomy impression (see Table), but on more careful examina-

Table showing Recovery Rates as found by Various Observers

Observer	Year	No. of Embol-ectomies	Successes	
			No.	%
Jefferson	1925	73	28	34.2
Danzis	1933	119	49	41.1
Pearse	1933	296	145	49.7
Key	1936	382	86	22.5
Hopkins	1945	42*	24	57.1

* This figure includes the 20 cases collected by Riddell (1935).

tion it becomes apparent that if *all* cases could be treated within 4 to 6 hours of the impaction of the embolus in the artery, then the total number of recoveries might be very much greater. Dickson Wright (1933) in a postgraduate lecture stated that there is a 50% chance of recovery after embol-ectomy. Without operation the mortality rate in patients who have a peripheral embolus has been put as high as 87% (Danzis, 1933), with the majority of the remaining 13% having to undergo amputation of the affected limb.

The ultimate prognosis, of course, is that of the causative disease.

FIG. 1.—After Pearse (1933). 296 cases.

FIG. 2.—After Danzis (1933). 119 cases.

Successful cases have been reported in which more than 24 hours have elapsed between the embolism and the operation; the present case was not admitted to hospital until 39 hours' delay had occurred, and it is very interesting to note that a limb is capable of recovering even after this period of complete anoxia, provided that a good circulation can be restored. Holst (cited by Danzis, 1933) was successful with his case after 27 hours' delay, and Girdlestone's (1935) case recovered after 24 hours. It is apparent, therefore, that operation should be undertaken even when an unduly long delay has occurred, although it must still be emphasized that the optimum time for embol-ectomy is within 4 to 6 hours of the lodgment of the embolus. This consequently means that early diagnosis is essential.

Diagnosis

The leading factor in the diagnosis of peripheral embolism is that *something sudden happens in a limb*. Often a patient will liken his illness to a “stroke,” and almost always he will be able to give the exact time of its onset. The most commonly described symptoms are pain and paralysis in the affected limb, but these are not constant. The pain may be agonizing; like a burning sensation; or merely be “pins and needles” or other