A Method of End-to-end Anastomosis in the Small Intestine

In lesions of the small intestine requiring resection speed is of very considerable importance in war surgery, and side-to-side anastomosis is therefore seldom done because, while giving a very satisfactory union, it is undoubtedly time-consuming. The common method of end-to-end anastomosis is to unite the cut ends with two layers of sutures—an inner "all-coat" and an outer "sero-muscular"—but this method has two disadvantages. In the first place it is often found that closure of the mesenteric angle gives rise to some difficulty, and at the end of this part of the anastomosis there is sometimes a feeling that one's technique at this stage has not been quite up to standard. Indeed, this fact has lately been uncomfortably proved at the necropsy of a patient dying of paralytic ileus from peritonitis, where a small localized abscess was found within the mesenteric angle. The second disadvantage is that the two layers of sutures produce a thick and rigid anastomotic line predisposing to mechanical obstruction.

The following method of anastomosis has been devised to overcome these difficulties and to reduce the operative time. The diagrams explain the stages of the operation.

METHOD

Having determined the line of bowel resection, the mesentery of the efferent loop is divided at this level, but with the afferent loop the mesentery is divided 1 in. proximal to this line, thus leaving 1 in. of bowel without mesenteric attachment (Fig. 1). Next, three cat-gut (00) sutures are introduced from without inwards into the efferent gut. These sutures are inserted 1 in. distal to the line of resection and are placed equidistantly around the circumference of the gut, with one suture at the antimesenteric border. Each stitch is then brought out through the end of the efferent loop, the corresponding part of the end of the afferent loop is picked up, and the stitch returns within the lumen of the efferent loop again, to emerge close beside its point of entry, where the ends are tied (Figs. 2 and 3). Thus the afferent loop is invaginated within the efferent loop and the cut ends of the mesentery are now in apposition. A continuous suture, picking up all coats of the cut end of the efferent loop, and the sero-muscular coats of the afferent loop at the level of mesenteric attachment, now completes the anastomosis, and the mesenteries are united by interrupted stitches (Fig. 4).

COMMENT

The advantages of this method are that it is quick and easy to perform and makes leakage at the mesenteric angle impossible. In addition it is felt that it offers less resistance to normal peristaltic movements along the gut.

Two criticisms may be anticipated: (1) Bleeding at the cut end of the afferent loop has not been controlled. In practice, noticeable bleeding has not occurred here, probably because of the distance below the mesenteric attachment at which the gut is divided. (2) The passage of the three interrupted sutures within the lumen of the gut might be considered to spread infection. This obviously is of no importance in war surgery, where gross soiling is inevitably present; and even for "clean" resections this soiling can be of little importance until we more nearly approach the ultimate goal of aseptic anastomosis.

Excluding patients who have died within 24 hours of operation from the severity of their wounds, this method has now been employed in 6 cases, 5 of whom survived. The one death occurred on the 4th post-operative day from gas gangrene of

the abdominal wall, and necropsy showed no intraperitoneal trouble and a sound anastomosis.

Access to the literature is impossible at present, but it is felt that an operation of this type has possibly been described before. Personal ignorance of the existence of such a technique has, however, prompted the description of this method of anastomosis.

J. SETON PRINGLE, F.R.C.S.I. & Eng.,
Major, R.A.M.C.

Two Cases of Air Embolism

In view of recent references to air embolism the following two cases may be of interest.

CASE I

Mrs. L., aged 30, suffering from tuberculous infiltration of the right apex. Left lung healthy. Has had an artificial pneumothorax since March, 1944. Collapse satisfactory and well maintained; patient making good general progress.

After screening, a refill was thought necessary, and at 11 o'clock one morning a needle was inserted in the fifth intercostal space in the anterior axillary line, as was customary with this patient. Pressure was -2, -4 and manometer movement was sluggish; a probe was used to make sure that the needle was clear, but readings persisted. A little air, less than 50 c.cm., was blown through to clear the needle. Patient remarked, "Something funny is happening, Doctor. I think I am going to faint." The needle was withdrawn forthwith. Mrs. L. then became unconscious with eyes wide open and stertorous breathing. Pupils were widely dilated; pulse almost imperceptible though colour remained good. A fine tremor was present throughout the whole body. About ten minutes later the pulse was quite imperceptible and paresis of the left side of the face was noticed. Two c.cm. of nikethamide solution was given intravenously. Dilatation of pupils decreased somewhat, the eyes closed and resisted opening; conjunctival reflex present. Tremor had ceased. The patient was put to bed between two hot blankets and with numerous hot-water bottles. About an hour and a half later she showed signs of returning consciousness and drank a few mouthfuls of tea. She relapsed again, and at 4.30 p.m. there was another fit of trembling with marked photophobia. A quarter of a grain of morphine was given. At 6 p.m. patient was conscious and rational but complained of severe post-orbital and frontal headache. She said that she was blind and could see nothing but a white sheet. Pupils were half dilated and she was obviously unable to see.

Twelve hours later, after a restful night, patient felt well and her vision had improved to the stage where she could see objects, but she complained of their being fuzzy. Optic disks showed a degree of venous congestion and arteries appeared to be contracted. Nothing abnormal was discovered in the nervous system. Thirty hours later she had returned to normal except for a residual fuzziness of vision. Next day this had disappeared. There have been no apparent sequelae. The question of a possible hysteria arose and was considered, but the patient has shown no antipathy to treatment and has been twice refilled since without untoward effect.

CASE II

J. M., aged 19. Extensive tuberculous disease with cavitation in right lung. Artificial pneumothorax induced Sept. 20, 1944, and refilled at regular intervals. On the last occasion, one month after the above case, on the insertion of the needle for refill, before any air was passed and while a reading of -2.0 was being recorded, he suddenly complained that he felt dizzy. The needle was immediately withdrawn and the tip was found to be bloodstained. This patient did not lose consciousness, but he too complained of loss of vision. His pupils, particularly the left, were somewhat dilated; examination of the left disk showed in at least two arteries a short, moving, silver thread which was taken to be air; this disappeared in a few seconds, and perhaps fifteen seconds later the patient remarked, "I can see again now, but I still feel dizzy." After a few minutes' rest he rose from the table unaided, and has had no more trouble other than a slight staining of the sputum about half an hour after the incident.

COMMENT

Hamilton and Rothstein (*J.A.M.A.*, June 22, 1935) consider that air embolism occurs once in 500-1,000 cases, and that it is more frequent in patients whose pleura is thickened and in inductions. It was noticed in both the above cases that the needle seemed to pass through a thickened soft pleura. This has been confirmed in the case of Mrs. L. on later occasions, when refills have been carried out without incident. More satisfactory pressure readings have been obtained on these occasions.

The similarity of the condition to so-called "pleural shock" is noted. It was suspected after the first case that the true mechanism was air embolism, and the opportunity taken for immediate retinoscopy in the second case seems to confirm this.

These cases emphasize the points made by Hamilton and Rothstein: that the common practice of "blowing a little air through the needle to clear it" is one not unaccompanied by danger and is in fact to be firmly discouraged; and, secondly, that the tubing from the apparatus to the needle should be as short as possible and of fine bore. In Case II this could have been the only source of air.

Noranside Sanatorium,
Angus.

PHILIP H. DALGLEISH, M.B., Ch.B.