LONGEVITY AND OLD AGE IN THE PUNJAB

BY

Khan Bahadur M. YACOB, M.B., B.S., D.P.H., D.T.M.&H.

Assistant Director of Public Health, Punjab

AND

SATYA SWAROOP, M.A.

Statistical Officer, Public Health Department, Punjab

(ABRIDGED)

To the medical profession hitherto the problem of old age has presented its chief interest in connexion with the methods of prolonging human life, such as Voronoff's technique of grafting slices of simian testes in the testes of man, Steinach's advocacy of vasectomy, and the use of testicular and ovarian extracts. From the eugenic, social, ecological, and other points of view, however, the question of longevity raises some serious and complex problems. For example, after a certain age, when man's productive capacity dwindles, he becomes a dependant and remains so until death. India is already supporting a large number of dependants—invalids, beggars, widows, and others—and might well ask whether an attempt to prolong human life and thus increase the proportion of the aged would not add to the heavy burden of her poor working population. In a leading article in the *British Medical Journal*¹ some interesting comments were made on ageing and the care of the aged, and the White Paper on Social Insurance was quoted to the effect that in Great Britain between 1965 and 1975 there will be an increase of about 800,000 in the pensioner population and a decrease of more than one million in the contributing population, supposing the present decline in the birth and death rates to continue. In India the question of pensioners is not yet important, but that of dependancy is present in many forms.

The Limit of Human Life

The limit given by the Psalmist to human life is often much exceeded both in the East and in the West. In India the Vedas [Brahman sacred writings] enjoined men to live 100 years, and this age is known to have been reached fairly often in the Punjab as well as elsewhere. There has been some speculation in scientific literature about the ideal duration of life. Rolleston² quotes Hufeland as laying down the law that animals generally live about eight times as long as they take to reach maturity, and Hufeland's figure of 200 years as the ideal span of life is obtained by regarding 25 as the termination of adolescence. Both figures—8 and 25—appear to be arbitrary. Flourens calculated that, throughout the animal kingdom, life was five times as long as the period taken to complete growth, as marked by the union of the epiphyses of the long bones, and as this union took place at 20 or 21 the span of human life should be about 100.

It is of interest to recall some extreme claims for longevity, admitting, of course, a considerable lack of accuracy in age records. Rolleston² quotes the cases, among others, of Henry Jenkins, who lived from 1501 to 1670; Thomas Parr, who lived from 1483 to 1635; and Katherine, Countess of Desmond, who lived 145 years, though some have said that the ages of two Countesses of Desmond were telescoped into one. The question of the longest life to be reasonably attested is of interest to actuaries, and in an American actuarial publication Dublin and Lotka³ cite the case of C. J. Drakenberg, a Danish seafarer, who was born in 1626 and died in 1772. It is declared that he was sturdy and robust and temperamentally very impetuous. At the age of 111 he married a 60-year-old widow, who died a few years later, and at the age of 130 he proposed to several women but without success.

In India an interesting case was recorded in the 1931 census report⁴ of one Siddi Wastad, the son of a negro father and Arab mother, who lived in Nagpur. At the time of the census he was supposed to have passed his 150th year, but the census superintendent for the Central Provinces stated that, even allow-

ing for much exaggeration, his age would then have been 130 years. We addressed an inquiry to Dr. R. L. Tuli, Director of Public Health, Central Provinces, who has informed us that Siddi Wastad died at the age of 158 years on August 14, 1943. He was a famous wrestler at the court of the Maharajah of Baroda. Until the influenza epidemic of 1918 he remained in perfect health, but he was attacked by that scourge and gradually lost his teeth and his hair turned white. Until then it had not been his habit to lie down or go to bed. A short nap in a chair at night was all that he needed, and he offered his prayers five times a day. His diet up to that time was six pounds of flour for one meal, as well as vegetables, rice, mutton, and four seers⁵ of milk each day. When ill he never took drugs, but he prepared his own medicines and medicated oils and treated himself.

Outstanding Age Records in the Punjab

We sent a request to all the district and municipal medical officers of health in the Punjab for recent information concerning long-lived persons in the Province, and some very interesting replies have been received, enabling us to state that this Province has a good number of persons who have passed the century. We further requested the officers to throw light on the state of health enjoyed by persons who had lived to advanced ages. It is not possible to comment in detail on the information supplied; in some cases further inquiry seemed necessary. Only a few salient points may be briefly mentioned.

The maximum age so far reported to us as having been reached in recent times in the Punjab is 138 years. This was the age of a person named Khan Dadan Khan, who died of fever four years ago in the Rawalpindi district, a hilly part of the Province. The death also occurred last year of Khanam Jee, in Rawalpindi city, at the age of 135 years. Two more deaths each at the age of 130 years have been reported from Lahore and Lyallpur districts. An interesting account has been received of one Saleh Mohammed Khokhar, who died recently at the age of 120 years in the Rawalpindi district. The district medical officer of health reports of this man that his grey hair had grown black and that new teeth had appeared. He was accustomed to eat maize, meat, and leafy vegetables, and he never drank tea. Quite a number of other instances have been reported to us of persons who have died recently at the ages of 110 and 130. Generally the cause of death has been stated to be "fever" or "old age," both vague terms. Without further inquiry nothing can be stated as to the precise cause, though we have little reason to doubt that malaria has been an important one.

Among the living in the Punjab a person is reported to have reached 130 years of age in Sialkot, 70 miles from Lahore. His state of health is said to be good. A dhobi [low-caste Hindu] 125 years of age is reported to be living in Amritsar. He attributes his longevity to his habit of taking the gum of the pipal tree [the sacred fig]. Two women of 120 years are reported—one in Ambala, near Simla, in a satisfactory state of health, and the other in a village in the Gujrat district, who is blind, deaf, and bedridden. The number of persons who claim to be over 100 years of age is fairly large. Most of these living persons claim that their longevity is due to the inclusion of ghee [clarified butter], milk, and dairy products in their diet. Some have been taking a few seers of *lassi* daily during both summer and winter. Abstinence from tea and tobacco, moderation in sexual life, and plenty of exercise when young are some of the other factors which these persons claim to have contributed to their longevity. Among the diseases from which they suffer in their old age malaria, eye troubles, bronchitis, and pneumonia are mentioned, but these, of course, are just those affections which constitute the recognized general causes of ill-health in the Punjab.

Expectation of Life in the Punjab

[The authors then turn from the discussion of a few cases of exceptional longevity to the length of life of the population of the Province as a whole, and they explain what is meant by the phrase "expectation of life" as used in census returns and by insurance corporations. The estimate is made by actuaries for each age and shows, on the average, the added number of years for which people of that age are likely to survive. They state that there are some

⁵ The Government seer in India is just over 2 lb. avoirdupois.

¹ *British Medical Journal*, 1944, 2, 634.

² Rolleston, H., *Medical Aspects of Old Age*, 1932, Macmillan.

³ Dublin, L. I., and Lotka, A. J., *Length of Life. A Study of the Life Table*, 1936, Ronald Press, New York.

⁴ *Census of India*, 1931, 1, 106.

objections to the use of expectation-of-life figures relating to India for making long-term comparisons, but for the immediate purpose of studying longevity in the Punjab such estimates may be regarded as fairly useful.]

The latest estimates of expectation of life in the Punjab, as in the rest of India, followed the census enumeration of 1931.

EXPECTATION OF LIFE AT BIRTH FOR MALES IN CERTAIN COUNTRIES.

CHART I

The war has stood in the way of obtaining more recent estimates. We have obtained figures for a number of countries corresponding to the 1931 estimates for the Punjab and for British India, and these are presented in Chart I, the countries being arranged in decreasing order of expectation of life at birth for the male inhabitants. The Netherlands are at the head of the list with as high an expectation of life as 65.1 years, and New Zealand and Australia rank next. British India is lowest in the chart, with 26.9 years, and the Punjab is only slightly higher (28.1). Generally speaking, all the advanced countries of the world, especially those in the West, have an expectation of life far in excess of the Punjab and British India. Among Eastern countries, the figure for Japan (44.8) is interesting; it is more than one and a half times the British India figure.

Within India itself the Provinces show variability, though not as large as that seen in the comparison of countries. Chart II gives the figures for the Provinces, and for males and females in each Province. The Provinces are here arranged according to an increasing expectation of life for females. The highest estimates are for Madras, where the females show a figure of 30.04 years (the only instance in which 30 years is reached). The lowest estimates are for the area adjoining Bengal, comprising the North-West Frontier Province, Sind, and Baluchistan. The Punjab figures are exceeded by those of Bihar and Orissa, the Central Provinces, Berar, and Hyderabad, and Madras.

Lower Female Longevity

It is almost a universal experience that females are longer-lived than males, but the Punjab affords a striking exception, as also do the North-West Frontier Province, Sind, and Baluchistan, Bombay Presidency, and Bihar and Orissa. Probably the lack of adequate facilities for maternal relief and care,

coupled also perhaps with a rather unfavourable attitude towards the weaker sex in the Punjab, are responsible for keeping female longevity at so low a level. In making the statement about an unfavourable attitude towards females we are supported by the great amount of evidence pointing to the practice of female infanticide in this Province in the past.

The attention of the British Government was directed to this subject immediately after the annexation [1849]. Lord Lawrence, while Commissioner of Trans-Sutlej States, promulgated three commandments, the second of which was, "Thou shalt not kill thy daughter." But writing in 1895 Sir James Douie recorded that "fifty years after Lawrence denounced the murder of female infants we are still discussing the best method of putting down that inhuman practice." Several causes tended gradually to restrict the extent of female infanticide, but while in recent years the killing of female infants may have altogether disappeared, the social customs and mental traits which found expression in such inhuman action may still exist among certain classes, with the result that females generally may lead a more neglected life than males.

A low expectation of life for females, which implies a higher mortality rate among them, affects indirectly the male population, because a large number of women die as the result of pregnancy and childbirth, which involves the potential loss of male as well as female children.

In order to illustrate the point that lack of maternity facilities, together with, perhaps, a general neglect of the female population, has brought about reduced longevity for females in the Punjab as compared with males, we quote the mortality statistics for the sexes at different age periods. The figures set out in Table I are based on the average experience

EXPECTATION OF LIFE AT BIRTH FOR MALES & FEMALES IN 1931 BY PROVINCES.

CHART II

of the quinquennium 1935-9, and the death rates are given per 1,000 of the corresponding populations (except that in the first age group the mortality rates have been calculated per 1,000 live births).

Female infants show a lower mortality than male infants; this is also true of the next age group. But from the age group 10-15 years onwards the female population must be considered to have started the reproductive span of life, and

they are found subjected to adverse mortality rates. The difference persists throughout the reproductive period—namely, up to about 40 years. In the ensuing two age groups the difference between male and female mortality rates becomes

TABLE I.—*Death Rates per 1,000*

Age Group	Males	Females
Under 1 year	172	157
1-5 years	37	34
5-10 "	9	9
10-15 "	7	8
15-20 "	8	10
20-30 "	7	9
30-40 "	9	11
40-50 "	14	15
50-60 "	24	25
60 and upwards	61	64

less, though the females still have the higher rate, and beyond 60 the difference again increases, to the disadvantage of the female population. On the whole, therefore, this table presents the picture of a female population born stronger than the male, but subjected during the prime of life to adverse circumstances, and continuing to live thereafter the life of a weaker sex, in the real sense of the word.

Further Deductions from Tables of Mortality

Mr. L. S. Vaidyanathan, the actuary appointed by the Government of India at the 1931 census, studied the same point—namely, the comparative mortality figures for each year of age.⁶ He did not believe that the recorded births and deaths afforded reliable information concerning mortality conditions in the population, and he proceeded to calculate mortality rates by making use of the indirect evidence available in the 1921 and 1931 census enumerations only. He found that the females started life with a relatively low mortality rate, and maintained this advantage for a few years. Thereafter the female mortality reached a higher level than the male, and this relative disadvantage continued until about the age of 50. After that, according to Mr. Vaidyanathan's calculations, the females scored over the males again and showed low mortality experience almost to the end of life.

Our own deductions from the recorded deaths in the Punjab agree in the main with Mr. Vaidyanathan's differently derived deductions with respect to women in the prime of life. We differ from him concerning the relative position of the male and female at advanced ages. The census enumeration figures of 1921 and 1931, from which he sought to deduce death rates by age and sex, were by no means entirely reliable. We are therefore unable to decide which estimate is subject to the greater error and which must accordingly be preferred.

Mr. Vaidyanathan is an eminent Indian actuary, and in this discussion of longevity and old age we would quote further from his figures. In Table II are set out the figures for the expectation of life for each decennium from 50 to 90 years. They are shown separately for males and females for the Punjab, for the whole of India, and for England and Wales, and they relate to the year 1931.

TABLE II.—*Comparative Expectation of Life at Decennial Ages*

Age	Males			Females		
	Punjab	India	England	Punjab	India	England
50	15.09	14.31	21.36	16.14	14.65	23.69
60	10.22	10.25	14.36	11.48	10.81	16.22
70	5.98	6.23	8.75	7.03	6.74	9.95
80	2.91	3.13	4.93	3.43	3.25	5.56
90	1.19	1.12	2.82	1.31	1.18	3.13

The table affords many interesting comparisons. At the higher ages, in India as well as in England, females enjoy a longer expectation of life. The reason usually given for this generally favourable position of older women is the fact that they have already passed the dangerous period of their reproductive life. From a biological point of view here is the evidence that, once the period of excessive maternal mortality is over, women again present a picture of a more favourable

make-up for survival. This argument also points the necessity for making maternity in the Punjab as easy and risk-free a process as is practicable.

While in England the rate of maternal mortality is as low as 2 or 3 for 1,000 births, in the Punjab it is generally believed to be from eight to ten times this figure.

Survival at Late Ages

Another point revealed by a study of Table II is that, although the figures showing the expectation of life at the higher ages are greater in England than in the Punjab, the difference is not as great as that between the figures for the two countries for the expectation of life at birth. Once, therefore, a person in the Punjab has reached an advanced age, he finds himself in a somewhat more favourable position, and may look forward to living to that greater age which his contemporaries reach in the more advanced countries of the world. At the age of 60, for instance, a male in the Punjab can on the average expect to live another ten years, while a similar expectation in England is about 14 years. Females at the age of 60 in the Punjab may hope to live another 11 years, and in England another 16 years. At the advanced ages females in the Punjab enjoy a longer expectation of life than the average for all India, but Punjabi males show no such appreciable advantage. In fact, at the ages 60, 70, and 80 the expectation of life for males in the Punjab is somewhat lower than that for males in India as a whole.

The smallest area within India for which separate figures for the expectation of life have been worked out is the province; but in order to study the variations existing within the Punjab itself we have calculated the proportion of persons above 60 years of age in the different districts to the total population.

The highest proportion of old persons to the total population is to be found in the hilly or sub-montane tracts of eastern and northern Punjab. In fact, four fairly well marked regions are seen as we proceed south and south-west. The lowest proportion is found in the dry and hot districts of the extreme south-east and south-west of the Province—Karnal and Rhotak on the one side and Muzaffargarh on the other. In a general way this geographical pattern lends support to the view that a cooler climate is conducive to the preservation of life. For example, Kangra, a district of 10,000 square miles in the north-east of the Province, although it has a relatively higher number of old people, is not a satisfactory area from the public health and nutritional points of view. The incidence of tuberculosis, pneumonia, dysentery, and diarrhoea is comparatively high, and goitre, rickets, and osteomalacia are peculiar to this area. Yet in spite of such unfavourable circumstances the population includes a higher proportion of older people, and we are led to the conclusion either that the population statistics are entirely unreliable or that the colder climate is a governing factor in this relative longevity.

Changes in Mortality Conditions

When is a person old? In other words, what age should we choose for a study of mortality conditions of old persons in the Punjab and the manner in which these have changed during, let us say, the present century? It is generally believed that when the period of reproduction wanes that of old age begins. In women this is marked by the menopause; in men the onset of senescence may occur at considerably varying periods and be so stealthy as to be undetected by the individual. A philosopher has said that the information that he is old may be suddenly conveyed to a person on overhearing the chance remarks of others, by catching the reflection of his bent back in a mirror, or by finding some girl offering him her seat in an omnibus. But statisticians and biologists have given ages ranging from 45 to 60 as indicating the onset of old age in man.

Vital statistics for the Punjab give mortality figures for ages over 60 in only one group. All we can study, therefore, is the group of persons aged 60 and above in a single combined estimate. The deaths occurring during each year from 1901 onwards in males and females over 60 have been made into rates per 1,000 of population in that particular sex group and are shown in Chart III for each year. The dotted curve is for the females and the continuous line for the males. In spite of considerable fluctuations occasioned by epidemics of

⁶ Vaidyanathan, L. S., Actuarial Report annexed to Census of India, 1932, Vol. I.

plague, cholera, malaria, and influenza, the general indication is that from 1900 onwards the mortality rate among old male and female populations have been on the decrease. Separate trend lines have been fitted by the method of least squares to the mortality experience of the sexes over the entire period of 41 years (1900-40), and these are shown on the chart. Both these lines are sloping downwards, which is an indication of a general improvement in the health conditions of the old. The significance of this decline has been tested statistically, and it has been found that chance alone cannot explain such a marked downward tendency.

**TRENDS OF DEATH RATES AMONG MALES & FEMALES
OVER 60 YEARS OF AGE IN THE PUNJAB FROM
1900 ONWARDS.**

CHART III

We conclude that, generally speaking, there has been during the present century a growing improvement in the condition of old persons as shown by mortality rates—an improvement somewhat more marked among the females.

Long-term Comparisons

Another method sometimes adopted of studying long-term changes in longevity is by comparing figures for expectation of life at decennial intervals, though we have grave doubts whether the Punjab or, for the matter of that, the all-India figures can be quoted with the same degree of confidence as the figures for Western countries.

TABLE III.—*Expectation of Life in Years for Males at Ten-yearly Age Intervals from 1881 to 1931*

Age	1881	1891	1901	1911	1931
0	25	27	23	21	28
10	36	38	35	31	38
20	30	32	30	26	31
30	25	27	25	22	25
40	20	20	20	18	20
50	15	16	15	14	15
60	10	11	11	11	10
70	6	8	6	7	6
80	3	5	3	3	3
90	1	2	1	1	1

Several factors tend to detract from the value of these figures for long-term comparisons. (1) The figures for 1921 are missing, as the actuary did not consider that they would be of any use. At that time the population had not fully recovered from the effects of the great influenza pandemic of 1918. (2) The estimates after each census were made by different actuaries who used varying methods of estimation. (3) The areas for which the estimates have been made have not always been the same. In 1931 the area included not only the Punjab but also the adjoining smaller Indian States, together with Kashmir, the biggest in area of the States, and noted for its highly salubrious and temperate climate. (4) There is some doubt, particularly for the figures for 1901, as to the precise year or period to which the estimates relate.

It is not possible to say how much allowance has to be made for these factors. However, if the broad indications have any significance it can be stated that the degree of improvement noticed at the lower ages between 1911 and 1931 has not been maintained at the advanced years of life. The 1931 census estimates were arrived at without any regard to the recorded death statistics. On a survey of the mortality rates among the aged in the Punjab during the period 1900-40 we have, however, demonstrated already a downward and statistically significant tendency. We therefore conclude that there has been some improvement in the mortality conditions of old people, and that females have had a somewhat greater advantage.

Summary and Conclusions

The present position of longevity in the Punjab has been studied in relation to other countries and different Provinces of British India. Some relevant information regarding remarkably long-lived persons has been set out, together with figures for the expectation of life for different countries and Provinces. An investigation has also been carried out separately for the two sexes.

As compared with the advanced countries of the world, the Punjab has pitifully low figures of longevity. Further, contrary to the general experience that females have a longer duration of life than males, the women in the Punjab, having been subjected to relatively more adverse circumstances, have not been able to enjoy that duration of life with which Nature equips them. This means the potential loss of male as well as of female children. The study indicates that one of the urgent needs is the making of reproduction an easy and risk-free process by all possible means, not excluding a general social improvement of the female population. Improvements in these directions will be doubly beneficial in that not only will there be a saving of the lives of mothers but the chances of survival of their offspring will increase.

It has been shown that some improvement has been occurring in the mortality rates at the advanced ages of both males and females.

MANCHESTER'S VITAL STATISTICS

Manchester in 1944 had the highest recorded birth rate for twenty years, the lowest general death rate for any year of the war period, and the lowest infantile mortality and puerperal causes mortality on record. Comparing the Manchester of seventy years ago with the Manchester of to-day, both the birth rate and the death rate have been almost precisely halved; the marriage rate has been considerably higher during the war years than during the previous sixty years, and the birth rate, although nothing like what it was before the war of 1914-18, has gone up steadily in each war year. Sixty years ago in Manchester the death rate from measles was 0.71 for every 1,000 persons living, it is now 0.02; from diphtheria it was 0.10, and is now 0.01; from whooping-cough it was 0.68, and is now 0.04; from pulmonary tuberculosis it was 2.42, and is now 0.80; from other diseases of the respiratory system it was 5.41, and is now 2.04; and from diseases of the digestive system it was 1.23, and is now 0.58. On the other hand, the cancer mortality has gone up from 0.50 to 2.09, and the mortality from diseases of the heart from 1.37 to 3.17. Nevertheless, the death rate in Manchester is higher than in England and Wales, and the infantile death rate is 54 per 1,000 births as against 46 for the whole country. But Manchester is in the proud position, not for the first time by any means, of having a maternal mortality rate distinctly lower than for England and Wales as a whole. In 1944 it was 1.83 per 1,000 births, the first time it has ever been below 2.0.