Electromyography

There is another interesting alteration in electrical activity which is of great significance in diagnosis; but, unlike electrical reactions, it can be of considerable prognostic importance. The action currents of contracting muscles can be recorded—the electrocardiogram is a well-known example. By electromyography it is possible to record the action currents in a voluntary muscle.

For about 12 to 28 days after denervation there is no recordable electrical activity in human muscle on attempted movement or at rest. From 12 to 28 days after denervation fibrillation occurs: this is shown by rhythmic fine spikes totally unrelated to attempted voluntary contraction. The exposed surface of such a muscle if observed by reflected light will show numerous small scattered twitches. Fibrillation persists as long as any contractile denervated fibres remain, or until re-innervation is taking place, but in the early stages of re-innervation a muscle which has previously shown vigorous fibrillation becomes silent. Motor-unit action potentials may precede recovery of voluntary function by as much as three months, and an isolated observation of this sort is no guarantee that further re-innervation is going to occur.

The finding of both fibrillation and motor-unit action potentials indicates a partial or recovering degenerative lesion of the lower motor neurone (Weddell, Feinstein, and Pattle, 1943, 1944; Bowden, unpublished).

Treatment

We have seen that electrical reactions and electromyography may be of help in diagnosis and to some extent in prognosis. The histological findings have shown that in the muscle itself atrophy of the muscle fibres and connective-tissue proliferation are the enemies of recovery. Surgical repair, if necessary, should be undertaken as soon as possible. There is experimental and clinical evidence that atrophy of muscle can be retarded by activity induced by adequate electrotherapy. The contraction of the muscle is essential; therefore faradic stimulation of denervated muscle is useless and a waste of time. Denervated muscle responds to galvanism, and this is therefore one method of treatment.

In a series of rabbits both the left and right tibialis anterior were denervated for the same length of time. One side was treated daily with galvanism, the other was untreated; and there was considerable difference in the naked-eye appearance of the muscles. The histological picture was even more impressive: the diameter of the treated muscle fibres is much greater than that of the untreated side and the formation of fibrous tissue is minimal (Gutmann and Guttmann, 1942, 1944). This is a most convincing proof of the value of regular adequate electrotherapy. Can this experimental finding be applied clinically?

Dr. Shirley Jackson investigated the value of galvanism in a series of cases of ulnar nerve palsies (personal communications; work in press). Volumetric measurements were made at intervals. It was seen that atrophy is most rapid in the early stages of paralysis, and that electrotherapy does not wholly prevent atrophy; it can only retard the rate. At no time was there an increase in volume of the hand in the absence of re-innervation. Therefore it is essential that electrotherapy not only should be regular and adequate but should begin as soon as possible after injury.

Connective-tissue formation is retarded by physiotherapy, but it cannot be entirely prevented. The late sequel of connective-tissue formation is contracture of the paralysed muscle. Whilst contractures can and do form in active unopposed muscles, they may occur in paralysed muscles, either early or late in the course of denervation. They can be wholly prevented by daily movements throughout the whole range of the joints of the limb. Splinting of paralysed limbs is never done to obtain immobilization unless there is a fracture or an unhealed wound; the object is to prevent overstretching and to encourage activity by compensating for the paralysis, if this is possible.

The author is indebted to the British Orthopaedic Association for permission to publish this paper, which was read at the association's meeting on Dec. 15, 1944. The work discussed has been carried out by members of the peripheral nerve injury unit, Department of Orthopaedic Surgery, Oxford, under the direction of Prof. H. J.

Seddon, and in conjunction with members of the Department of Zoology and Comparative Anatomy, and of the Department of Human Anatomy, University of Oxford. A fuller discussion, and bibliography, will be found in the papers quoted.

REFERENCES

- Bowden, R. E. M., and Gutmann, E. (1944). *Brain*, **67**, 273.
 Erb, W. H. (1876). In *Cyclopaedia of the Practice of Medicine*, **11**, H. v. Ziemssen, London.
 Gutmann, E., and Guttmann, L. (1942). *Lancet*, **1**, 169.
 — (1944). *J. Neurol., Neurosurg., Psychiat.*, **7**, 7.
 — Medawar, P. B., and Young, J. Z. (1942). *J. exp. Biol.*, **19**, 14.
 — and Sanders, F. K. (1943). *J. Physiol.*, **101**, 489.
 — and Young, J. Z. (1944). *J. Anat., Lond.*, **78**, 15.
 Seddon, H. J. (1942). *British Medical Journal*, **2**, 237.
 — (1943). *Brain*, **66**, 237.
 — Medawar, P. B., and Smith, H. (1943). *J. Physiol.*, **102**, 191.
 Weddell, G., Feinstein, B., and Pattle, R. E. (1943). *Lancet*, **1**, 236.
 — (1944). *Brain*, **67**, 178.

VALUE OF SUPPLEMENTS OF VITAMIN C IN PREVENTING LACTATIONAL MASTITIS

BY

A. A. FULTON, M.D., D.P.H.

*Maternity and Infant Welfare Medical Officer, Corporation of
Dundee*

One theory put forward for the high incidence of lactational mastitis in an industrial Scottish town (Fulton, 1945) was that a deficiency—probably due to wartime restrictions—in some food factor, or factors, was a predisposing cause. Nixon (1941) has suggested that there is a relationship between dietary deficiencies and puerperal infections, and has cited as evidence the fact that during the last war mortality from puerperal fever in Germany was almost doubled as a result of the blockade. One of the most obvious of wartime food curtailments in this country is shortage of fresh fruits and vegetables, the chief source of vitamin C. Widdowson and Alington (1941), in studying middle-class diets, have shown that in 1935 fruit supplied ten times as much vitamin C as it did in 1941 and that the average vitamin C intake in 1935 was more than double what it was in 1941.

At the time when this experiment was begun (Dec., 1941) the following assumptions could reasonably be made.

- (1) There was a high, if not an increased, incidence of mastitis.
- (2) The incidence of mastitis appeared to be greater in the winter months (Moon and Gilbert, 1935).
- (3) Some degree of vitamin C deficiency was common. Many writers had brought forward evidence to support this view (Harris and Ray, 1935; Abbasy *et al.*, 1935; Orr, 1936; Harris *et al.*, 1936; Portnoy and Wilkinson, 1938; Harris, 1940a; Youmans, 1941).
- (4) Hypovitaminosis C was likely to have increased under wartime conditions. This has since been proved by Widdowson and Alington (1941), Harris (1942), Francis and Wormald (1942), and Widdowson and McCance (1942).
- (5) Hypovitaminosis C is more likely to occur during pregnancy and lactation. There is evidence that vitamin C requirements are increased during these periods (Harris and Abbasy, 1937; Smith, 1939; Neuweiler, 1935).
- (6) There is a seasonal fluctuation in the intake of vitamin C, as fresh fruits and vegetables are more plentiful in the summer and autumn months. Evidence of this has since been published by Harris (1942), McNee and Reid (1942), Harris and Olliver (1943), Prunty and Vass (1944), and Craig *et al.* (1944).
- (7) The time of seasonal increase in the incidence of mastitis (the winter and spring months) roughly corresponds to the period when intakes of vitamin C are lowest.
- (8) Lack of vitamin C diminishes the resistance to infection (Harris, 1937; Smith, 1939; Harris, 1940b; Youmans, 1941).

If these assumptions were accepted there seemed to be some justification for postulating that vitamin C deficiency might be responsible, at least in part, for the occurrence of mastitis. An experiment was devised to attempt to verify or disprove this hypothesis. Although, admittedly, a diet functions as a whole and deficiency of one constituent only is unlikely to occur, it was thought better to investigate the effects of varying the intake of one single factor in order to attempt to obtain a clear-cut result regarding that factor. The project was to give one group of pregnant women supplements of ascorbic acid and to compare them, especially as regards the incidence of mastitis, with a group of women, similar in other respects, who did not receive supplements.

Plan of the Experiment

There are two ante-natal clinics in the town concerned, each serving approximately half the town and attended by patients similar in most respects, including social status. The expectant mothers attending one of the clinics (Clinic A), who had booked to be delivered in the Maternity Hospital, were given a daily supplement of ascorbic acid from the sixth month of pregnancy until the end of the sixth month after delivery—i.e., over a period of nine months covering the later months of pregnancy and almost the whole period of lactation. The supplements consisted of 50-mg. tablets of ascorbic acid, and the patient was requested to take one tablet a day for the nine-months period. The tablets were given out at the ante-natal clinic, and supplies were renewed at subsequent visits, in the Maternity Hospital, at the child welfare clinics, or delivered at the home. The dosage chosen was based on a daily requirement of 30 mg. for an adult as recommended by the Technical Commission on Nutrition (1938), but some surplus (20 mg.) was allowed owing to the increased demands of pregnancy and lactation. The total daily intake was probably in the region of 70 mg., as these women were presumably consuming some vitamin C in their ordinary diet—probably not less than 15 to 20 mg. a day—and the 50-mg. tablet was considered to be in the nature of a supplement to bring what might be an inadequate intake up to a satisfactory level. Graham and Cameron (quoted in the Departmental Report on Infant Mortality in Scotland, 1943) found that in Glasgow the average values of vitamin C in the diets of 300 pregnant women varied from 30 to 42 mg. a day.

The great disadvantage in dealing with out-patients is that it is impossible to be sure that the tablets are actually ingested. Most experiments of this nature have been carried out in hospitals, schools, and other institutions, where the subjects are under direct control and supervision, and where supplements can be added to the diet. In this experiment the difficulty could not be overcome in any way, but tests were carried out on every patient while in the Maternity Hospital to estimate the degree of saturation, and this was very helpful in assessing the previous regularity of ascorbic acid consumption.

The group of patients attending Clinic A, who received supplements of ascorbic acid, will be referred to as Group A, and consisted of 124 women. The patients attending Clinic B did not receive any supplement, and acted as controls; they numbered 126, and will be referred to as Group B. Comparisons between the two groups as regards age, parity, and income levels were made. The average age of Group A patients was 27.30 years, which was very much the same as the average for Group B—namely, 27.62. The average weekly income per household was slightly higher for the Group B patients, being 75s. 10d., as compared with 68s. 9d. for Group A. There was a greater proportion of primiparae in Group B (52.4%) than in Group A (47.6%).

On admission to the Maternity Hospital, where Group A patients continued to have a daily dose of 50 mg. of ascorbic acid, all patients in both groups were examined and particulars taken as to previous diet, with special reference to the consumption of fruit and vegetables, weekly incomes, and number of members in the household. In the case of Group A inquiries were made as to the regularity of the tablet consumption. The vitamin C nutrition of both groups was assessed by means of the saturation test devised by Harris and Abbasy (1937).

In the early part of 1943 the Maternity Hospital had to be closed owing to an outbreak of puerperal infection and, for this and other reasons, 15 women in Group A were delivered at home instead of in the hospital, as had been originally intended. Supplies of ascorbic acid tablets to these women were continued, but it was not possible to carry out saturation tests. As the Group B patients were only "enrolled" on admission to hospital all in this group were tested.

It was found that a fair number in Group A were not saturated for several days, which suggested either that the tablets had not been taken regularly or that the demands were greater than the amount of vitamin supplied.

It is considered by Harris (1943) that subjects who saturate on the second day of test-dosing have a satisfactory intake as judged by the League of Nations standard, and that with daily intakes of 50 to 75 mg. saturation will occur on the first day. On the other hand, when intakes are appreciably below the League of Nations standard, three days or more are required for saturation.

According to this standard, only one-half (50.5%) of the 109 women in Group A who were tested could be considered to have had an adequate intake prior to testing. This state of saturation of the women in Group A was judged to be not entirely satisfactory, but the degree of saturation in Group A was much better than in Group B, in which only 10 women (7.9%) were saturated by the second day. It may also be noted that, whereas 20 (18%) of Group A were unsaturated by the end of the fourth day, 87 (70%) in Group B took more than four days to reach saturation level.

The average amount of ascorbic acid required for saturation was 1,673 mg. for Group A—just a little more than half of the 3,076 mg. required for Group B. These figures may be compared with the average number of milligrammes of ascorbic acid required to

TABLE I.—Number of Days of Test-dosing required to Attain Saturation

Days	Group A	Group B
1	31	3
2	24	7
3	21	19
4	13	10
5	12	24
6	6	32
7	2	22
8	—	6
9	—	2
10	—	—
11	—	—
12	—	—
13	—	*1
Total	109	126

* This patient showed no evidence of scurvy. The gums were healthy, but she was completely edentulous.

saturate members of the staffs (nursing and medical) of two municipal hospitals (1,165 mg. for the staff of one hospital and 1,421 mg. for the staff of the other).

It would appear from the results of the saturation tests that pregnant women require supplements of vitamin C to bring the intake up to a satisfactory level; but Harris (1943) is careful to point out that saturation is not necessarily desirable and that "sub-standard" is not synonymous with "clinically deficient." Although many members of both groups were found to be deficient in vitamin C there was a big enough difference in the saturation levels of the two groups to allow some comparisons to be made.

Effect of Vitamin C Administration on Incidence of Mastitis

The incidence of suppurative mastitis in both groups was very high, and more than double the rate found by a notification scheme elsewhere described (Fulton, 1945). It must be remembered, however, that these patients were very carefully observed and followed up for a period extending to the end of the first six months after delivery, whereas under the notification system referred to cases may have been missed. It was impossible to complete the follow-up of five patients (four in Group A and one in Group B) who left the district before the six-months period had expired and could not be traced, and two in Group B who died. Consideration must also be given to the fact that the great majority (all but the 15 in Group A who were delivered at home) were hospital patients, and most authorities agree that patients delivered in hospital are much more prone to develop mastitis.

Table II shows that the incidence of mastitis, including cases which resolved without abscess formation, was almost equal (30% in Group A and 30.9% in Group B), but there was a slightly greater percentage of suppurative mastitis in Group A (26.7%, as compared with 23.6% in Group B). It would not have been surprising if these figures had been reversed, as Group A was more favourably situated as regards two factors considered to be predisposing causes—viz., primiparity and delivery in hospital.

TABLE II.—Incidence of Mastitis in the Two Groups

	Group A	Group B	Total
Mastitis (suppurative) ..	32 (26.7%)	29 (23.6%)	61 (25.1%)
" (non-suppurative) ..	4 (3.3%)	9 (7.3%)	13 (5.3%)
No mastitis ..	84 (70.0%)	85 (69.1%)	169 (69.5%)
Untraced ..	4	3	7
Total ..	124	126	250
Primiparae ..	59 (47.6%)	66 (52.4%)	
Number confined in hospital	109 (88.0%)	126 (100.0%)	

From these results it would appear that vitamin C administration *per se* has no influence in reducing the incidence of mastitis, and this is borne out when the available information is analysed more fully.

The number of patients with mastitis in whom the state of vitamin C nutrition was estimated by means of the saturation test was 69—57 with abscess formation and 12 in whom the condition subsided.

Mastitis Cases tested by Saturation

It can be seen from Table III that the average amount of ascorbic acid required to saturate the patients with suppurative mastitis was 1,221 mg. in Group A and 3,167 mg. in Group B; the comparable figures for non-suppurative mastitis were 1,600 mg. for Group A and

3,067 mg. for Group B. These may be compared with the average amounts required to saturate all the members of each group.

TABLE III.—Average Amount of Ascorbic Acid in milligrammes required for Mastitis Cases to Attain Saturation

	Suppurative Mastitis	Non-suppurative Mastitis	Average for Whole Group
Group A ..	1,221 (28 cases)	1,600 (3 cases)	1,673 (109 cases)
Group B ..	3,167 (29 ")	3,067 (9 ")	3,076 (126 ")
Both groups ..	2,210 (57 ")	2,700 (12 ")	2,425 (235 ")

As regards the time required to attain saturation of the patients who developed mastitis, 17 (29.8%) of the suppurative cases (all in Group A) and 3 (25%) of the non-suppurative cases (two in Group A and one in Group B) were saturated by the end of the second day—i.e., they could be considered to have a satisfactory intake according to the League of Nations standard. Therefore the attainment of a satisfactory level of saturation by itself did not prevent mastitis in one-quarter of the cases (Table IV).

TABLE IV.—Number of Days of Test-dosing required to Attain Saturation of Mastitis Cases

Days	Group A	Group B	Total
(a) Suppurative Mastitis			
1	13	0	13
2	4	0	4
3	9	0	9
4	1	2	3
5	1	7	8
6	0	6	6
7	0	7	7
8	0	1	1
	28	29	57
(b) Non-suppurative Mastitis			
1	0	0	0
2	2	1	3
3	0	0	0
4	1	1	2
5	0	4	4
6	0	1	1
7	0	2	2
8	0	0	0
	3	9	12

The incidence of suppurative mastitis in 62 patients of both groups who had reached a satisfactory level of saturation by giving a response before the third day was 27.4% (Table V).

From the results of the experiment it would appear that supplements of vitamin C, or a level of saturation denoting a satisfactory intake of this vitamin, had no beneficial effect on the incidence of mastitis.

TABLE V.—Incidence of Mastitis in Patients who Reached the "Satisfactory" Standard—i.e., Response Before the Third Day

	Group A	Group B	Total
1st and 2nd day responses ..	55	10	62
Suppurative mastitis ..	17	—	17 (27.4%)
Non-suppurative mastitis ..	2	1	3 (4.8%)
No mastitis ..	33	9	42 (67.7%)
Untraced ..	3	—	3
Total ..	55	10	65

Discussion

The results of this experiment would seem to indicate that a deficiency of vitamin C is not the factor concerned in the high incidence of mastitis. There is no doubt that a large number of the women, when admitted to the maternity hospital, showed marked deficiency as judged by saturation tests, but they did not seem to be more susceptible to mammary infections than did those showing no deficiency. It was possible to carry out the saturation tests only when the women were in hospital, and these tests were done with two objects in view; (1) to assess the degree of saturation of the women receiving supplements, in order to get some indication of the regularity of ingestion of the tablets; (2) to estimate the degree of saturation of the ordinary hospital patient who had had no vitamin C apart from what was obtained in the ordinary diet.

It is not claimed that one testing during the period under consideration would necessarily indicate the general saturation level for the whole period; the state of vitamin C nutrition may

have varied—some women improving their intake after delivery, and vice versa. However, as the normal individual tends to be conservative in the matter of diet the tests probably gave a broad indication of the vitamin C nutrition of the subjects concerned. Apart from the difficulty of carrying out tests except under hospital conditions, it must be remembered that the controls—Group B, who had no supplements—entered the lactation period in a "saturated" condition as a result of the test-dosing. This was unavoidable, and it was assumed that these women would not materially change their dietary habits and would therefore soon revert to their former state of low vitamin C intake. In this connexion it is of interest to note that, in an endeavour to find out the duration of the state of "saturation" following test-dosing, some members of the hospital staff were tested at intervals of three weeks. Although, when considered over a long period, some seasonal fluctuation was noted, there was very little change from one testing to the next—e.g., a subject who in mid-December required three days of test-dosing to attain saturation required as many days in the beginning of January. It was thus reasonable to assume that the effects of saturation from test-dosing did not last longer than about three weeks.

If, however, it had been feasible to make more frequent tests and these had been carried out, the use of Group B as controls would then have given rise to a possible fallacy, in so far as these subjects could no longer have been considered to be having no supplements. As it was, the effect on the results of the experiment of Group B women entering the lactation period in a state of saturation is not likely to have been important, since the state of "saturation" following test-dosing does not appear to be of long duration. Moreover, the experiment was designed as an attempt to assess the possible part played by the administration of supplements of vitamin C, and Group A women, who were also saturated on the completion of the test-dosing, continued to have extra vitamin C in the form of a daily tablet containing 50 mg. of ascorbic acid. It can be argued that if vitamin C does increase resistance to infection, only in the event of the women in both groups already having such optimum intakes that they were in a state of full "saturation" and their vitamin C nutrition incapable of further improvement could additional supplies have been expected to yield no result. As test-dosing showed that this was far from being the case, Group A subjects, who had additional supplies of ascorbic acid, should have shown a lessened incidence of infection, which was not borne out by the results of the experiment.

It is apparent from the results of the saturation tests that many members of both groups, consisting of pregnant women of the type normally delivered in a maternity hospital, were deficient in vitamin C, and many, if testing had not been carried out, would normally have entered the lactational period with a gross deficiency. Judging by the average amount of ascorbic acid required to attain saturation, women in Group B showed a high degree of deficiency, while those in Group A, even with a daily supplement of 50 mg. of ascorbic acid, barely attained the level reached by institutional staffs on an ordinary diet without any supplement. Two factors may have been responsible for this deficiency: (1) greater utilization of the vitamin, due to the increased demands of pregnancy; (2) low intakes due to poverty, ignorance, or faulty methods of preparation and cooking of food.

Although many members of Group A showed some deficiency, it would be unfair to suggest that the daily supplement of ascorbic acid was necessarily insufficient, as there was no guarantee that all the women in this group actually ingested the supplements, and any defaulters in this respect would raise the apparent level of deficiency for the whole group.

In considering these negative results it must be remembered that the experiment was concerned with the intake of a single food factor. Proof of a possible relationship between mastitis and deficiency of any single dietary constituent may be difficult to obtain on account of the almost inevitable coexistence of other deficiencies. If a deficiency of one factor is present the simultaneous existence of other deficiencies is extremely likely. To increase the supply of one deficient factor without at the same time increasing the supply of the others may not be sufficient to affect the health and resistance of the subjects concerned. It is also to be remembered that the supplement

used was a synthetic product, and the maximum benefit of the protective substances may be obtained only when given in the form of natural foods.

Summary

When comparing two groups of women for a period extending from the later months of pregnancy until six months after delivery, one of the groups being supplied with supplements of ascorbic acid, it was found that there was a deficiency of vitamin C, as measured by saturation tests, in both groups; the deficiency was naturally greater in the group without the supplements.

The incidence of suppurative mastitis in both groups taken as a whole was very high. When approximately 250 expectant mothers were followed up over a period of six months following delivery it was found that 25% suffered from breast abscess. It is to be noted that, with few exceptions, these patients were delivered in hospital.

There was no reduction in the incidence of mastitis in the group receiving the supplements or in those who gave evidence of a satisfactory intake as judged by saturation tests.

REFERENCES

- Abbasy, M. A., Harris, L. J., Ray, S. N., and Marrack, J. R. (1935). *Lancet*, **2**, 1399.
 Craig, A. A., Lewis, F. J. W., and Woodman, D. (1944). *British Medical Journal*, **1**, 455.
 Department of Health for Scotland (1943). *Infant Mortality in Scotland*, Scientific Advisory Committee Report, H.M.S.O., London and Edinburgh.
 Francis, G. E. C., and Wormald, A. (1942). *Lancet*, **1**, 647.
 Fulton, A. A. (1945). *British Medical Journal*, **1**, 693.
 Harris, L. J. (1937). *Ibid.*, **1**, 774.
 — (1940a). *Lancet*, **2**, 259.
 — (1940b). *Ibid.*, **2**, 539.
 — (1942). *Ibid.*, **1**, 642, 644.
 — (1943). *Ibid.*, **1**, 515.
 — and Abbasy, M. A. (1937). *Ibid.*, **2**, 1429.
 — Yudkin, J., and Kelly, S. (1936). *Ibid.*, **1**, 1488.
 — and Olliver, M. (1943). *Ibid.*, **1**, 454.
 — and Ray, S. N. (1935). *Ibid.*, **1**, 71.
 McNee, G. Z. L., and Reid, J. (1942). *Ibid.*, **2**, 538.
 Moon, A. A., and Gilbert, B. (1935). *J. Obstet. Gynaec. Brit. Emp.*, **42**, 268.
 Neuwiler, W. (1935). *Klin. Wschr.*, **14**, 1793.
 Nixon, W. C. W. (1941). *British Medical Journal*, **2**, 703.
 Orr, J. B. (1936). *Food, Health and Income*, London.
 Portnoy, B., and Wilkins, J. F. (1938). *British Medical Journal*, **2**, 554.
 Prunty, F. T. G., and Vass, C. C. N. (1944). *Lancet*, **1**, 180.
 Smith, S. L. (1939). *Food and Life—Yearbook of Agriculture*, 235. United States Government Printing Office, Washington.
 Technical Commission on Nutrition (1938). *Bull. Hlth. Org. L. o. N.*, **7**, 461.
 Widdowson, E. M., and Alington, B. K. (1941). *Lancet*, **2**, 361.
 — and McCance, R. A. (1942). *Ibid.*, **2**, 689.
 Youmans, J. B. (1941). *Nutritional Deficiencies*, J. B. Lippincott Co., London and Philadelphia.

ACUTE IDIOPATHIC PORPHYRIA

BY

G. DISCOMBE, M.B., B.S., B.Sc.

Senior Demonstrator of Pathology

AND

JOHN L. D'SILVA, D.Sc., M.R.C.P.

Lately Chief Assistant

St. Bartholomew's Hospital

In the porphyrias uroporphyrins are excreted in the urine, in contrast with coproporphyrinuria, which occurs in many diseases. The porphyrias have been reviewed in detail by Mason *et al.* (1933), Waldenström (1937), Chandler *et al.* (1939), Roger *et al.* (1939), Dobriner and Rhoads (1940), and Nesbitt (1944).

Waldenström (1937) has described five types of acute porphyria: (a) Latent porphyria—an asymptomatic form in which uroporphyrin is excreted continuously or intermittently, for years, in the urine. (b) Abdominal type—onset sudden, with colic, vomiting, obstinate constipation, and tympanites. (c) Nervous type—onset sudden, with pareses of the extremities, convulsions, confusional states, coma, or bulbar paralysis. (d) "Classic type"—both abdominal and nervous symptoms. (e) Comatose type—abdominal colic, rapid onset of coma, and death.

In all types tachycardia and hypertension occur during the acute attack. In nearly all cases the urine contains uroporphyrin-III, with but traces of the series-I isomer; occasionally the latter predominates (Turner, 1938; Golden, 1943). Sometimes acute porphyria occurs during the prolonged administration of trional, tetralon, or sulphonal, and these have been regarded as "toxic" porphyrias (Mason *et al.*, 1933; Nesbitt and Watkins, 1942), but Waldenström considers that the division between toxic and idiopathic groups is very questionable.

The diagnosis is established by the demonstration of porphyrin in the urine. In some cases the freshly voided urine is normal in colour and contains porphobilinogen (Vahlquist, 1939), which changes to porphyrin on standing or heating. The presence of porphobilinogen is recognized by Watson and Schwartz's test (1941). The reddish-brown colour of the urine which contains porphyrin is due to the formation of coloured substances such as urofuscin, uroscopine, and indirubin from colourless chromogens (cf. Turner, 1938; Watson, 1939).

The porphyrin may be present as a zinc complex (Turner, 1938; Chandler *et al.*, 1939; Nesbitt and Watkins, 1942) which has absorption bands indistinguishable in position, if a hand spectroscopy only is used, from those of oxyhaemoglobin—differing, however, in that the β band is more intense than the α band, whereas in oxyhaemoglobin the reverse is true.

A few cases have been reported (Backer-Gröndahl, 1935; Schie, 1939) in which the symptoms closely resembled those of the "abdominal type" of the disease but the dark-coloured urine contained no detectable traces of porphyrin. Waldenström (1939) described a similar case, but during a remission of the disease the patient excreted pathological amounts of porphyrin.

The case described below is put on record for three reasons: (a) it was under continuous observation for two years, during which four exacerbations of the disease occurred; (b) in one of these the porphyrin output was followed quantitatively from day to day; and (c) in several relapses the clinical picture varied.

Case History

The patient, a sheet-metal worker aged 19, was admitted to St. Bart's for investigation on April 4, 1942. He was quite well until June, 1941, when he had a fairly sudden onset of dull aching pain round and below the umbilicus. This soon became colicky in nature and severe, but the pain did not radiate. Attacks lasted for two or three hours and had no relation to food or exercise. Nausea, vomiting, and obstinate constipation were prominent features of the illness. Some two weeks later he had five "fits" in one morning and was admitted to a hospital with a *Bact. coli* infection of the urine and a B.P. 150/110 but no other abnormal signs. The urine was free from lead, and the C.S.F. and a radiograph after a barium enema were normal. The urinary infection cleared satisfactorily, the B.P. returned to normal, and he was discharged in three weeks.

Second Attack.—In Jan., 1942, he was readmitted to hospital with symptoms similar to those in the first attack. The urine was reported as normal, but the colicky pains persisted for a few days. B.P. 160/100. He was discharged within three weeks.

Third Attack.—On March 28, 1942, he complained of aching in the limbs followed by colicky abdominal pain, vomiting on one occasion, and obstinate constipation. On April 4 he was admitted to this hospital for investigation, when, in addition to the above history, he reported having lost a considerable amount of weight during the past six months. On examination the fauces were injected, there was slight tachycardia (pulse 90), and the B.P. was 160/120. There was slight tenderness in the abdomen below the umbilicus and the liver margin was just palpable. The urine was dark brown in colour, of sp. gr. 1032, without protein, ketone bodies, or blood. It reduced Benedict's reagent, and Schlesinger's test for urobilinogen was strongly positive. Examination of the fresh urine with the direct-vision spectroscopy showed no abnormal absorption bands, but on standing for some hours it darkened and the two-banded spectrum of porphyrin became visible in a half-inch layer. An amyl alcoholic extract of the fresh or of the stale urine showed the α , β , and γ bands of alkaline porphyrin, converted by hydrochloric acid to acid porphyrin. Toluene-soluble red pigments were also present.

Investigations.—Hb, 118%; R.B.C., 5.5 millions; W.B.C., 6,800, with a normal differential count. There was no punctate basophilic. E.S.R., 1 mm. in the first hour. Serum bilirubin was normal. Blood W.R. and Kahn tests were negative. The reducing substance in the urine was proved to be glucose. A glucose-tolerance curve gave the following result (blood sugar in mg. per 100 c.cm. followed by time in hours, in parenthesis, after the test dose of glucose): 88 (0); 148 (1/2); 140 (1); 125 (1½); 115 (2). Urine specimens before the test dose and two hours afterwards contained sugar, which indicates a low renal threshold for glucose. Radiographs of the lungs, abdomen, and kidneys were normal. The skull was normal except for a small pituitary fossa (transverse, 8 mm.; vertical, 3 mm.; and interclinoid measurement, 1 mm.).

Course of the Illness

Nausea and constipation were marked, but abdominal pain did not amount to more than discomfort below the umbilicus. The