times after 78 subcultures. On subcultivation in plain broth "fastness" was maintained for two days; it then began to fall until after 76 subcultures the original "bacteriostatic" dose of 0.08 unit of penicillin per c.cm. was again effective. There is, however, considerable doubt whether 0.08 unit per c.cm. was the true "bacteriostatic" dose for the original culture, since smaller effective doses were observed on a number of occasions at the beginning of the experiment. A direct comparison of the original culture, which had been kept in a dry state, with the 82nd subculture of the resistant strain in plain broth gave the following results:—"Bacteriostatic" dose: original culture 0.02 unit per c.cm.; 82nd subculture, 0.08 unit. Nevertheless the general appearance of Fig. 1 suggests that "fastness" of haemolytic streptococci to penicillin is a temporary acquired character which is gradually lost on subcultivation in broth.

Pneumococci

Rake *et al.* (1944) subcultured *Pneumococcus* Type III in penicillin broth 55 times, thereby increasing the resistance of the strain to penicillin 30 times. After 32 subcultures of their resistant strain in ordinary blood broth the "bacteriostatic" dose of penicillin remained unchanged.

Fig. 2 shows the results of subculturing *Pneumococcus* Type III first in penicillin broth and then in plain broth. Resistance to penicillin was increased 27 times after 46 subcultures, and this level was maintained after 114 subcultures in penicillin broth. "Fastness" to penicillin was maintained after 82 subcultures in plain broth. This confirms the observation of Rake *et al.* (1944) that acquired "fastness" of pneumococci to penicillin is a permanent character.

Discussion

Although the number of strains used in these experiments is too small to draw any general conclusions there are certain facts which seem to be established. The first of these is that staphylococci become penicillin-"fast" much more easily than haemolytic streptococci or pneumococci on subcultivation in penicillin broth. The accompanying Table, which is compiled

Table showing the Number of Subcultures in Penicillin Broth required to produce Penicillin-"fast" Cultures, and the Degree of "Fastness" obtained with Various Organisms

Authors	Organism	Strain	Number of Subcultures in Penicillin Broth	Increase of Resistance to Penicillin
Rake <i>et al.</i>	Staphylococcus	"Smith"	59	6,000 times
" "	"	C 1	28	1,000 "
" "	"	C 2	28	4,000 "
Todd <i>et al.</i>	"	Oxford H	72	5,000 "
" "	"	A	77	5,000 "
Rake <i>et al.</i>	Pneumococcus	Type I	54	6 times
" "	"	" II	62	6 "
" "	"	" III	55	32 "
Todd <i>et al.</i>	"	" III	114	27 "
Rake <i>et al.</i>	Str. pyogenes	C 203	58	27 times
Todd <i>et al.</i>	"	C 203	78	4 "

from the figures of Rake *et al.* (1944) and of Todd, Turner, and Drew, shows a 4,000-fold average increase of resistance to penicillin for five strains of staphylococci, compared with an 18-fold average increase for four strains of pneumococci and a 16-fold average increase for a single strain of *Str. pyogenes*,

although the average number of subcultures in penicillin was greater for the pneumococci and streptococci than for the staphylococci.

Secondly, the observation of Rake *et al.* (1944) that penicillin-"fast" pneumococci maintain their "fastness" on subcultivation in ordinary laboratory media, without penicillin, is confirmed. On the other hand a single strain of *Str. pyogenes* lost its "fastness" on subcultivation in plain broth, behaving in this respect more like staphylococcus than pneumococcus.

Summary

Staphylococci are easily made penicillin-"fast" by subcultivation in penicillin broth; they lose their acquired "fastness" when subcultured in plain broth.

Haemolytic streptococci, so far as can be judged from the behaviour of a single strain, are more difficult to make penicillin-"fast" than staphylococci; they also appear to lose "fastness" when subcultured in plain broth in a similar way to staphylococci.

Pneumococci also acquire "fastness" much less readily than staphylococci; they maintain acquired "fastness" when subcultured in ordinary laboratory media.

REFERENCES

- Rake, G., McKee, C. M., Hamre, D. M., and Houck, C. L. (1944). *J. Immunol.*, **48**, 271.
 Todd, E. W., Turner, G. S., and Drew, L. G. W. (1945). *British Medical Journal*, **1**, 111.

Medical Memoranda

Hypersensitivity to Quinine and Mepacrine

In view of the great importance and widespread use of chemotherapy in the suppression of malaria at the present time, the following rarity is of topical interest.

CASE REPORT

The patient was an air-crew officer, aged 25, who took one tablet (0.1 g.) of mepacrine hydrochloride at 1 p.m. on Nov. 19, 1944. Two hours later he began to notice a certain itchiness and painfulness of the skin of his hands and feet, and shortly afterwards these and the scrotum, penis, and nipples began to swell. At the same time the skin of his whole body became red and his throat felt as if it was swollen. About one hour after the onset of his symptoms he began to feel hot; he had a slight frontal headache and marked anorexia and nausea, but no actual vomiting. There were no other symptoms. This was the first mepacrine tablet he had ever taken.

On examination his temperature was 104° F. and pulse 96. The skin of the whole body showed a marked generalized erythema, with no blanching on stroking it. The lips were slightly cyanosed. Some conjunctival injection was present. There was considerable oedema of hands, feet, scrotum, penis, nipples, and eyelids, with slight swelling of the fauces. Otherwise nothing abnormal was found.

Treatment with adrenaline, ephedrine, and magnesium sulphate was given, but without any effect. His temperature, however, settled in 18 hours, and all the oedema and erythema had disappeared in 48 hours. The next week the skin of his body generally, and of his scrotum in particular, peeled, and during this time it remained itchy.

His previous history showed that about three weeks before this reaction he had taken one quinine hydrochloride tablet (6 gr.), and this had produced very similar effects to the reaction described above, with the following two differences: (i) the onset of the quinine reaction occurred four hours after he had taken the tablet; (ii) he had had no rise in temperature after taking the quinine.

When the erythema and oedema of the mepacrine reaction had disappeared I scratch-tested him with both quinine and mepacrine

suspensions in normal saline, using a normal saline control; marked localized urticarial reactions occurred with both the quinine and the mepacrine, while the control showed no reaction. As a result of these tests he was considered unfit for service in malarious areas.

Such a condition must be very rare indeed, and I can find no reference to this double hypersensitivity in any textbook available to me at present. In fact, in *Memorandum on Medical Diseases in Tropical and Subtropical Areas* there occurs this statement: "Those with an idiosyncrasy to quinine can take mepacrine with impunity. Mepacrine appears to provoke no similar condition."

JOHN ORR, M.B., Ch.B.,
Fl. Lieut., R.A.F.V.R.

A Source of Error in the Determination of Cell Counts

The flatness of the cover-slips used for counting-chambers is usually taken for granted. It is perhaps not as widely known as it might be that slips which are not flat are occasionally met with. As all such slips are ground flat, one can only assume that they warp after manufacture as the result of internal stresses. Inaccurate cell counts result.

A degree of deformity too small to be perceptible by inspecting the cover-slip edge-on may cause gross inaccuracy. Thus a slip which is only 10μ "off the flat" at the highest point of the convexity will cause an inaccuracy of 10% if the correct depth of the counting chamber is 0.1 mm. A curved cover-slip, used sometimes with the convexity upwards, sometimes downwards, will cause puzzling fluctuations in the cell count of the same patient.

Once a cover-slip has been found to be defective it may be thought advisable to determine the degree of inaccuracy with a view to adjusting counts that have previously been made with it. This is possible only when one can be sure that the curve has been upwards or downwards in all estimations, when the amount by which the curve is off the flat, expressed as a percentage of the correct chamber depth, can be deducted from

Distortion of the image of a spherical bulb caused by a deformity of 10μ off the flat.

or added to the counts, as the case may be. If, as has probably been the case, the convexity was sometimes upwards and sometimes downwards, measurement of the degree of deformity will at least show whether it is so great as to place previous counts beyond the permissible limits of inaccuracy.

RECOGNITION OF DEFORMITY

Any such deformity may be easily and quickly perceived and measured by the following method:

1. Hold a corner of the cover-slip lightly between the tips of thumb and forefinger.
2. Stand with the back towards a lighted lamp, and about 1 metre from it.
3. Hold the slip at arm's length, and at such an angle that the reflection of the light bulb can be seen in it. If the slip is deformed the image will be distorted. Such deformation is usually cylindrical, resulting in an image which is elongated or compressed in one axis so that a circular light-bulb appears oval.
4. Hold the slip with the concavity facing the eye as shown by an elongation of the long axis of the oval reflection.
5. Note by what proportion approximately the long diameter exceeds the short. Then, assuming that the distance between eye and slip is 30 in., it may be shown, by reasoning beyond the scope of this paper, that if the excess is 28% the slip is 10μ off the flat (see Fig. 2), if $12\frac{1}{2}\%$ it is 5μ , and if 6% it is $2\frac{1}{2}\mu$. In practice an approximate estimate of this excess is sufficient.

The whole process is a matter of a few moments, and large numbers of slips can be examined in a short time.

The presence and extent of such deformity can, of course, be determined with the aid of a microscope with a calibrated fine adjustment. A few small drops of a solution of a coloured substance—e.g., ink—are allowed to dry on the surface of the slip, which is then placed, coloured surface downwards, in position on the counting-chamber. The dried ink is first focused, then the ruled squares, the fine adjustment being read on each occasion and the true depth of the chamber noted. The correct depth subtracted from this will give the figure required.

The first method described has the advantage of simplicity and speed, and does not entail the use of a microscope with a calibrated fine adjustment.

St. Giles' Hospital (L.C.C.), BERNARD FREEDMAN, M.B., M.R.C.P.
Camberwell.

Salivary-gland Tumour of the Upper Lip

The following record of a case of some rarity may be found of interest.

CASE REPORT

A girl aged 18 came under my care in the Royal Infirmary, Edinburgh, in March, 1940, on account of a painless swelling of the upper lip which had been present for many years but had become larger and rather disfiguring recently. It was centred strictly in the midline and protruded on the buccal surface, covered by normal mucous membrane. Using a local analgesic, a transverse incision was made through the mucous membrane, and a solid round tumour, one inch in diameter, was enucleated very easily and completely. The wound was sutured and healed without difficulty, a perfect cosmetic result being obtained.

Histological examination demonstrated the characteristic appearance of a mixed tumour of the parotid (see Fig.). There were hyperchromatic epithelial cells, arranged in the form of alveoli and in solid masses, and a stroma showing the appearance of the so-called cartilage. There was no evidence of malignancy.

COMMENT

The mixed tumour of the salivary glands occurs most frequently in the parotid and less so in the submandibular and sublingual glands, but it also arises in uncommon sites such as palate, tongue, pharynx, larynx, neck, face, orbit, scalp, auditory canal, lacrimal gland, and lips. Sir James Paget in 1851, according to Cade (1937), was the first to describe a

A photomicrograph of the specimen.

parotid tumour in the lip, and of the cases noted since then in the upper and lower lips four were recorded by Wood (1904), one by Schreiner and Mattick (1929), one by Patey (1930), two by Ahlbom (quoted by Cade), and four by Robbins (1943). An aberration in the development of the salivary-gland tissue may possibly account for some of the unusual positions. Ahlbom (1935), however, believed that the tumours in all sites should be classified as mucous-gland and salivary-gland tumours, in which case a tumour of the lip can be explained as arising from a mucous gland and not from an aberrant portion of parotid tissue. In his opinion, also, such tumours are quite benign at the age of the patient here reported.

I wish to thank Dr. R. F. Ogilvie for the histological examination and Mr. D. Smith for the photomicrograph. I am indebted to Mr. John Harkness, who saw the patient with me and actually suggested the correct diagnosis.

Edinburgh.

J. F. CURR, F.R.C.S.Ed.

REFERENCES

- Ahlbom, H. (1935). Quoted in *British Medical Journal*, **2**, 1213.
Cade, S. (1937). In *Post-Graduate Surgery* (ed. R. Maingot), **3**, 5014, Medical Publications, Ltd., London.
Patey, D. H. (1930). *Brit. J. Surg.*, **18**, 241.
Robbins, G. F. (1943). *Surgery*, **14**, 924.
Schreiner, B. F., and Mattick, W. L. (1929). *Amer. J. Roentgen.*, **21**, 541.
Wood, F. C. (1904). *Ann. Surg.*, **39**, 57.

The *Annals of Science*, the quarterly review of the history of science since the Renaissance, which has been in abeyance since December, 1942, has resumed publication with the issue of July 15.