CuSO₄ Method of Phillips et al. (1943).—As an indication of the ease and speed of this method it may be remarked that once the saturated solution of copper sulphate had been prepared the whole procedure of making the separate dilutions and finding the specific gravities of the plasmas took only four hours.

Results

The values for the total plasma nitrogen of the 45 boys are given here in the form of a distribution diagram.

The distribution of total plasma nitrogen in a group of 45 boys.

Taking these figures and multiplying by the factor 6.25 (on the assumption that the protein contained 16% N), the values obtained by the micro-Kjeldahl method are shown in contrast with those obtained by the copper sulphate method (Table I).

TABLE I.—Total Plasma N, Protein (N×6.25) and Protein as estimated by the CuSO₄ Method in the Plasma of 45 Boys aged 14 to 15. N.P.N. in 11 Samples of Plasma is also given: Values as g. per 100 ml.

	Total Plasma N	Protein (N × 6.25)	Protein by CuSO ₄ Method
Range ..	0.95–1.26	5.80–7.80	5.50–8.25
Mean ..	1.078 ± 0.14	6.65 ± 0.42	6.97 ± 0.73

N.P.N. (11 samples): Range 0.022–0.033. Mean 0.025 ± 0.0048.

The discrepancy between the values given for N content and those given for protein (i.e., after multiplying N by 6.25) is due to the fact that the latter were obtained from N values taken to one significant place of decimals only. An error of not more than ± 2% may be assumed for the N values. We now regard these values as the more accurate expression of the protein content.

Table II shows a comparison of the results in respect of 12 specimens of plasma in which the proteins were determined

TABLE II.—Plasma Proteins (g./100 ml.) by Different Methods

Micro-Kjeldahl (Total N × 5.25)	Gravimetric	CuSO ₄ Method
7.70	6.30	6.85
7.20	6.23	6.85
6.45	7.05	6.15
6.55	6.84	7.20
6.70	7.46	8.20
6.15	6.19	7.20
6.25	6.13	6.50
6.30	6.56	7.20
6.55	6.00	7.10
7.30	6.75	7.35
7.50	7.39	7.15
7.10	7.24	7.00
Mean 6.81 ± 0.53	6.68 ± 0.48	7.06 ± 0.51

by three different methods. It will be seen that collectively the agreement is fairly good, but individually the agreement is far from perfect.

In the Tables ± is the standard deviation calculated by the method of Bradford Hill (1942).

Discussion

One of us has determined the nitrogen content of the acetone-treated heat-coagulated plasma proteins and finds it to be 15.2%. This value gives a protein factor of approximately 6.6 (Cook, 1945). The nitrogen distribution of plasma is so complex that a conversion of any part of it into protein by multiplying by a simple factor is not justifiable. For example, the view that only proteins are precipitated by acid tungstate can easily be shown to

be erroneous, because in this precipitate are brought down the N-containing phosphatides. Another important factor is that the N content of globulin and albumin may well be different. Such a state of affairs may explain the individual differences seen in Table II. The whole subject merits reinvestigation. If methods dependent on N determination are used it would be advisable to express results as g. protein N or as g. N precipitated by some particular precipitant—say acid tungstate or trichloroacetic acid. For all practical purposes, however, multiplying total N by 6.25 produces a reasonably accurate result and one that is now generally accepted as giving an exact value, even if of the pseudo variety, to the proteins of plasma. It should be stressed, however, that the final court of appeal ought always to be the gravimetric method. Probably on account of its laborious nature this will never be popular.

These experiments also indicate that, for a quick and reasonably accurate method of analysis, determination of the plasma proteins by the specific gravity method is of great value.

Summary

The plasma proteins of 45 boys aged 14 to 15 have been estimated by (a) determination of nitrogen by the micro-Kjeldahl method, and (b) the CuSO₄ specific gravity method. The values obtained are shown in Table I.

In 12 cases the results from the above two methods were compared with those of the gravimetric method. Collectively, agreement between the three methods is good, but there are considerable individual differences.

The gravimetric method is the most accurate. It is considered that if a method dependent on N determination is used the result should be stated as g. N per 100 ml.

As a rapid approximate determination the CuSO₄ method is justified.

We are grateful to Dr. W. L. Burgess, C.B.E., and Prof. R. C. Garry for their interest in this work, and to Mr. H. Macdonald for able technical assistance.

REFERENCES

- Chibnall, A. C., Rees, M. W., and Williams, E. F. (1943). *Biochem. J.*, **37**, 354.
 Cook, R. P. (1945). *Proc. biochem. Soc.* In press.
 —, Davidson, W. A., Keay, D. M., and McIntosh, D. G. (1944). *British Medical Journal*, **2**, 443.
 Guillaumin, C. O., Wahl, R., and Laurencin, M. L. (1929). *Bull. Soc. Chim. biol.*, **11**, 387.
 Harrison, G. A. (1937). *Chemical Methods in Clinical Medicine*, 2nd ed., Churchill London.
 Hill, A. Bradford (1942). *Principles of Medical Statistics*, Lancet Ltd., London.
 Peters, J. P., and van Slyke, D. D. (1932). *Quantitative Clinical Chemistry Methods*, Baillière, Tindall and Cox, London.
 Phillips, R. A., van Slyke, D. D., Dole, V. P., Emerson, K., Hamilton, P. B., and Archibald, R. M. (1944). *Copper Sulphate Method for Measuring Specific Gravities of Whole Blood and Plasma*, John Macy, jun., Foundation New York.

EARLY TREATMENT OF BELL'S Palsy

BY

H. P. PICKERILL, C.B.E., M.S., F.R.A.C.S.

AND

C. M. PICKERILL, M.B., Ch.B.

(From the Plastic Surgery Department, Wellington Hospital, New Zealand)

It is said that 75% of cases of Bell's palsy recover spontaneously, and therein we think lies the cause of lifelong disfigurement for many people. It seems to be always hoped, sometimes presumed, that the particular case in hand is or will be one of the 75, in spite of the fact that there is often no means of knowing if it will turn out to be so. We should regard Bell's palsy rather from the point of view of those who do not recover spontaneously, and say that in 25% of cases the present usual line of treatment results in complete failure and the patient is permanently disfigured; this is a more arresting point of view, and is likely to result in a considerable diminution of such failures. In every fresh case of Bell's palsy we suggest that it should be assumed not that it will probably get well if little or nothing is done, but that it is possibly one of the 25 which will not recover by itself, and measures should be taken accordingly. That the treatment nowadays of any surgical condition should have so high a failure rate as 25% should of itself call for attention and inquiry. In every case of Bell's palsy the only safe procedure is to presume that it will not recover unless something more than an attitude of masterly expectancy is

adopted. We are concerned primarily with cases due to exposure to cold and not with cases due to disease or trauma of the temporal bone or to central causes, though the treatment to be advocated will be beneficial and helpful in all, even when from the start no recovery can be hoped for in such cases as severe intentional or accidental trauma. This paper is not intended to be a dissertation on "facial paralysis," but simply a contribution to assist the general practitioner, for we are convinced that the possibility of reducing the present 25% failures rests in his hands.

Probably the chief factor in non-recovery of what should be the temporarily paralysed muscles is their immediate and continued overstretching by the non-paralysed muscles of the opposite side of the face. A common method of attempting to overcome this is by the use of a hook in the corner of the mouth attached by a loop to the ear; but, as will be shown, this is often worse than useless. A more simple and effective method of restraining the overaction of the muscles of the sound side is by means of strapping. It is applied thus.

The Strapping Method

The lower half of the paralysed side is first smeared thinly with vaseline or oil. Two strips of half- or three-quarter-inch strapping are then applied to the clean sound side on a level with the upper and lower lips, starting about three-quarters of

at the end of a week or ten days. The latter method is probably the most effective that can be employed to aid recovery, but it must be done while the normal side is out of action. That is to say, conditions must be such that the only thing the paralysed orbicularis oris or zygomatic muscles have to do is to respond, even if ever so feebly, to the will to contract; they must not be concerned also with opposing the pull of the muscles of the opposite side. The probable reason why massage and exercises are so often ineffective is that in the early stage of recovery the muscles are overpowered with work by having two jobs to do at once, and, again becoming overstretched, anything gained is soon lost. The paralysed orbicularis oculi muscle should be supported by a sling composed of two quarter-inch strips of strapping running downwards from each end of the eyebrow and meeting below the centre of the lower lid, as shown in Fig. 1. This serves to keep the lid in apposition to the eye, to prevent epiphora, and to counteract the weight of the paralysed cheek and so prevent the orbicular muscle from losing tone. Fortunately this muscle is not concerned with opposing its fellow on the opposite side, and so this method of strapping is nearly always successful in its object; it may be said that if it is applied early enough, and persisted in, there will certainly be no ectropion to follow in simple Bell's palsy.

Electrical Measures.—During the third week, if recovery is not obvious, the muscles may with advantage be stimulated with

FIG. 1.—Showing method of applying strapping. S=strapping stuck to skin on sound side. N.S.=strapping not stuck to skin, on paralysed side. Also showing sling for paralysed orbicularis oculi muscle to prevent ectropion.

FIG. 2.—The strapping stuck to the skin on the sound side; it should not be carried back further than shown.

an inch in front of the ear (Figs. 1 and 2). The sound side is pushed across to the paralysed side, and while in this position the other ends of the strapping are firmly stuck down to the mastoid process on the paralysed side (Fig. 3). Good and very adhesive rigid strapping must be used—not elastoplast. The strapping, once on, must be left indefinitely on the sound side. If it is removed the skin becomes sore and there is difficulty in reapplying it firmly; moreover, the patient is discouraged from persisting with the treatment. When it gets soiled another piece may be placed on top of it, and only in some real necessity should it be removed, gently, by the surgeon or nurse, with the aid of a solvent, and reapplied immediately. Non-adhesion to the paralysed side is necessary in order to avoid the possibility of a pull on the paralysed muscles and so that gentle massage may be given; also so that the patient may endeavour to make voluntary movements of the affected side in front of a mirror

weak faradic currents. Minimal doses only should be used—just sufficient to cause visible contractions in the orbicularis and zygomatic muscles. The aim should be just to maintain tone and not to tire the muscle with violent contractions. A current which causes any painful sensation is much too strong, and will do harm. The current should be tried first on the operator himself and then on the sound side of the patient before being applied to the paralysed muscles. If a current which causes contraction on the patient's sound side causes no contraction on the paralysed side, no attempt should be made to produce contraction by using a stronger current. Short daily treatments with very weak current should be ordered. The strapping should be kept on and the above treatment continued indefinitely. It may be for only a week or two, but it may be for two or three months. The patient must be handled very strictly, and be impressed with the serious risk of taking the strapping off even for one occasion, no matter how seem-

ingly important it may be. One laugh without the strapping may undo weeks of treatment, and to leave it off for several days just when recovery is beginning will almost certainly result in partial failure.

The Mouth Hook.—From what has already been said it will be obvious that if the point of decussation of the orbicularis oris muscle on the paralysed side is made a fixed point, then, when the overacting sound fibres contract, the effect on the paralysed fibres will be to stretch them still more—much more than if there were no hook. If the hook is so adjusted that it actually pulls the angle of the mouth back, not only will the stretching be more severe but the fasciculi radiating from it, the depressor and levator anguli oris muscles will be stretched too. The only muscles which the hook really rests are the zygomatics, and in cases in which the hook has been used we often see that these are the only muscles which have recovered, the orbicularis being particularly flaccid. The rule should be, "Look after the sound side—leave the paralysed side alone; it is the sound side which does the damage."

Other Methods of Treatment

Nerve grafting in traumatic (mastoid) cases is obviously the method of choice and, equally obviously, should be employed

FIG. 3.—Showing method of strapping on the paralysed side. N.S.=Not stuck. S=Stuck very firmly to the mastoid process and a little way beyond the hair line.

early if it is to be successful, as also should decompression of the aqueductus fallopian if thought necessary; these operations, however, fall within the sphere of otolaryngology and will not be discussed here.

Fascia Lata and Muscle Grafts.—These are the plastic surgeon's methods of dealing with facial paralysis in cases usually referred in the later stages, when R.D. is complete. We would, however, like to advance a plea that all traumatic cases (even those few for which nerve grafting is chosen) should be referred to the plastic surgeon as early as possible after the injury. Slings of fascia lata to take the weight of the paralysed cheek, to restrain the overacting muscles of the opposite side, are ideal for the purpose; they are permanent and invisible, and should be used instead of strapping, which can be regarded as only a "tide-over" method in this class of facial paralysis, but which should nevertheless be adopted immediately after the trauma,

and replaced as soon as possible by fascia lata. For instance, if the facial nerve is implicated in a parotid tumour and the surgeon judges that it is better to sacrifice the facial nerve and save the patient's life, the strapping described above should be applied at the end of the operation, and the case referred to the plastic surgeon in three weeks' time. There is nothing to be gained by waiting, but much to lose. It is surprising how little tone the muscles seem to lose when they receive early and permanent support in this way. Here as elsewhere it is easier, and the result is infinitely better, when deformity is forestalled than when reliance is placed on correcting it after it is well established. The difference between the results of the same operation (fascia lata grafts) done three weeks after total facial paralysis and three years after, is almost incredible to those who have not had the opportunity of such comparison. There is some strange persistent urge to "wait and see" if recovery will take place or whether the paralysis will become bad enough to need operation. The answers to this are: (1) recovery will be hastened and not retarded by the fascia lata operation, which may make recovery possible in some cases where it would otherwise be impossible; (2) if nothing is done in the case of total paralysis it is *certain* to become bad enough to require operation, and the golden opportunity of obtaining the best result has been lost.

In well-established long-standing cases (which in future we think should not be permitted to exist) the choice lies between (1) fascia lata slings for the orbicular muscles, and perhaps temporal and masseter muscle slips to give some voluntary movement after re-education, and (2) anastomosis with the hypoglossal nerve. The former methods give balance to the face, with a little muscular movement of the angle of the mouth and full closure of the eye; while the latter method, when successful, provides mass movements only, and facial balance in repose is not often so good as in the sling method.

OSTEOMYELITIS OF THE SKULL DUE TO SALMONELLA TYPHI

BY

J. M. LEVER, M.B., Ch.B.

Clinical Pathologist, Atkinson Morley Emergency Hospital

AND

G. B. BARKER, M.R.C.S., L.R.C.P.

First Assistant, Neurosurgical Unit, Atkinson Morley Emergency Hospital

Cases of osteomyelitis due to the typhoid bacillus are uncommon; the following case is also of interest owing to the great length of time that elapsed between the initial infection and the occurrence of osteomyelitis, the site of infection, and mode of onset of the condition.

Case History

A man aged 40, a linesman on the L.P.T.B., was admitted to this hospital with a history of headaches for two years. These had occurred almost daily, and were of a stabbing character lasting varying times from one hour to a whole day. They did not interfere with his work; he had accepted them as part of his life and did not seek medical advice.

For the three months before admission they had been getting steadily worse; he had been losing sleep and was beginning to feel weak and exhausted. On Dec. 24, 1944, he was sitting in his kitchen when he felt a tingling sensation in his right hand. He rose from his chair and found that he had lost control of his right leg, which was useless. This attack lasted one hour and then passed off, leaving him apparently normal in every respect. He consulted his doctor and was admitted to hospital, where he had a similar attack. Investigations proved negative and he was discharged. He still lacked energy, however, and the headaches had increased in severity. Shortly after this he had a severe pain in the back of the head, agonizing in nature; he consulted his doctor, who found a soft fluctuant swelling in the occipital region, and he was again admitted to hospital.

During the next four days the swelling increased steadily in size, but the severe pain passed off, to be replaced by a dull ache. On admission to us, 10 days later, he had a soft, diffuse, smooth, very tender, hot swelling 2½ in. in diameter over his occipital region, mainly on the right side, and considerable oedema of the surround-