BRITISH MEDICAL JOURNAL

LONDON SATURDAY FEBRUARY 5 1955

MEDICAL SCIENCE AND PROBLEMS OF FLYING*

BY

Sir HAROLD E. WHITTINGHAM, K.C.B., K.B.E., LL.D., F.R.C.P., F.R.C.S.Ed.

Director of Medical Services, British Overseas Airways Corporation

In the development of flying, medical science has played a very important though naturally an unobtrusive part in accordance with the traditions of the profession. It has helped to solve problems in the selection and training of aircrew and of air traffic control personnel, the design of such flying equipment as oxygen systems, goggles and glasses to counteract glare, intercommunication sets to obtain good intelligibility, safety harness and other survival devices, including ejection seats and parachutes. Medical science has also contributed to the interior design of aircraft to promote and conserve efficiency of human performance, in particular pressurization, the lay-out of flight decks and gun-turrets, the shape and position of controls, the presentation of instruments to give the best grouping and siting of the most important of them, and to provide legible dials, together with suitable lighting to facilitate dark adaptation as well as map reading by night. In addition, it has aided the development of equipment to prevent blacking-out in aerobatics, of runway lighting for ease of landing and take-off, and of many other matters such as the design of aircraft seats to promote comfort and lessen fatigue, anti-air sickness remedies, and techniques for disinfecting and disinsecting aircraft.

It is impossible in the space at my disposal to deal with this vast subject, so I shall give a brief review of the part played by applied physiology to war in the air, mainly as it concerned the Royal Air Force. During the first world war the work of Flack,^{41 42} Head,⁵² Birley,¹³ and Bazett¹¹ gave us certain selection tests, which ensured that those who passed were alert in body and mind, had accurate vision, good muscle tone, quick reaction time, and a proper sense of balance. These tests did not predict the individual's aptitude to learn to fly or his suitability for high-altitude flying or aerobatics, but they aided in the assessment of general physical fitness to withstand the stress of flying.

Applied physiology in relation to efficiency in flying had only really begun when the first world war finished, and it remained dormant in this country until 1935. During the ensuing four years, tests were introduced to assess suitability for particular aircrew duties, the main ones, which were developed at the R.A.F. Central Medical Establishment, being for power of co-ordination,¹¹² cardiovascular stability,⁸⁸ tolerance of rarefied atmospheres,⁷⁴ night visual capacity,^{57 66} and hearing efficiency when operating in aircraft^{29 33 35 74}; this hear-

ing efficiency test was replaced later during the war by one simulating speech sounds as actually transmitted through an electrical communication system with a background noise as encountered in Service aircraft.^{36 37} The problems of glare,⁵⁴ ocular muscle balance,⁵⁵ and the design of flying goggles⁵⁶ and helmet^{30 31 32 34} were also investigated.

Early in 1939 the Secretary of State for Air, on the instigation of Air Marshal Sir Victor Richardson, then Director-General of Medical Services, formed the Flying Personnel Research Committee, under the chairmanship of Sir Edward Mellanby. As a result, research was intensified and planned to cover all current medical problems connected with Service flying. Under the aegis of this committee a certain amount of fundamental physiological research was instituted, but, subsequent to the outbreak of war eight months later, many of the researches had to be of an *ad hoc* nature. It is convenient to consider these researches in relation to the main phases of the war—namely, the phony period, the Battle of Britain, the night raids on London and the industrial cities, the retaliation by the R.A.F., the great air offensive in the West, and the air war in the Orient or the setting of the Rising Sun.

The Phony War

In the phony period from October, 1939, to Spring, 1940, the chief physiological problem which demanded attention was an adequate oxygen supply for those engaged in high-altitude reconnaissance when the main shooting was photographic. Though there was little fighting during the first eight months of the war, the Air Force was busy developing the system of photographic reconnaissance which later proved such a great factor in its success. In this work it was important to attain altitudes ever higher than the enemy, in order to escape detection or interference. This necessitated the development of an adequate oxygen system. At the outbreak of war the continuous-flow system of oxygen supply in use allowed two-thirds of the gas to go to waste, and the mask did not fit comfortably with the flying helmet, goggle, and microphone assembly. Work carried out in the laboratory by B. H. C. Matthews and staff,⁷⁵ and in the field by D. A. H. Robson (who later lost his life in an experimental flight of one of the first Boeings used in this country), led to the early development of an improved oxygen mask and an economizer, which prevented the oxygen from going to waste during expiration and was foolproof against

*A special university lecture in military studies given at the University of London on May 25, 1954.

stoppage of the flow of the gas even if the breathing of the wearer became minimal, as during shock or unconsciousness (Fig. 1). This improvement in the oxygen supply more than doubled the operational range of aircraft from an oxygen point of view. The anoxia meter devised by E. A. G. Goldie^{48 50} was of inestimable value in assessing the relative merits of different types of oxygen masks and revolutionized the study of anoxia, as the amount of oxygen in the blood could be estimated with this apparatus at any moment and for any length of time without puncture of an artery as was necessary hitherto (Fig. 2).

As decompression sickness occurred in some pilots engaged in reconnaissance at altitudes over 25,000 feet (7,600 metres), E. A. Carmichael,¹⁷ B. H. C. Matthews,⁷⁴

and I. de Burgh Daly²⁶ studied this matter and made the first appreciation of its effects. Later, the staff of the R.A.F. Physiological Laboratory at Farnborough, working in liaison with the Royal Canadian Air Force, intensified these researches and, as a result, tests were devised to select those suitable for high-altitude flying, and measures were adopted to alleviate or prevent the occurrence of decompression sickness.^{38 46 92 95 97 119} This pioneer work materially aided in Bomber Command's retaliation on the enemy, which began in 1941.

The aircraft used in these reconnaissance flights were of the single- or two-seater type, had reasonably warm cabins, and were engaged only on flights of relatively short duration, so that the effect of cold on the individual and his flying equipment was not markedly noticeable and did not call for research then.

Work done in connexion with the oxygen supply during this phony period proved of inestimable value for the Battle of Britain.

The Battle of Britain

The enemy, having failed to attain victory by force of numbers, tried to circumvent the air barrier by flying over it. The famous few had therefore to climb to equal or greater altitudes to claw the enemy out of the sky, which would not have been possible without an adequate oxygen supply.

At this time two other problems demanded immediate attention—namely, the temporary loss of vision or blacking-out which resulted from high acceleration during dog-fighting, which was then the vogue; and fatigue consequent on the large number of sorties that had to be flown without a rest.

Fortunately, in the absence of a centrifuge for the fundamental study of blacking-out, a considerable amount of research in this matter had already been carried out by W. K. Stewart and G. E. Watt^{108 109 107} at Farnborough in an aircraft specially fitted for use as a flying laboratory. In this aircraft a cine-camera was suitably mounted to take shots of pilot and passenger, so as to record their physical appearances and visual reactions under various forces of *g*, together with the readings of the various instruments indicating force of *g*, time for which applied, air speed, and altitude (Fig. 3).

In this way it was proved that abdominal belts *per se* were of no use as an anti-*g* device,^{99 102} but that resistance to blacking-out could be increased by 1½ to 2 *g* if the rudder-bar pedals of the aircraft were raised by 6 in. (15 cm.), and the pilot assumed a crouched position, so as to lessen the amount of blood that pooled in the legs and to reduce the differences in the levels between the brain and the heart (Fig. 4). This minor modification to the aircraft was readily carried out at the airfields without interfering with the operational effort—a most important consideration at a time when every available aircraft had to be in the air or at the alert. The raising of the blacking-out threshold by 1½ to 2 *g* proved adequate to prevent blacking-out with the centrifugal forces encountered in aerobatics at that period.^{99 102}

The Battle of Britain was such a critical period that little or no rest was possible for our pilots in their all-out effort. Medical aid was sought to give relief from sleeplessness and fatigue without, if possible, loss of performance.⁸ Several drugs were tried, but attention was mainly directed to the use of caffeine¹⁴ and amphetamine ("benzedrine")^{7 14 15 28 99 115 117}; the former was found to stave off fatigue for a short period without interfering with performance. The popular belief at the time was that amphetamine was the wonder

FIG. 1.—Oxygen economizer designed to prevent oxygen going to waste during expiration.

FIG. 2.—Goldie's anoxia meter. The apparatus is lightly clamped to the ear and the oxygen content of the blood is measured by means of two photoelectric cells. (Air Ministry Crown copyright.)

drug to banish fatigue and improve performance—in fact, some thought it would give them a mental acumen that they never possessed.

Professor F. C. Bartlett and his co-workers¹ showed that amphetamine, though it helped to banish sleep, did not im-

prove performance and was mainly a stimulant and acted as a whip to the flagging horse, so that when its effect passed off the need for rest was greater than ever. There was the danger that if it were taken to lessen fatigue prior to a landing its stimulative effect, something like that of alcohol, would so warp judgment that undue risk might be taken, with dire results. The stimulative effect varied according to the type of individual, and in correct dosage the dull appeared somewhat brighter, the average was perceptibly enlivened, while the above average who was normally a "live wire" became "hay-wire" with frayed nerves. Consequently, orders were issued that no drugs were to be given to aircrew, except by the unit medical officer, and then only to individuals in accordance with their known reactions, and that as a general rule no drug should be used except to ward off sleep when the danger of that was greater than any ill effects of the drug. The importance and necessity for natural sleep, even for short periods, was stressed.²⁻⁸

Night Raids by the Enemy

When, in the last quarter of 1940, the enemy changed tactics from daylight bombing to terror raids at night on London and the industrial cities, anti-measures by the Air Force called for intensive research into the problems of night vision in general, and the protection against searchlight glare, to help our pilots and air gunners to detect the enemy during interception flights.⁹⁻¹¹ Night-vision-capacity tests were devised to help in the selection of aircrew most suitable for night operations,¹²⁻¹⁴ and investigations were carried out to assess the effects of vitamins A and C and various drugs on night vision.¹⁵⁻¹⁷ At the same time, physiological and psychological aspects of night vision had to be considered in relation to aids for taking-off and landing in the dark. This necessitated improved methods of pre-adapting the eyes for night vision,¹⁸⁻²⁰ first by wearing dark smoked glasses²¹ to exclude all light, and later by the use of ruby-red glasses²² which permitted the user to read while standing by for taking-off.

Cockpit and cabin lighting were studied to ascertain the best type and intensity of illumination that could be used to read instruments and maps without materially interfering with dark adaptation.²³⁻²⁵ This led to the development of a satisfactory lighting system whereby fluorescent or red lighting²⁶⁻²⁸ could be used either together or separately, and so controlled that the maximum permissible luminosity could not be overstepped. At the same time, night-vision training schools were established to teach all members of aircrew how to maintain and improve their night vision and how to see or observe, including the value of oxygen in preserving night vision.²⁹⁻³¹ Those mainly concerned in these researches were K. J. W. Craik,³²⁻³⁴ and P. C. Livingston.³⁵⁻³⁷ They also played an important part in the development of devices to counteract searchlight glare³⁸⁻⁴⁰ and in the lay-out of airfield lighting systems to facilitate taking-off and landing in the dark.

Retaliation

Starting in 1941, retaliation bombing by the R.A.F. became ever stronger, and for this purpose the range of flight was increased and operations were carried out at higher altitudes, partly to permit surprise attacks and partly to evade anti-aircraft fire. At the same time, bomber aircraft were provided with more guns for defence, which meant more draughts through the gun apertures. Intense cold was therefore experienced by aircrews, particularly tail-gunners. It was not practicable, owing to operational needs, to install efficient fuselage heating in the aircraft then in use; the immediate necessity was to devise clothing that would keep the aircrews reasonably warm without hampering their movements.⁴¹ Hot-air suits were tried, but were not a success.⁴² However, as a result of the combined researches of physiologists and material scientists, special flying clothing, consisting of electrically heated suits, boots, and gloves, was devised to apply warmth to those parts of the body most in

FIG. 3.—Original blacking-out experiments in a flying laboratory in Battle aircraft showing cinecamera record of effects of high g on man. (A) At 8 seconds and a force of 6.2 g: beginning of blacking-out. (B) At 18 seconds and a force of 2 g: still blacked out, but consciousness returning. Dials on left side record performance. (Air Ministry Crown copyright.)

FIG. 4.—Diagrammatic representation of raising the threshold for blacking-out by adopting a crouched position with legs raised on rudder bar to lessen the pooling of blood in the dependant parts of the body.

need of it; this flying clothing combined lightness and flexibility to facilitate movement, waterproofness against rain, and permeability to air to allow for dispersal of sweat.^{69 71 72}

Another ill-effect of the intense cold encountered was the freezing-up of the oxygen-mask valve under icing conditions.⁸¹ J. C. Gilson⁴⁶ tackled this problem and produced an improved type of mask which did not freeze up at high altitudes and had the additional qualities of comfort, smallness, lightness, and perfect fit even during strenuous aerobatics. For these long bombing flights it was necessary to devise a light, portable oxygen set to enable aircrew to move about away from their stations at altitudes over 15,000 feet (4,600 metres), especially when loading guns or moving ammunition.⁸⁹ Before the introduction of this portable set, there were instances of crews collapsing from oxygen want when away from their normal oxygen points.²⁰ A further development of this set was the bale-out bottle to be used either fixed to the person or combined in the parachute pack to ensure an adequate oxygen supply for 10 minutes to enable aircrew to escape from aircraft and descend safely in emergency.^{83 93} These oxygen supply devices were introduced mainly as the result of researches by B. H. C. Matthews and his staff,^{4 76} and played an important part in maintaining the efficiency and morale of aircrew⁸ on hazardous missions.

At this time, anti-submarine activities were intensified and the medical branch was asked to advise on the best visual means of detecting submarines in various waters. This involved camouflage problems and the drawing up of a scheme for teaching aircrew how to look for submarines, and the best means of scanning.^{1 44} Coupled with this the optimum period of watch and the maximum period permissible to avoid fatigue of such a degree that further watch was useless had to be assessed.^{81 114} Another method of countering submarine and air attacks on our convoys was the use of catapult aircraft on merchantmen. These had often to operate in northern waters, and at the end of the sortie the pilots had to bale out, as they could not land their aircraft on the ships. This meant immersion, perhaps in

icy waters, for half an hour or more, before action was broken off and they could be rescued. It was necessary, therefore, to devise an immersion suit which would permit adequate perspiration during standby and still keep the wearer dry and warm enough to remain alive while immersed.¹¹⁶ E. A. Pask^{88 89} was mainly instrumental in the development of such a suit, the principle of which was subsequently adopted by the Royal Navy and the Allied Air Forces (Fig. 5).

Life-jackets were naturally provided for aircrew operating over the Atlantic to escort convoys, spot enemy submarines, and give defence against enemy aircraft, but it was necessary to ascertain whether these life-jackets would keep the nose and mouth of an unconscious individual above water in a rough sea. To test this, E. A. Pask⁷⁸ was anaesthetized on two occasions for several hours while immersed in water wearing various types of life-saving jackets in turn (Fig. 6). It was

found that not only the R.A.F. "Mae West" but also life-saving jackets of other Air Forces and Navies, though adequate in giving safe flotation to a fit person, could not be relied on for an unconscious individual. Following these researches, suitable modifications were made to the life-saving jackets to ensure adequate flotation for the unconscious.

These long, hazardous flights over the ocean and enemy territory were accompanied by considerable fatigue and an increased incidence in nervous breakdown. In connexion with the latter, C. P. Symonds and D. J. Williams⁸ made extensive surveys of the psychological disorders of aircrew, and their findings enabled authoritative advice to be given

FIG. 6.—Assessing flotation properties of life-saving jackets to keep nose and mouth of unconscious individual above water. Subject was anaesthetized through leads to nose. (Air Ministry Crown copyright.)

on operational limits to conserve man-power and maintain efficiency. Researches by physiologists and psychologists into the problem of flying fatigue pointed the way to lessen this by properly designed cockpits and cabins, including seating position, instrument lay-out and dial markings,⁹ lighting to preserve dark adaptation, the best position for all controls so as to obtain the necessary reaction with the minimum of energy, sound-proofing to minimize engine noise, and air-conditioning to maintain a reasonable temperature in all climates and at all altitudes.

This work, and subsequent improvements in the aircraft and flying equipment for maintaining the maximum efficiency of aircrews, proceeded steadily and in parallel with the increasing striking power of the R.A.F. in preparation for the great offensive in the West.

The Great Air Offensive in the West

The air offensive on enemy and enemy-occupied territory was stepped up considerably in 1943, and reached crescendo in 1944 and 1945. Larger and faster bombers were used, flying at still higher altitudes, and these offensives were interspersed with low bombing attacks by much faster-flying aircraft, such as the Mosquito, followed later by the dive-bombing attacks by the very high speed rocket-carrying Typhoons. Operations were carried out both by night and by day, and pin-point bombing became the routine, especially for the "Pathfinders." The physiologists and psychologists were consulted regarding the design of the bomb-aiming apparatus, the best colours for target-markers, and the pattern of the marking.^{6 10 80}

At this time the enemy's anti-aircraft defences were considerably strengthened, and violent evasive action was often necessary to avoid the heavy flak. The enemy's night fighters were frequently concentrated to attack our raiders, so it was essential that both pilot and gunners should have the clearest vision possible to forestall the fighters' attacks. Researches by A. Luczak,⁶⁷ of the Polish Air Force, showed that a clear-vision panel was the ideal, but, failing this, plate

FIG. 5.—Original immersion suit made of fabric which allows the passage of water vapour—i.e., perspiration—but is waterproof. It was designed to keep a man's clothing dry after ditching before boarding his dinghy or being picked up. (Air Ministry Crown copyright.)

glass set as near the vertical as practicable was to be preferred to "perspex." Goldie⁴ demonstrated how markedly visibility was reduced by scratches and dirt on windscreens, and devised an apparatus for measuring the scatter of light and visual loss which resulted from using dirty perspex and other transparent screens: this gadget did much to convince aircrews of the need to maintain clean, unscratched vision panels.

To facilitate the detection and pin-pointing of targets in the dark and through cloud or mist, and to assist in homings under adverse weather conditions, various instrument aids, visual and auditory, were introduced. Physiologists and psychologists collaborated in this work, not only in the laboratory but also in the air in actual operations. The "beam approach" was a good example, as it concerned both visual and auditory aids. The pilot relied on the beam approach device in homing at night or in bad visibility by day at a time when he was likely to be tired and particularly apt to make errors. In addition to this apparatus, the pilot had at least ten instruments on his flying panel which had to be read and their meaning fitted into a whole before he could make a safe landing under blind flying conditions. It was of supreme importance, therefore, that all instruments should present their messages as simply and clearly as possible.

The beam approach consisted of a radio beam which gave two types of clue—one auditory, the other visual. Originally, these were wrongly presented, as the auditory clue did not use the morse letters that could be picked up with least error; the visual clue, consisting of a small luminous plan of the aircraft and its relation to the beam, was confusing, as the pilot had to remember to reverse the signal for right or left turn when approaching the airfield in the opposite direction to the normal. Psychological researches by R. C. Browne¹⁴ showed that the best presentation was that which necessitated the least thought, and as a result correct clues were introduced. The application of this work to the promotion of safety in flying needs no stressing.

Stratosphere Flying

Both sides were striving for the key to victory through mastery in the air, so two projects received high priority—namely, the development of the pressurized cabin and jet-propelled aircraft.

The pressurized cabin aircraft was needed for duration flights at altitudes over 38,000 feet (11,600 metres), and certain physiological aspects of pressure differentials and air-conditioning had to be solved.¹⁷ An important point that required early clarification was the effect that sudden decompression, following holing by a missile or the blowing of a port, would have on the occupants. It was thought that such an accident would lead to so much distension of the lungs as to cause serious injury or death. R. H. Winfield,¹¹⁸ by personal experiment, demonstrated that explosive decompression at the differential pressures visualized caused no perceptible harm to a fit man provided his glottis was open.^{94 108} Experiments on animals carried out by Professor de Burgh Daly and his team at Edinburgh University gave similar results.^{26 38 39} To meet the enemy's challenge for altitude supremacy, it was necessary to develop some interim measure, and here Professor H. C. Bazett¹² and A. P. Gage,^{44 45} of U.S.A., with their pressure-breathing waistcoat and mask, respectively, saved the situation, for this equipment so raised the intrapulmonary gas pressure as to maintain sufficient arterial oxygen saturation to enable aircrews to fly

efficiently at altitudes up to 44,000 feet (13,400 metres). This was a requirement of the high-altitude photographic reconnaissance flights, whose work was of the utmost value in deciding on targets and in assessing the result of operations.

Other physiological problems of stratosphere flying concerned ventilation, oxygen needs, visibility, and methods of escape in emergency. In connexion with escape descents by parachute from the stratosphere, physiological aid was sought as to how to guard against the intense cold (-55° C. or less), lack of oxygen, and the high forces of deceleration at the moment of opening of the parachute. Suitable clothing was designed to protect against cold. The problem of oxygen lack was studied by E. A. Pask, W. K. Stewart, and S. L. Cowan,⁴⁷ who carried out a series of experiments in the laboratory to ascertain the length of time the average man could live at altitudes between 44,000 and 25,000 feet (13,400 and 7,600 metres) during a parachute descent, although unconscious. These volunteers were suspended in the air by means of the standard parachute harness and made to breathe nitrogen-air mixtures to simulate the atmospheric conditions from 44,000 feet (13,400 metres) downwards at the rate of descent of an open standard parachute. It was found that man could remain unconscious for $7\frac{1}{2}$ minutes and yet live during such a descent without additional oxygen: the bailout bottle was therefore designed to give a 10- to 12-minute supply of oxygen to allow a sufficient margin of safety to cover the oxygen used by the energy expended in getting out of the aircraft.

Experiments in U.S.A. by R. Lovelace and M. W. Boynton by personal jumps, and by G. A. Hallenbeck,⁵⁵ using large animals and dummy men fitted with accelerometers, indicated that decelerating forces up to 30 g occurred at altitudes of 35,000 to 40,000 feet (10,700 to 12,200 metres) if the parachute opened almost immediately after baling-out, and this force severely damaged the canopy and its rigging. It was therefore essential to warn all parachutists to allow the speed of fall to slow to a steady rate before pulling the ripcord, and this free-drop time had to be worked out for various altitudes. In one of these very gallant experiments Boynton lost his life. B. J. O. Winfield,¹¹³ who personally made many parachute jumps, added considerably to our knowledge of this art, and his observations helped in the design of parachutes and their attachments to the user, and in establishing the best stance to adopt for landing so as to minimize injuries.

Physiological Problems of High Speed

The advent of jet-propelled aircraft with speeds approaching that of sound soon raised other physiological problems,

FIG. 7.—Shows the first successful ejection from aircraft in flight of a dummy man complete with seat following a 40-foot-per-second explosive charge: the airspeed was 250 m.p.h. The parachute, seen in lower photos, operated automatically after ejection. (Air Ministry Crown copyright. Photograph of official test done on October 1, 1945.)

an early and very important one being emergency escape during flight. As a result of a fatality in January, 1944, when a pilot tried to escape from a jet machine which was out of control, it was evident that the force of the slipstream of high-speed aircraft was so great that the unprotected human body could not withstand the air blast for more than a fraction of a second, else limbs were fractured or evulsed against the edge of the escape hatch, the skin of the face torn, and the lungs over-expanded. Researches done on the ground with a Martin-Baker test rig, then in fast-moving aircraft, first with dummies and later with men, showed that it was practicable to eject a pilot, complete with seat and parachute, clear of an aircraft travelling up to speeds of 500 miles (800 km.) an hour (Fig. 7). For safe ejection, however, it was necessary to devise a canopy to prevent undue flexion of the head during the upward thrust, to protect the face and lungs against blast, and to keep the oxygen mask from being torn off by the force of the slipstream.⁴⁷⁻⁵³ In the final design, the pulling down of the protective canopy automatically activated an explosive charge under the pilot's seat only when the body of the canopy reached nipple level, and the resulting force of ejection ensured that the pilot's hands drew the hood tightly over his head.⁵⁴⁻¹⁰⁵

While the war in the air was proceeding apace, important events were taking place in enemy-occupied territory from resistance groups with whom we were in touch by agents dropped by parachute and at times picked up from air. Owing to the difficulty of landing aircraft for this purpose, a technique had to be developed to pick up these agents from the ground while the aircraft was in flight. The physiological and physical problem here was how to accelerate man from rest on the ground to approximately 120 m.p.h. (190 km.p.h.) in the air within half a second. W. K. Stewart and R. H. Winfield¹⁰⁶ did most of the physiological work in connexion with this problem and took part in practising and perfecting the technique (Fig. 8).

Looking ahead to the landing in Normandy and the advance to Germany, it was expected that there would be aerial dog-fights at higher speeds than hitherto, so researches were done in Britain, Australia, Canada, and the United States to develop and perfect anti-g garments that could be rapidly donned and would be comfortable to wear even in hot weather and during long sorties. The combined efforts of these researches led to the production of a very light (3 lb.—1.4 kg.) pneumatic suit or high-waisted trousers fitted with bladders, so designed as to equalize automatically the internal pressure built up in the body fluids by the accelerating forces, thus preventing the blood from pooling in the lower part of the body (Fig. 9). By wearing such a suit the threshold for blacking-out was raised by 2 g or more, and, in addition, by ensuring good circulation of blood to the head, lessened fatigue and helped to maintain quick reaction time during

long and arduous flights.²⁷⁻¹⁰⁶ Much of the credit for this work belonged to those pioneers W. R. Franks,⁴⁴ of Canada, and F. S. Cotton,¹¹⁷⁻¹¹⁸ of Australia, to S. Davidson,⁵⁷ a medical officer who was test pilot for these anti-g devices, to W. K. Stewart,¹⁰⁰⁻¹⁰¹⁻¹⁰⁴ who acted as a guinea-pig in over 200 black-outs, and to our American Allies, particularly the Mayo team—E. J. Baldes, C. F. Code, E. H. Lambert, and E. H. Wood.¹²⁰ The Allied air supremacy was so overwhelming, however, that dog-fights did not occur, so these suits were not required by the R.A.F., though they were used in large quantities by the Americans, chiefly to lessen fatigue on long flights under turbulent atmospheric conditions.

The Air War in the Orient

Following the surrender of Germany in May, 1945, researches in connexion with physiological problems peculiar to the Orient were intensified for the final showdown against the Japs. Special attention was paid to air-conditioning and cooling of aircraft, particularly when on the ground, to protect against the ill-effects of heat, and to the development of escape suits and equipment for survival in the jungle and in warm climates after baling out or crash-landing. The results of these researches were not adequately tested in the field, owing to the abrupt and dramatic end of hostilities in the East, as a result of atomic warfare. Nevertheless, much progress was made and sound foundations were laid for post-war research into problems of climatology, clothing, and human performance in temperature extremes.

FIG. 9.—Pneumatic anti-g suit. A: Shows position of abdominal, thigh, and lower leg bladders as indicated in solid black. The shaded areas are pressurized by tension of the fabric. Pressure cuts in automatically at 2 g. B: Shows how the pressure suit is worn under uniform. (Air Ministry Crown copyright.)

FIG. 8.—Initial experiments in pick-up from ground by aircraft in flight at 120 m.p.h. Person to be picked up is crouched facing the oncoming aircraft. Drag from trailing rope of aircraft catches nylon loop fixed between two poles. (Air Ministry Crown copyright.)

Post-War Position

Though there was a lull in researches in connexion with aviation medicine after the first world war, it is most heartening to know that the reverse has been the case since the end of the second world war. Such researches have been intensified as regards both Service and civil aviation, and the R.A.F. Institute of Aviation Medicine at Farnborough has had a considerable increase in staff as well as extension of all its departments, which are fitted with the most up-to-date equipment capable of tackling all current and many future problems. Researches have been actively pursued into space and pace problems of the upper troposphere and the stratosphere, and of supersonic speeds, as well as into the noise and vibration associated with jet-propelled aircraft. Stratosphere investigations include the effect on man of solar and cosmic rays, mesons, and neutrons, and how to protect the body from explosive

decompression should it occur in pressurized aircraft at altitudes over 60,000 feet (18,300 m.), where the blood would boil if some means of compression were not immediately forthcoming.

In addition, the terms of reference of the Flying Personnel Research Committee were extended in 1948 to cover aviation interests other than the R.A.F. and the Fleet Air Arm. The Committee's terms of reference are now to advise the Secretary of State for Air on investigations into the medical aspects of all matters affecting personnel, particularly those which might conduce to safety and efficiency in flying, including research into problems associated with the scientific selection and training of flying personnel and into measures designed to maintain their physiological and psychological efficiency. In addition, the Committee's terms of reference include the following: to advise the Board of Admiralty on these matters in relation to naval aviation; to advise the Ministry of Civil Aviation and the civil corporations on problems of flying staff and passengers of civil aircraft; and to advise the Ministry of Supply on matters in relation to human requirements in flying. The Committee also advises the War Office so far as problems of airborne troops are concerned.

APPRECIATION

This is only a sketchy survey of medical science as applied to the war in the air, but it gives some idea of the part it played in helping to win the war. Those concerned in this work deserve the highest praise for the fine results attained and for the very gallant way in which many of the researches were carried out with great risk to life. Special tribute is due to the excellent work done by physiologists under the guidance of B. H. C. Matthews at Farnborough and to practical psychologists working with Professor F. C. Bartlett at Cambridge, and to those medical officers who were also pilots (termed "flying personnel medical officers") and did liaison duties between aircrews and the laboratory, including two who gave their lives in this great endeavour—Group Captain H. W. Corner and Squadron Leader D. A. H. Robson. These results, however, could not have been attained without the whole-hearted co-operation and pooling of knowledge that existed between biological and material scientists, aircraft designers, and the executive and medical branches of the Allied Air Forces.

The Royal Air Force is greatly indebted to the foresight of Sir Edward Mellanby and Sir Victor Richardson for conceiving and initiating the formation of the Flying Personnel Research Committee, whose activities have proved so valuable in the progress of aviation medicine. Thanks are also due to the Medical Research Council for the help given in certain problems by many of their staff.

REFERENCES

- 1 Air Ministry Pamphlet 129 (1941).
- 2 — 154 (1943).
- 3 — 158 (1943).
- 4 — 165 (1944).
- 5 Air Ministry Publication 3139 (1947). *Psychological Disorders in Flying Personnel of the R.A.F.*
- 6 — 3139B (1950). "Planned Seeing."
- 7 Bartlett, F. C. (1941). Air Ministry Report, Flying Personnel Research Committee, No. 308.
- 8 — (1941). Ibid., No. 488.
- 9 — (1943). Ibid., No. 565.
- 10 — Horn, V., and Mackworth (1945). Ibid., No. 423 (W).
- 11 Bazett, H. C. (1920). *Spec. Rep. Ser. med. Res. Coun. (Lond.)*, No. 53, 184.
- 12 — (1943). Communication to Air Ministry.
- 13 Birley, J. L. (1920). *Spec. Rep. Ser. med. Res. Coun. (Lond.)*, No. 53, 141.
- 14 Browne, R. C. (1941). Air Ministry Report, Flying Personnel Research Committee, Nos. 285 and 292.
- 15 — (1942). Ibid., No. 507.
- 16 — (1943). Ibid., Nos. 418 (a)–(d).
- 17 Carmichael, E. A. (1939). Ibid., No. 79.
- 18 Corner, H. W. (1940). Ibid., No. 134.
- 19 Cotton, F. S. (1941). Ibid., No. 358.
- 20 Craik, K. J. W. (1940). Ibid., No. 188.
- 21 — (1941). Ibid., No. 289.
- 22 — (1941). Ibid., No. 342.
- 23 — (1942). Ibid., No. 415.
- 24 — and Vernon, M. D. (1940). Ibid., No. 193.
- 25 — (1942). *Brit. J. Psychol.*, 32, 206.
- 26 Daly, I. de B. (1940–1). Air Ministry Reports, Flying Personnel Research Committee, Nos. 100, 174, 200, and 238.
- 27 Davidson, S. (1944). Ibid., Nos. 599 and 622.
- 28 Davis, D. R. (1941). Ibid., No. 234.
- 29 Dickson, E. D. D. (1939). Ibid., No. 8.
- 30 — (1939). Ibid., No. 21.
- 31 — (1939). Ibid., No. 55.
- 32 — (1939). Ibid., No. 56 (c).
- 33 — (1939). Ibid., No. 74.
- 34 — (1942). *Proc. roy. Soc. Med.*, 35, 248.
- 35 Ewing, A. W. G., and Littler, T. S. (1939). *J. Laryng.*, 54, 531.
- 36 — (1943). Air Ministry Report, Flying Personnel Research Committee, No. 532.
- 37 Dickson, E. D. D., Simpson, J. F., Fry, D. B., Swindell, G. E., and Brown, R. E. C. (1946). *J. Laryng.*, 61, 139.
- 38 Eggleton, P., Elsdon, S. R., and Hebb, C. O. (1942). Air Ministry Report, Flying Personnel Research Committee, No. 495.
- 39 — (1945). *J. Physiol. (Lond.)*, 104, 129.
- 40 Evelyn, K. (1941). Air Ministry Report, Flying Personnel Research Committee, No. 401.
- 41 Flack, M. F. (1920). *Spec. Rep. Ser. med. Res. Coun. (Lond.)*, No. 53, p. 93.
- 42 — and Bowdler, A. P. (1920). Ibid., No. 53, p. 81.
- 43 Franks, W. R. (1941). Air Ministry Report, Flying Personnel Research Committee, No. 301.
- 44 Gagge, A. P., and Allen, S. C. (1944). *Air Surg. Bull.*, 1, 1.
- 45 — and Marbarger, J. P. (1945). *J. Aviat. Med.*, 16, 2.
- 46 Gilson, J. C. (1942). Air Ministry Report, Flying Personnel Research Committee, No. 444.
- 47 — Macdonald, A. C., and Stewart, W. K. (1946). Ibid., No. 659.
- 48 Goldie, E. A. G. (1940). Ibid., No. 211.
- 49 — (1943). Ibid., No. 551.
- 50 — (1946). *J. Sci. Instrum.*, 19, 23.
- 51 Hartley, C. H., and Young, J. L. (1941). Air Ministry Report, Flying Personnel Research Committee, No. 262.
- 52 Head, H. (1920). *Spec. Rep. Ser. med. Res. Coun. (Lond.)*, No. 53, p. 214.
- 53 Hallenbeck, G. A. (1944). A.A.F. Air Material Command Memo. Rep. No. ENG 49-696-66.
- 54 Livingston, P. C. (1932). *Brit. J. Ophthalmol.*, 16, 577.
- 55 — (1935). *Proc. roy. Soc. Med.*, 29, 59.
- 56 — (1939). Air Ministry Report, Flying Personnel Research Committee, No. 5.
- 57 — (1939). Ibid., No. 80.
- 58 — (1940). Ibid., No. 104.
- 59 — (1940). Ibid., No. 164.
- 60 — (1940). Ibid., No. 222.
- 61 — (1941). Ibid., No. 262 (a).
- 62 — (1941). Ibid., No. 287.
- 63 — (1941). Ibid., No. 297.
- 64 — (1941). Ibid., No. 346.
- 65 — (1942). Ibid., No. 438.
- 66 — (1942). *Brit. J. Surg.*, 29, 339.
- 67 Luczak, A. (1943). Air Ministry Report, Flying Personnel Research Committee, No. 525.
- 68 Lyle, T. K., Macrae, T. F., Yudkin, S., and Gardiner, P. A. (1942). Ibid., No. 473.
- 69 McArdle, B. (1940). Ibid., Nos. 147 and 171.
- 70 — (1940). Ibid., Nos. 169 and 169 (a).
- 71 — (1940). Ibid., No. 173.
- 72 — and Doupe, J. (1940). Ibid., No. 117.
- 73 Macdonald, A. C., and Stewart, W. K. (1946). Ibid., No. 657.
- 74 Matthews, B. H. C. (1939). Ibid., Nos. 60 and 70.
- 75 — (1940). Ibid., No. 128.
- 76 — (1941). Ibid., No. 375.
- 77 — and Roxburgh, H. L. (1941). Ibid., No. 402.
- 78 Macintosh, R. A., and Pask, E. A. (1943). Ibid., Nos. 550 and 550 (a).
- 79 Mackworth, H. N. (1944). Ibid., No. 423 (s).
- 80 — (1944). Ibid., No. 423 (v).
- 81 — (1944). Ibid., No. 586.
- 82 Miles, W. R. (1943). *Fed. Proc.*, 2, 109.
- 83 Ministry Aircraft Production (1941). Air Ministry Report, Flying Personnel Research Committee, No. 249.
- 84 Ministry of Supply (1946). M.O.S., Ballistics LPE. Note No. 130.
- 85 Pask, E. A. (1942). Air Ministry Report, Flying Personnel Research Committee, No. 423 series.
- 86 — (1946). *Maker-up*, 14, 225.
- 87 — Stewart, W. K., and Cowan, S. L. (1943). Air Ministry Report, Flying Personnel Research Committee, No. 511.
- 88 Rook, A. F. (1939). *J. roy. nav. med. Serv.*, 25, 203.
- 89 Royal Air Force Physiology Laboratory (1939). Air Ministry Report, Flying Personnel Research Committee, No. 81.
- 90 — (1940). Ibid., No. 196.
- 91 — (1941). Ibid., No. 232 (b).
- 92 — (1941). Ibid., No. 305 and Air Ministry Pamphlet 120.
- 93 — (1942). Ibid., Nos. 423 (a) and (b).
- 94 — (1942). Ibid., No. 437.
- 95 — (1943). Ibid., No. 521.
- 96 Royal Australian Air Force Med. Service (1941). R.A.A.F. Report No. FP. 27.
- 97 Royal Canadian Air Force Med. Service (1941). Air Ministry Report, Flying Personnel Research Committee, No. 319.
- 98 Stewart, W. K. (1940). Ibid., No. 176.
- 99 — (1940). Ibid., No. 177.
- 100 — (1940). Ibid., Nos. 212 and 213.
- 101 — (1941). Ibid., Nos. 233 and 269.
- 102 — (1941). Ibid., No. 300.
- 103 — (1942). Ibid., No. 364.
- 104 — (1942). Ibid., Nos. 407 and 458.
- 105 — (1946). Ibid., No. 671.
- 106 — (1945). *J. Neurol. Neurosurg. Psychiat.*, 8, 24.
- 107 — and Watt, G. E. (1940). Air Ministry Report, Flying Personnel Research Committee, No. 159.
- 108 — and Winfield, R. H. (1942). Ibid., No. 437.
- 109 — (1945). Ibid., No. 648.
- 110 Stiles, W. S. (1942). Ibid., No. 494.
- 111 Stiles Committee, Flying Personnel Research Committee (1941). Ibid., No. 353.
- 112 Williams, G. O. (1939). Ibid., No. 15.
- 113 Winfield, B. J. O., and Crichton-Miller, C. (1942). Ibid., No. 445.
- 114 Winfield, R. H. (1941). Ibid., No. 355.
- 115 — (1941). Ibid., No. 361.
- 116 — (1942). Ibid., No. 468.
- 117 — (1942). Ibid., No. 493.
- 118 — (1942). *Proc. roy. Soc. Med.*, 35, 257.
- 119 — (1943). Air Ministry Report, Flying Personnel Research Committee, No. 542.
- 120 Wood, E. H., Lambert, E. H., Baldes, E. J., and Code, C. F. (1946). *Fed. Proc.*, 5, 327.

Lady Derby opened a new centre for spastic children at Stockport on January 5, "The Heaton Moor Clinic." It will provide physiotherapy, speech therapy, and other treatment and training for 14 local spastic children.