the localized molecular disruption there results a gradual spread towards the centre. The analogy to fatigue fractures in metals presents itself and would repay further study. It would be interesting to note whether the characters of the fractures in both types are similar. In those of metals the fracture is comparatively smooth at its outer zones where the surfaces have been rubbed together on innumerable occasions prior to the sudden spread of the fracture transversely, this suddenly fractured zone showing relative coarseness. Difficulty remains in affording an explanation for the predominance of this group of fatigue fractures of the fibula in females.

Summary

A case of bilateral fatigue fractures of the fibula in a middle-aged woman is reported.

Point tenderness, and pain at the same site when the ankle mortise was widened by everting the heels or dorsiflexing the foot, were features of the clinical findings.

The site of the fractures was identical with that described by Burrows (1948) in women of middle age.

Treatment advised was stirrup strapping and rest.

REFERENCES

- Burrows, H. J. (1948). *J. Bone Jt Surg.*, **30B**, 266.
Hartley, J. B. (1943). *Brit. J. Radiol.*, **16**, 255.
Richmond, D. A. (1945). *Lancet*, **1**, 273.
Watson-Jones, Sir Reginald (1952). *Fractures and Joint Injuries*, 4th ed. Livingstone, Edinburgh.

INTUSSUSCEPTION OF VERMIFORM APPENDIX IN ALLERGIC CHILDREN

BY

R. G. MITCHELL, M.D., M.R.C.P.Ed., D.C.H.

Lecturer in Child Health, University of St. Andrews; lately Medical Registrar, the Royal Hospital for Sick Children, Edinburgh

Chronic intussusception is relatively rare in childhood and the symptoms are often ill defined, especially when the appendix is the site of intussusception. The following two cases are of interest because the presence of allergic disease caused difficulty in diagnosis. The age of the second patient makes her case additionally interesting, because this is only the second report of intussusception of the appendix in a child under 2½ years of age, the first being that of a boy aged 10 months (Forshall, 1953).

Case 1

A boy of 3 years was sent to the Royal Hospital for Sick Children because of attacks of diarrhoea with blood in the stools. A tentative diagnosis of gastro-intestinal allergy had been made because there was a history of infantile eczema.

After spontaneous delivery at term the child was breast-fed for three weeks and then put on to cow's milk, following which he developed eczema. He had bronchitis at 6 months and measles at 1 year. There was no family history of allergic diseases. The patient was well until seventeen days before admission, when he passed a small quantity of bright red blood per rectum. Two days later he had a little diarrhoea and again passed blood. Treatment with a chalk mixture was prescribed and no further bleeding occurred for ten days. Five days before admission he passed several loose stools containing a larger amount of blood; he looked pale but not ill. On the day of admission he had an attack of abdominal pain and passed more blood in the stool.

Examination on admission showed a poorly nourished child with rough dry skin. A soft systolic murmur was

heard at the apex of the heart, otherwise physical examination was negative. Blood examination revealed a moderate hypochromic anaemia.

On the day following admission he passed a dark stool, and a small, smooth, mobile mass was felt in the right iliac fossa which disappeared and reappeared at intervals during the next two days. He passed no further stools, but complained on one occasion of abdominal pain lasting for a few minutes. On the fourth day a tumour was felt extending from the right iliac fossa to the right hypochondrium, and at operation (Miss R. M. Mackay) a mass was found consisting of the caecum, into which the appendix had completely intussuscepted. There was no obvious impairment of the blood supply, but all layers were so oedematous that reduction was impossible. The caecum was opened, and the completely inverted and very swollen appendix was seen inside; the tip was just becoming gangrenous, but the rest was only congested. The appendix was removed and the caecum closed. Following blood transfusion, recovery was uneventful.

Case 2

A girl, aged 16 months at the first admission, was born spontaneously at term and was fed on dried milk. At the age of 6 weeks she developed infantile eczema, and at 6 months the introduction of egg into her diet resulted in vomiting and urticaria, which recurred whenever eggs were given. She had no other illnesses until 14 months, and there was no family history of allergy.

At 14 months she began to have attacks of abdominal pain and vomiting, lasting for a day or two at a time, and her appetite became poor. She passed normal stools between and during the attacks. On admission to Maryfield Hospital, Dundee, at 16 months of age, she was found to be a well-nourished child with extensive eczema, involving principally the shoulders, chest, and knees. No other abnormality was found and no mass was palpable in the abdomen. The haemoglobin was 11 g.%, the leucocyte count 12,000 per c.mm., and the erythrocyte sedimentation rate was 40 mm. in the first hour (Westergren). The urine was normal. There was mucus and a little red blood in the stools, culture of which showed no pathogenic organisms. The Mantoux test was positive, but x-ray examination of the chest showed only increased broncho-vascular markings.

While in the ward she was inadvertently given an egg custard and had an immediate reaction consisting of generalized urticaria, vomiting, and abdominal pain, which promptly subsided on treatment with an antihistaminic drug.

A provisional diagnosis of gastro-intestinal allergy was made, but a barium enema examination was carried out to exclude any other condition in the bowel. There was a temporary hold-up of the barium in the ascending colon, but eventually normal filling was obtained. The possibility of an intussusception which had been reduced by the enema was considered, but as the symptoms had subsided she was allowed to go home.

She had no more trouble until two months later, when she was readmitted because of further attacks of abdominal discomfort followed by the passage of blood in the stools. Apart from extensive eczema, no abnormality was found on physical examination. A second barium enema showed good filling of the colon but defective filling of the caecum. Gastro-intestinal allergy was still thought to be the most likely cause of her symptoms, and as she was thriving she was discharged from hospital.

One month later she had three more attacks of abdominal colic with blood in the stools, and this time a horizontal thickening was palpable in the upper abdomen. At laparotomy (Mr. W. G. Campbell) a basal appendicular intussusception was found which had further intussuscepted into the colon. The intussusception was easily reduced except for the part formed by the base of the appendix, and the fundus of the caecum with the appendix was removed. Microscopical examination showed diffuse post-inflammatory fibrosis in the subserosa of the appendix, while the mucosa

and submucosa of the caecum showed diffuse and heavy infiltration with eosinophil leucocytes. Post-operative progress was satisfactory, and one year later she was very well, although the eczema was still present.

Discussion

The outstanding symptom of intussusception of the appendix is intermittent abdominal pain, often accompanied by nausea and vomiting, and the stools are usually normal (McSwain, 1941). Many symptoms have been attributed to gastro-intestinal allergy, but intermittent abdominal pain and vomiting are those most commonly mentioned. Blood in the stools is often described as a feature of gastro-intestinal allergy, especially in infants, while it is usually scanty or absent in chronic intussusception (Gaymer Jones, 1924). The difficulties of proving that gastro-intestinal symptoms are allergic have been discussed by Ingelfinger, Lowell, and Franklin (1949), who concluded that absolute confirmation is virtually impossible. They considered such a diagnosis reasonable "if gastro-intestinal symptoms invariably follow contact with a normally inactive substance, and if other causes such as psychic effects and mechanical irritation are excluded." The history of sensitivity to egg of the second patient, and the vomiting and urticaria produced by its administration, favoured the diagnosis of gastro-intestinal allergy, and it seemed reasonable, in the absence of other demonstrable cause, to ascribe the abdominal pain and blood in the stools to allergy as well. Beaven (1929) reported a somewhat similar case, his patient being a girl of 3 years who had florid eczema, and who developed urticaria if she ate eggs. The child had had peri-umbilical pain for one month before laparotomy was carried out. In this child, however, an intussusception was found involving a Meckel diverticulum.

In the present Case 1 there was less justification for attributing the symptoms to allergy, but the history of eczema, the absence of abnormal signs in the abdomen, and the duration of symptoms were misleading. The initial diagnosis was not sustained after admission and the true cause became apparent when the abdominal mass was felt.

Intussusception of the appendix was first recorded in a boy aged 7 (McKidd, 1859), and the majority of cases since reported have occurred in the first decade of life (Forshall, 1953). The condition is usually chronic, the average duration of symptoms before operation being eight months in the 77 cases reviewed by McSwain (1941). Although all cases may not arise in the same way, it is thought that the condition is usually due to peristaltic contractions of the appendix, initiated by attempts to expel a foreign body or by a local thickening of the appendix itself (Fraser, 1943).

It seems possible that in the present cases increased peristalsis resulting from allergic reactions facilitated the occurrence of intussusception, although the eosinophilia in the caecal wall in the second case cannot be adduced as evidence, because the significance of such infiltration is not clearly understood. Moreover, since neither McSwain (1941) nor Fraser (1943) mentions allergy in reviewing the literature on intussusception of the appendix, it appears unlikely to be a common cause, and the association may have been fortuitous.

Summary

Two cases of chronic intussusception of the appendix in young children are reported.

The symptoms were at first attributed to allergic reactions because of a history of infantile eczema in the first child and the presence of infantile eczema and gastro-intestinal allergy in the second.

I am grateful to Professor J. L. Henderson for permission to report these cases, and to Dr. Kenneth Rhaney for the pathological report.

REFERENCES

- Beaven, P. W. (1929). *Amer. J. Dis. Child.*, **37**, 373.
Forshall, I. (1953). *Brit. J. Surg.*, **40**, 305.
Fraser, K. (1943). *Ibid.*, **31**, 23.
Ingelfinger, F. J., Lowell, F. C., and Franklin, W. (1949). *New Engl. J. Med.*, **241**, 303, 337.
Jones, J. Gaymer (1924). *Brit. J. Surg.*, **12**, 378.
McKidd, J. (1859). *Edinb. med. J.*, **4**, 793.
McSwain, B. (1941). *Sth. med. J.*, **34**, 263.

SURGERY IN OCTOGENARIANS

BY

RONALD W. RAVEN, O.B.E., F.R.C.S.

Surgeon, Westminster (Gordon) Hospital; Surgeon,
the Royal Marsden Hospital

The problems caused by lesions in aged patients which constitute either an immediate or a delayed threat to life require careful consideration if an adequate solution of them is to be found. It is recognized that the number of these patients in this country is gradually increasing, and these matters will assume greater proportions in the years to come. A study of a personal series of 34 octogenarians who underwent surgical operation demonstrates that much can be done to help these aged patients even though they are afflicted with the worst forms of malignant disease. The patients were divided into two groups: with malignant and with non-malignant lesions. Their details are set out in Tables I and II.

There were 24 patients with malignant disease; for the majority a major operation was required, and four died. The follow-up of the survivors is shown and six patients were still alive up to twenty-nine months after the operation. Three patients died of non-malignant causes 3 years, 4½ years, and 4 years 5 months later. Of those who died of cancer four lived for a minimum period of two years.

Distribution and Management of the Lesions

In 15 patients a carcinoma of the alimentary tract was present, and in 9 it was situated in the colon or rectum. One patient suddenly developed a perforation of the pelvic colon proximal to an annular constricting carcinoma. Another suffered from complete intestinal occlusion caused by carcinoma of the pelvic colon.

For an aged patient with an obstructing carcinoma of the transverse or pelvic colon Paul's operation is a satisfactory procedure, provided any involved regional lymph nodes can be removed at the same time. When the affected loop of bowel cannot be exteriorized—for example, in the lower part of the pelvic colon, or at the recto-sigmoid junction—Hartman's operation is useful. For non-obstructing lesions of the transverse or distal colon a partial colectomy with an end-to-end anastomosis is usually radical enough rather than a more extensive left hemicolectomy. When the carcinoma affects the right half of the colon a right hemicolectomy in one stage is advisable if there is no obstruction.

In one patient a large pelvic abscess complicated a carcinoma of the pelvic colon and caused severe toxæmia. A series of staged operations were performed, including drainage of the abscess, the institution of a defunctioning transverse colostomy, partial colectomy, and closure of the colostomy later. The timing of these staged operations is important, so that the patient's general condition is at its optimum when the next stage is performed. By this means many aged patients, even when seriously ill, can be brought through safely. It is worth noting that a palliative colostomy for a carcinoma of the rectum in two patients was followed by survival periods of 2½ and 3½ years. Another patient lived 4½ years after a perineal excision of the