EMERGENCIES IN GENERAL PRACTICE**SHOCK***

BY

Sir HENEAGE OGILVIE, K.B.E., M.D., M.Ch., F.R.C.S.*Consultant Surgeon, Guy's Hospital*

The study of shock has for many years attracted the ablest investigators in Britain and America. Men like Aub, Blalock, Bywaters, Grant, Green, Gregerson, Haist, Moon, Phemister, and Wilson (the order is alphabetical rather than chronological) have paid particular attention to this problem, and by their research work in the laboratory and their clinical observations at the bedside they have added greatly to our knowledge of the reactions of the animal and human body to various forms of injury.

The treatment of shock is essentially a practical problem. We must study the research work that has been done in the laboratory, and if it has provided an answer we must apply the lesson it brings to our clinical tasks. Very often, however, the answer is "So what?" and it is extremely important that we should leave philosophical doubt behind when we come to the side of the pale listless man who has just been rushed in by the ambulance. It is right, therefore, that from time to time a clinical speaker should try before a clinical audience to bring this important subject from the laboratory to the bedside, to turn aside from a consideration of what shock might be and consider what shock is as we see it, what steps we can take to prevent it, and what we can do to save the shocked patient.

What is Shock?

We are often criticized by our scientific friends for using the word "shock" without explaining exactly what we mean by it. The word "shock" has a very definite meaning to me, and to most surgeons. To me it means a state of collapse following injury, and I shall go on calling it shock till I find I am wrong. The definitions of the experts are no more helpful than this simple statement. Blalock defines shock as "peripheral circulatory failure." Why peripheral? Why failure? In the type of shock most commonly encountered in clinical practice, that due to haemorrhage or trauma, nothing has failed. The central pump and the peripheral mechanism are working with supreme efficiency. The motor is spluttering only because the tank is running dry. Moon defines shock as "a circulatory deficiency characterized by decreased blood volume, decreased cardiac output, and increased concentration of the blood." This may be the shock of the laboratory mouse. It is not the shock of the wounded soldier; unless he is burned or crushed his blood concentration is normal or below that figure, and it falls in subsequent days. These definitions, in attempting to give an exact meaning to a comprehensive and general term, force upon it a significance foreign to its accustomed clinical use. This is hardly surprising, since many of the experiments whereby shock is produced in the laboratory bear no relation to the manner of its causation in surgical practice.

I will return to my first definition of shock and expand it slightly. "Shock is a state of collapse following injury, characterized by a depression of the circulation." All we know when we are called to the patient's side is that he has been injured, that he is collapsed, and that he lies apathetic

with an ashen grey face and cold clammy hands. The diagnosis is a clinical one, and, having made it, we must set ourselves urgently to seek the cause and decide the treatment.

Mechanism of Shock

The circulation of the blood depends on three things: on the action and force of the heart beat, on the volume and viscosity of the blood, and on the peripheral resistance of the arterioles. Interference with one, two, or all of them may be found in the condition of acute circulatory depression known as shock. Heart failure may resemble shock. Many factors associated with injury, particularly with battle injury—peritonitis, blast damage to the lungs, fat embolism, acute anaerobic infection, absorption of toxins from damaged muscle—may complicate and be partly responsible for the clinical state of shock. But to label them as "shock" is to make the term completely meaningless.

Traumatic shock is, in the overwhelming majority of cases, the result of a diminution of the effective circulating volume of the blood, and the word "shock" should be used for this condition alone. The other shock-like states should be called pseudo-shock. The later changes caused in the circulatory mechanism by anoxaemia, infection, or tissue toxins of various sorts should be referred to as epiphenomena or sequelae of shock, but not as shock.

In shock the diminution of volume may be due to actual loss, or to a sudden increase in the capacity of the vascular bed, the volume being unchanged or diminished to an extent that would not in itself cause shock. In wound shock the loss is one of blood, either from the wound or into damaged tissues. In the shock which follows crush lacerations or severe surgical operations the loss is of blood and plasma. In shock from burns the loss is of plasma only, from the burned surface and into the tissues underlying it. In the vasovagal syndrome, known also as primary or nervous shock, a state which may be found alone or preceding or accompanying true oligæmic shock, the circulating volume is inadequate to the vascular bed, whose capacity is increased by widespread dilatation.

Primary and Secondary Shock

Phemister has observed that "primary and secondary shock should be recognized on the basis of difference in aetiology rather than difference in time of occurrence." Primary, or nervous, shock is a condition of circulatory disturbance associated with low blood pressure due to dilatation of the peripheral arterioles, and a slow pulse due to stimulation of the vagi. It is a cardiovascular upset of reflex origin, the result of painful stimuli or of emotional disturbance, very often a combination of the two. A painful injury is more likely to cause it than a large one, and an apprehensive individual is more likely to suffer from it than a phlegmatic one. The main features distinguishing primary from secondary shock, apart from the nature of the causative injury, are that the patient does not look ill, that his skin is warm, that he is torpid or unconscious rather than apathetic, and that his pulse is slow.

Primary shock is essentially a transient phenomenon that responds to rest, relief of pain, warmth, hot drinks, and removal from distressing surroundings, and a good response to these simple measures may be the surest method of distinguishing primary from secondary shock. It is, however, less important to recognize those who respond well than it is

* Based on a contribution to a symposium at Bristol University on "Medical Aspects of Atomic War" (see *Journal*, November 27, 1954, p. 1283).

to watch anxiously for those who do not or who relapse after temporary improvement, for they are suffering from secondary shock and need immediate and active treatment. If there is any doubt, needless transfusion of a patient in primary shock is a harmless mistake compared with delay in starting the transfusion of one suffering from secondary shock.

Secondary shock must be assumed where there has been much blood loss or tissue damage. Some estimate of the loss can often be made from the history given by the patient or his friends, or from observing the amount of blood on the clothing or the stretcher. But fluid may be lost into the tissues without external loss, a mechanism repeatedly stressed by Blalock. A man can bleed to death into the thigh muscles round a fractured femur without any free haemorrhage occurring. Grant, in his observations on wound shock in the Italian campaign published in 1951 in the Medical Research Council Special Report Series, No. 277, has pointed out that the amount of blood and plasma lost into the damaged tissues in extensive muscle injuries is far greater than had been previously realized, and that the loss continues and is responsible for a relapse into clinical shock after resuscitation. The view that shock leads to endothelial damage, with loss of fluid into tissue spaces throughout the body and concentration of corpuscles in the peripheral capillaries, is no longer held.

Assessment of the Shocked Patient

A man suffering from shock looks ill. His attitude is apathetic, but he is mentally alert and answers questions readily. His skin is cold, moist, and pale, with a greyish colour, particularly about the ears and lips. His hands and feet are icy cold. His eyes, with pupils dilated, appear abnormally bright. His respirations are little altered. The pulse is rapid and of low tension. He passes little urine, and is prone to vomit any fluids given by mouth. Air-hunger and spontaneous vomiting are indications of severe shock.

An index to the presence and degree of shock, but not an entirely reliable one, is the blood pressure. While the fall in systolic pressure is usually commensurate with the loss of blood volume and the degree of shock, there are surprising exceptions. Some men show a low blood pressure rising subsequently, some show periodic rises and falls, and some maintain a high pressure till just before the end. A low pulse pressure is often of more significance than a low systolic pressure.

The whole clinical picture of shock can usually be accounted for by the loss of circulating fluid alone. The degree of shock is roughly proportional to the loss of fluid: a loss of 2 pints (1 litre) leads to moderate shock, one of 3 pints (1.5 litres) to severe shock, one of 4 pints (2 litres) (roughly 40% of the blood volume) to profound shock. A known loss of 4 pints (2 litres), even though the patient may, by supremely efficient compensation, be able temporarily to keep his blood pressure up, will eventually lead to profound shock, and, conversely, profound shock means that at least 4 pints (2 litres) have been lost and must be replaced.

Restoration of the Circulation

The treatment of oligæmic shock is the restoration of the circulating volume to its former level. If possible, the volume should be restored qualitatively as well as quantitatively—that is, lost blood should be replaced with blood, and plasma with plasma—but the immediate need is for volume, and the fluid must usually be introduced directly into the circulation. Water by mouth or by rectum or crystalloid fluids intravenously are of little value when the circulatory loss has been large enough to give rise to clinically recognizable shock; under such conditions water is poorly absorbed in the alimentary canal and poorly retained in the circulatory system.

The restoration of volume should be early and rapid. While a patient is in a state of shock, all his tissues are

poorly supplied with oxygen; organs such as the kidneys and liver are by-passed to conserve blood for the heart and brain and may suffer permanent damage, and bacteria in his wounds, particularly the anaerobes, are given a chance to establish themselves. When a total of 4 pints (2 litres) or more is needed, the first two should be run in rapidly, using pressure to assist gravity, and full restoration in four hours should be aimed at.

The choice between blood and plasma depends to some extent on the availability of the two. Plasma can always be obtained, blood only when donors have been prepared, and blood in quantity only when arrangements have been made to supply it. Plasma is free from the dangers of reaction due to incompatibility, infection, or mechanical destruction of the red cells which are present with blood, but may transmit the virus of hepatitis. It is better than blood for the shock of burns, indeed for all shock characterized by haemoconcentration. In haemorrhage it is as good as blood for restoring volume, and not greatly inferior in therapeutic results when the blood loss has been moderate. It flows more easily, so that even in transfusion for severe haemorrhage it has been advised that plasma should be used for the first 2 pints (1 litre), which are given rapidly. Blood should be used to replace a haemorrhage of more than 2 pints (1 litre), and when the haemorrhage is not merely severe but is expected to continue during resuscitation or operation, blood alone should be used. Plasma cannot bring a man who has lost more than 2 pints (1 litre) of blood into a state fit to withstand a severe operation, or enable him to overcome infection or to repulse anaerobes.

"Plasma-savers" are solutions containing large molecules—of gum acacia, a polyvinyl compound, or a polysaccharide—that remain in the circulation and can replace in volume and viscosity an equal amount of blood. What is their place in the treatment of shock? The answer is in the seventh chapter of St. Matthew's Gospel: "What man is there of you, whom if his son ask bread, will he give him a stone?" The plasma-savers provide volume and viscosity and can restore the blood pressure, but they do not carry oxygen, and they do not provide energy or any substance used by the body. On the other hand, they tend to clump the red cells, thus interfering with grouping, and they are finally fixed by the reticulo-endothelial system. By restoring the blood pressure before the cause of hypotension has been arrested, they may restart a haemorrhage that has stopped and wash out the life fluid with a fake substitute.

There is one legitimate place for the plasma-savers: to keep a patient alive in a desperate emergency till blood or plasma can be obtained. If a man has lost 5 pints (2.8 litres) of blood rapidly, or if his systolic pressure has dropped to 50 mm., his central nervous system and his kidneys will suffer irreparable damage and his acid-base equilibrium will be upset beyond recovery unless the pressure can be raised within half an hour. Plasma-savers must be given till blood arrives, for desperate situations demand desperate remedies.

I hold that every surgeon who is called upon to treat acute injuries in healthy men should act upon the belief that no haemorrhage need be fatal if it can be stopped and the lost blood can be replaced early enough and fast enough, and no shock need become irreversible if it can be reversed in time and with sufficient energy. Shock becomes irreversible chiefly because prolonged anoxia has been allowed to wreck the machinery of recovery. Many more lives are lost by tardy, hesitant, or inadequate transfusion than by excessive transfusion or transfusion for the wrong indications.

Ancillary Measures

I must touch briefly on some of the other measures used in the treatment of shock. Warmth may be beneficial in primary shock, but in secondary shock it is harmful. Experimentally shocked animals are more likely to die in a warm room than in a cold one. The cold skin of shock is an index of a peripheral vasoconstriction the purpose of which is to reduce the volume of the vascular bed and to maintain

the arterial pressure, and a drop in body temperature is beneficial in reducing the general metabolism to a level more adapted to the reduced oxygen-carrying capacity of the blood. The "shock-cradle" can be a shock producer. Patients expend their ebbing strength in trying to throw it off, and in this demonstrate that the physiological call of instinct is wiser than the teaching of the textbooks.

Morphine must be used with great caution. No more than $\frac{1}{4}$ gr. (16 mg.) should be given at a time, and it should be given into a vein, since in a state of shock it is absorbed slowly from the subcutaneous tissues. A second and larger subcutaneous dose is apt to be given if the first fails, and when later the circulation is restored the accumulated amount is absorbed and may produce a profound stupor.

The use of pressor drugs, including the adrenal preparations, is unwise. They may do good in primary shock accompanied by vasodilatation, but here they are unnecessary. In oligæmic shock, which is associated with vasoconstriction, the real need is for an increased supply of blood and oxygen to the tissues rather than for a raised systolic pressure, and further vasoconstriction may therefore do more harm than good.

When Normal Services Fail

The treatment I have discussed demands, not necessarily hospital conditions, but at any rate facilities for careful examination and constant supervision, the provision of shelter and bed-clothing, and a supply of the necessary fluids, drugs, and apparatus.

Should a casualty be moved to hospital before shock has been combated? A man in profound shock can easily be killed by a rough journey, particularly if his fractures are not adequately splinted. But part of the shock seen immediately after an accident is nervous in origin, and the patient will recover considerably after a short period of rest during which he is kept warm, reassured, and given fluids. The apparatus for intravenous fluid replacement is seldom available at the scene of an accident, but it may arrive with the ambulance. Should blood or dextran be given before the casualty is loaded for the journey? The answer depends on the state of the patient, the nature of the injury, and the length and quality of the coming journey. When in doubt it is better not to transfuse, for transfusion without control of the blood loss is apt to increase that loss, particularly if it raises the blood pressure above shock level during a bumpy trip. However, if the bleeding can be controlled by ligature or tourniquet, it may be wise to start transfusion before the man is put into the ambulance, and to send him off, with the transfusion running, as soon as a palpable pulse has returned.

What should be done in a major disaster when normal services are absent, inadequate, or disrupted? Three things that can be given to all are reassurance, warmth, and fluids. Reassurance alone will restore most cases of primary shock and benefit all. No one can dispense it better than the doctor, who by his quiet confident actions can do most to assure the victims that the crisis is over and the situation is in hand. Warmth and fluids must be dispensed with more caution, but here, fortunately, the customer is always right. The man who feels cold needs blankets. The man who is cold and does not feel it is better left uncovered. With water, also, personal desires can usually be trusted; the man who asks for a drink can usually be given one, the man who does not want water or vomits when it is given him should be left alone or given fluids another way. Pain must be allayed if the means are available, and morphine, given intravenously, is the best analgesic in shock. The administration of blood plasma and dextran will depend on whether these fluids are available; they cannot be improvised on the spot.

We must always remember that even the most severe shock is not necessarily fatal. Men have recovered after lying out all night on an icy battlefield who might have died had they been found earlier and subjected to the additional trauma of movement. The one thing that must

be remembered is that a shocked man will not stand an anaesthetic or an operation, and if the means of treating shock actively are not available he will stand a better chance if nothing is done to his injuries beyond splinting fractures, stopping haemorrhage, dressing his wounds, and keeping infection at bay by the administration of sulphonamides at regular intervals, than if he is subjected to surgical excision of wounds and mechanical cleansing of burns while still in a state of shock.

Next article on Emergencies in General Practice.—
"The Differential Diagnosis of Coma," by Dr. M. J. Parsonage.

Refresher Course Books.—Copies of the first two volumes of collected articles from the Refresher Course for General Practitioners published in the *Journal* are still available at 25s. (postage 1s.) each. The first volume contains 55 articles and the second 60. Each article has been revised and brought up to date by its author.

These books may be obtained from the Publishing Manager, B.M.A. House, Tavistock Square, London, W.C.1, or through any bookseller.

BACTERIAL GENETICS

[FROM A SPECIAL CORRESPONDENT]

An informal meeting on bacterial genetics was held at the London School of Hygiene and Tropical Medicine from December 14 to 17, 1954. The intention was to provide reviews of the main topics in terms comprehensible to those not well acquainted with the subject, and for the most part the speakers succeeded reasonably well.

Experimental Production of New Strains

The first day was devoted to some of the methods by which two bacterial strains can interact to produce a new "hybrid" strain. In the pneumococcus, the influenza bacillus (*Haem. influenzae*), and the meningococcus, cells treated with deoxyribonucleic acid (D.N.A.) extracted from some other strain of the same species may acquire some heritable character from that strain. Many different traits can be transferred in this way; and it has recently been discovered that some pairs of traits are linked—that is, that absorption of a single particle of D.N.A. by a bacterium may confer on it two distinct new hereditary properties. These facts make it at least very probable that D.N.A. constitutes the genetic apparatus of the bacteria concerned, and that single macromolecules of D.N.A. contain differentiated regions regulating different bacterial characters—that is, different genes. The latest development in this field was reported from the Biophysics Research Unit at King's College, London. By chromatography on a column of histone adsorbed on kieselguhr it has been possible to divide D.N.A. from streptomycin-resistant pneumococci into a series of fractions which differ not only in their chemical composition (ratio of bases) but also in their ability to transform streptomycin-sensitive pneumococci into resistant ones. It remains to be seen whether the ability to effect different transformations will be found concentrated in a single fraction or distributed amongst different ones. In the latter case the isolation and chemical characterization of a specific gene may not be quite so remote a possibility as it has hitherto seemed.

The recently discovered phenomenon of transduction of characters from one salmonella strain to cells of another by treatment with a cell-free filtrate of a lysed culture now appears in essence similar to the D.N.A.-mediated transformations; the main difference being that in salmonella the genetic material requires a phage particle to carry it across from the lysed cells to the cells of the recipient strain. The chemical nature of this phage-transported salmonella