hope that the foreign body would be vomited, a dangerous practice when pointed or sharp objects have been swallowed. Furthermore, children who have swallowed foreign bodies should be admitted to hospital without delay so that skilled surgical treatment is at once available should the necessity arise.

The danger of delayed perforation by a hair-grip from pressure necrosis of the bowel wall will be only too apparent, especially if the grip has been impacted for any length of time.

RALPH H. GARDINER, D.M., M.Ch.,
F.R.C.S., F.R.C.S.Ed.,
Surgeon, Aylesbury.

Effect of "Venostasin" on Serum Cholesterol Levels

In recent years attention has been drawn to the correlation in some instances between the serum cholesterol level and the occurrence of atheroma. The subject has been reviewed in the *Annals of Internal Medicine* (1950) and more recently by Gutman (1953). In 1953 Shimert and Schwarz reported that an extract of the horse chestnut, "venostasin," caused a significant reduction in the serum cholesterol level, which fell in 25 subjects from a mean level of 204 mg. per 100 ml. to 154 mg. after five daily intravenous injections of 5 ml. They found the extract inactive when given by mouth, but no toxic effects were noted after as many as 50 daily intravenous injections of 5 ml. each. The mode of action of venostasin is unknown, but if this effect is confirmed the material might prove of value in studying factors which influence the serum cholesterol level and in treating or preventing atheroma.

METHOD

Six patients were studied by a "double-blind" technique using venostasin and an inert solution of burnt sugar indistinguishable in appearance from the venostasin. Neither the analyst nor the person giving the material knew whether the patient was receiving the horse chestnut extract or the inert solution. After a control period of three days the patients were given a daily injection of 5 ml. of one substance for five or ten days, followed by the other substance for a similar period of time. Blood was collected in the fasting state at approximately the same time each morning, and the cholesterol level estimated by a modification (MacIntyre and Ralston, 1954) of the method of Zlatkis *et al.* (1953). Throughout the period of study the patients received an ordinary ward diet.

RESULTS

The results are shown in the Table and two illustrative examples in the Figure. In all six patients venostasin had no

Failure of venostasin or a solution of burnt sugar ("inert substance") to influence the serum cholesterol level in two patients.

detectable influence on the serum cholesterol level, whether this was in the high or in the low normal range.

Effect of Venostasin and an Inert Substance on the Serum Cholesterol Level

Case No.	Diagnosis	Average Cholesterol (mg./100 ml.)		
		Control Period	Venostasin Period	Inert Substance Period
1	Essential hypertension ..	193	191	193
2	Peptic ulcer ..	188	197	198
3	Coronary sclerosis ..	339	327	340
4	Essential hypertension ..	312	297	326
5	Coronary sclerosis ..	228	198	191
6	Mitral stenosis ..	183	201	218

We are indebted to Dr. R. I. S. Bayliss, at whose suggestion and on whose patients this investigation was carried out.

DOREEN AILWARD,
D. HYWEL DAVIES, M.A., B.M.,
R. HECKER, M.B., M.R.A.C.P.,
Departments of Medicine and Biochemistry,
Postgraduate Medical School of London, W.12.

REFERENCES

- Annals of Internal Medicine*, 1950, 33, 251.
Gutman, A. B. (1953). *Amer. J. Med.*, 14, 1.
MacIntyre, I., and Ralston, M. (1954). *Biochem. J.*, 56, xliii.
Shimert, G., and Schwarz, K. (1953). *Klin. Wschr.*, 31, 1063.
Zlatkis, A., Zak, B., and Boyle, A. J. (1953). *J. Lab. clin. Med.*, 41, 486.

Nitric Oxide Haemoglobin in Urine Due to Contamination

Nitric oxide haemoglobinaemia is probably incompatible with life. Banham, Haldane, and Savage (1925) described the finding of nitric oxide haemoglobin in the blood of a patient who died of pneumonia; they postulated a nitrifying organism in the blood as the agent producing the nitrification, though they did not confirm the identity of the abnormal pigment by spectroscopic examination. Barton (1954) described a fatal case of nitric oxide haemoglobinaemia due to sodium nitrite poisoning.

Nitric oxide haemoglobinuria has not been described. Below we report the identification of the pigment in urine, although the pigment was an artifact and not present in the patient.

A dark urine, which had been collected from a female patient in an open container eight hours previously, was examined in the laboratory. Nitric oxide haemoglobin was identified by spectroscopic and chemical tests (Harrison, 1947). As the patient was not desperately ill, but was suffering from stone in the urinary tract, the urine specimen was re-examined. It was found to be heavily infected with a mixed growth—one organism was a member of the Bacillus group. A catheter-specimen of urine was sterile. The mixed bacterial growth was cultivated in normal urine to which a little fresh blood had been added. The nitrifying action of the bacteria produced nitric oxide haemoglobin within 24 hours. The contaminated mixture was examined by a bacteriologist, but he was able to give a no more detailed identification than the one here presented.

The conclusion reached was that the patient's urine originally contained blood: the haemoglobin was converted to nitric oxide haemoglobin as a result of bacterial contamination from the faeces, the container, or the air.

We wish to thank Professor Sir Charles Dodds for his interest in this case.

D. N. BARON, M.D.,
Department of Chemical Pathology,
Royal Free Hospital.
C. F. M. ROSE, A.R.I.C.,
S. A. Courtauld Institute of Biochemistry,
Middlesex Hospital Medical School.

REFERENCES

- Banham, H. A. L., Haldane, J. S., and Savage, T. (1925). *British Medical Journal*, 2, 187.
Barton, G. M. G. (1954). *Lancet*, 1, 190.
Harrison, G. A. (1947). *Chemical Methods in Clinical Medicine*, 3rd ed. London