- Flynn, F. V., Harper, C., and de Mayo, P. (1953). *Lancet*, 2, 698.
 Garrod, A. E. (1923). *Inborn Errors of Metabolism*, 2nd ed. London.
 Greenwald, I. (1930a). *J. biol. Chem.*, 88, 1.
 — (1930b). *Ibid.*, 89, 501.
 — (1931). *Ibid.*, 91, 731.
 Hagedorn, H. C., and Jensen, B. N. (1923). *Biochem. Z.*, 135, 46.
 Horrocks, R. H., and Manning, G. B. (1949). *Lancet*, 1, 1042.
 Hough, L., Jones, J. K. N., and Wadman, W. H. (1950). *J. chem. Soc.*, 2, 1702.
 Janeway, T. C. (1906). *Amer. J. med. Sci.*, 132, 423.
 Larson, H. W., Blatherwick, N. R., Bradshaw, P. J., Ewing, M. E., and Sawyer, S. D. (1937). *J. biol. Chem.*, 117, 719.
 — Chambers, W. H., Blatherwick, N. R., Ewing, M. E., and Sawyer, S. D. (1939). *Ibid.*, 129, 701.
 Lasker, M. (1950). *Amer. J. clin. Path.*, 20, 485.
 — and Enklewitz, M. (1933). *J. biol. Chem.*, 101, 289.
 — Enklewitz, M., and Lasker, G. W. (1936). *Hum. Biol.*, 8, 243.
 Levene, P. A., and La Forge, F. B. (1914). *J. biol. Chem.*, 18, 319.
 Margolis, J. I. (1929a). *Amer. J. med. Sci.*, 177, 348.
 — (1929b). *J. Amer. med. Ass.*, 93, 173.
 Minot, A. S., Frank, H., and Dziewiatkowski, D. (1949). *Arch. Biochem.*, 20, 394.
 Orr, W. F., and Minot, A. S. (1952). *Arch. Neurol. Psychiat. (Chicago)*, 67, 483.
 Partridge, S. M., and Westall, R. G. (1948). *Biochem. J.*, 42, 238.
 Peterman, M. G. (1945). *J. Pediatr.*, 26, 296.
 Rabinowitch, I. M. (1925). *J. clin. Invest.*, 2, 457.
 Salkowski, E., and Jastrowitz, M. (1892). *Zbl. med. Wiss.*, 30, 337.
 Schmidt, O. T., and Treiber, R. (1933). *Ber. dtsch. chem. Ges.*, 66, 1765.
 Wolfson, W. Q., Cohn, C., and Devaney, W. A. (1949). *Science*, 109, 541.

TRUE HERMAPHRODITISM

BY

A. F. BROMWICH, M.B., F.R.C.S.

Surgical Registrar, Charing Cross Hospital

An individual who has both ovarian and testicular tissue is a true hermaphrodite. Proof of the presence of both types of gonad in the same individual is histological, as the secondary sex characters may be affected by other factors, such as overactivity of the adrenal cortex.

Adopting these criteria, an examination of the literature has disclosed 59 unequivocal cases of true hermaphroditism, to which two further cases are added below. The further progress of the case originally recorded by Broster *et al.* (1938) is also noted.

Case 1

J. O., the third child of healthy parents of English extraction, was first seen at the age of 16 months, having been brought for determination of the sex. No family history of any endocrine disease or congenital abnormality was forthcoming, except for talipes in the second child. The mother had suffered no illness during the pregnancy.

On examination the child appeared healthy and apparently normal except for the genitalia, which showed the following abnormalities. The phallus, about 3 cm. long, curved downwards and had a hooded prepuce. About 3 cm. behind the phallus lay an orifice into which opened the urethra and a short recess. This recess admitted a probe for 2.5 cm. and presumably represented the lower part of the vagina. The labio-scrotal folds were rugose and contained no testes or hernial sacs.

Investigations.—Urinary 17-ketosteroids, 0.3 mg. a day (this is at the lower limit of assay, and indicates only that the 17-ketosteroid excretion was not increased). X-ray examination of the pituitary fossa showed no abnormality. The epiphysal bone age was 2–3 years—that is, about one year more than the actual age.

Operation.—Mr. L. R. Broster performed a laparotomy on July 17, 1953. A small (4 cm.) uterus was observed, from which ran two Fallopian tubes, terminating in fimbriated ends. In the usual position of the right ovary lay a structure about 2 cm. long, resembling a testis, a portion of which was removed for microscopical examination. No epididymis or other Wolffian elements could be found on either side. The adrenal glands felt normal in size and consistency. Samples of skin were taken from both sides of the abdomen for examination of the somatic nuclei by the method described by Moore *et al.* (1953). Dr. W. E. Evans reported

that the whole of the piece of tissue (8 by 3 mm.) from the right gonad consisted of immature testicular tissue, with seminiferous tubules separated by a loose, clear-celled stroma. Interstitial cells were not identified. A narrow fibrous capsule lay along one edge of the biopsy. The specimen, about 3 by 3 mm., from the left gonad consisted of a strip of young ovarian tissue beneath which was loose fibrous tissue. No ovarian "germinal" epithelium was found on the surface. Several ova were present, each surrounded by rather flattened cells with deeply stained nuclei, apparently derived from the ovarian stroma.

Skin Biopsies.—Professor M. L. Barr reported that a chromatin mass was present in less than 10% of epidermal nuclei. It has been observed (Moore *et al.*, 1953) that, in sections of tissue taken from a normal female, more than 80% of the nuclei seen contain a speck of chromatin on the periphery of the nucleus. In normal males such a mass is seen in less than 10% of nuclei in tissue sections. This observation is in accordance with the difference between the female sex chromosome pattern (XX) and the male pattern (XY). The nuclear picture in this case therefore conforms to the male type.

Case 2

P. F. was first seen in July, 1940, at the age of 20 months. The child had been brought up as a girl, but was already rougher and more masculine in behaviour than the elder sister, who was a normal girl. The patient was a muscular child with a deep voice and of normal intelligence. The phallus was about the normal size for a male of that age, but the urethra terminated at its base. No gonads could be felt in the scrotum, which was split, and a small uterus was palpated rectally.

Investigations.—Urinary 17-ketosteroid excretion, 0.3 mg. a day. Perirenal insufflation gave no evidence of adrenal enlargement.

Operation.—Laparotomy was performed on September 1, 1940, by Mr. L. R. Broster. A small infantile uterus with fimbriated Fallopian tubes was found. Two gonads lay in the position of the ovaries. The left gonad was the size of a pea and the right gonad was slightly larger. No epididymis or vas deferens was seen on either side. Both gonads were removed for microscopical examination.

Professor H. W. C. Vines reported that the left gonad consisted of ovarian tissue and the right gonad of fibrotic testicular tissue.

The child was clearly more male than female in its behaviour, and was therefore brought up as a boy. From the age of 4 years a series of operations were performed to correct the hypospadias. Injections and local inunction of testosterone from the age of 6 years have produced some increase in the size of the penis, which was, however, still smaller than normal when he was last seen in 1951 at the age of 13. He was then psychologically male, but rather shy, tending to cling to his mother.

Unfortunately, it has not proved possible to get in touch with the patient in order to obtain a sample of skin for nuclear examination.

Case 3

B. T. (Broster *et al.*, 1938) was regarded as a female at birth. At the age of 3 years, however, the child's social and legal sex was changed to male. Two years later hair began to grow on the face, legs, and pubis, and the voice cracked. When seen at the age of 9 years the child had a small penis with complete hypospadias and a split scrotum containing no testes. He was psychologically male. At laparotomy on November 30, 1936, at the age of 9, an infantile uterus with tubes was present. The gonads, which were in the position normally occupied by the ovaries, were shown by biopsy to be both ovotestes.

The patient's hypospadias was successfully repaired at the age of 14, and since then injections and implants of testosterone have been given to encourage penile growth.

When last seen, in January, 1954, he had the appearance of a normal adult male except for a small penis and no

testes in the scrotum. The nipples and areolae have a brown pigmentation, but there is no gynaecomastia. Psychologically, he is a normal male and has been married for over a year. Intercourse has been achieved with some difficulty owing to the small size of the penis. Professor Barr has, however, reported that the nuclear picture in skin samples from both sides of the body is that of a normal female. The chromosomal determination is therefore at variance with the psychological sex.

Discussion

Diagnosis.—In many of the reported cases of true hermaphroditism the diagnosis was made incidentally during operation for other conditions such as hernia, or at post-mortem examination. The question of clinical diagnosis only arises in those patients brought for opinion because of genital malformation and consequent doubt about the sex. Such patients may be female or male pseudo-hermaphrodites or, much more rarely, true hermaphrodites. The female (adrenal) pseudo-hermaphrodite presents clinically with an enlarged clitoris and a uterus palpable on pelvic examination. There may be a history suggesting Addisonian crisis. The diagnosis is suggested by a raised urinary 17-ketosteroid level and confirmed by demonstration of enlarged adrenal glands, usually at laparotomy. The male pseudo-hermaphrodite or intersex with hypospadias and cryptorchidism will have no uterus and a normal urinary 17-ketosteroid level for his age. These two conditions may also be differentiated by examination of the epidermal nuclei, as outlined above. Most cases of true hermaphroditism have a uterus, and a phallus intermediate in size between penis and clitoris. The 17-ketosteroid level is, however, not raised. The diagnosis is made finally by laparotomy and biopsy of the gonads.

Classification.—From the published cases it appears that in true hermaphroditism either gonad may be a testis, an ovary, or ovotestis, independently of the condition found in the other gonad. The various possible combinations have been classified by Hinman (1935) as follows :

Type	One Side	Other Side	Reported Cases
Lateral	Testis	Ovary	22
Unilateral	Ovotestis	Ovary or testis	17
Bilateral	"	Ovotestis	14
(Indeterminate)	"	Not sectioned	8

It will be seen that Cases 1 and 2 reported here are lateral hermaphrodites and Case 3 is a bilateral hermaphrodite. In the majority of lateral cases reported the adnexae on each side corresponded to the gonad on that side. In the case of Brachetto-Brian *et al.* (1943) the asymmetry involved the whole body, one half being male and the other half female. In Cases 1 and 2 above, and that reported by Davis and Scheffey (1946), however, the abnormality was restricted to the gonads, the adnexae being female on both sides.

Aetiology.—The case mentioned above in which one-half of the body was male and the other female is analogous to the condition of gynandromorphism found in lower animals, such as insects and birds. Crew (1927) has shown that gynandromorphism in insects is due to loss of sex chromatin at the time of the first division of the fertilized ovum. It is well known that anomalies are not uncommon at this stage of development, such as the complete or partial separation of the primary cells which is thought to give rise to identical or conjoined twins. Since genetic maleness appears to depend on the absence of a second X chromosome rather than on the presence of the Y chromosome, the loss of one sex chromosome during the first division of the female zygote would result in one male and one female cell. It is possible that some such mechanism may operate in man. In the higher animals, however, sex may not be absolutely predetermined by the sex chromosomes, and the genetic direction of development may be overridden by other factors, of which hormones appear to be the most important. The relative significance of genetic, hormonal, and environmental factors in producing the many different varieties of human hermaphroditism is as yet unknown.

Management.—The major problem in the management of these cases is to decide on the predominant sex. In patients first diagnosed in adult life it is usually found that the emotions are appropriate to the sex of upbringing (the "social" sex) and that removal of the inappropriate gonad and suitable hormone therapy to reinforce the remaining gonadal tissue is satisfactory. In children, however, it is difficult to predict the direction the emotions will take at puberty. This is well shown in the case reported by Lattimer *et al.* (1943) and Engle *et al.* (1946). This patient had been brought up as a girl, but was masculine in behaviour. At the age of 11 an ovotestis and an ovary were removed, leaving a third gonad, which was thought to be a testis. The patient was then regarded as a boy, but at puberty three years later a strong female trend became apparent. The remaining gonad was removed (and proved to be an ovotestis) and the social sex changed back to female.

In Case 1 above, the parents were advised to bring up the child as a boy in view of the testis being much the larger gonad and the large size of the phallus, and in spite of the presence of female adnexae. This advice was reinforced by Professor Barr's opinion on the somatic nuclei. In consequence of this decision the hypospadias will be corrected at a later date.

The third patient mentioned above is psychologically a normal male, although the epidermal nuclear picture is that of a female. It must be remembered that the patient has been treated with testosterone and also emotionally conditioned to some extent by his environment. Nevertheless it would appear that considerable caution is required when assessing the results of skin biopsy in true hermaphrodites. It is of interest in this connexion that 16 cases of true hermaphroditism have been social females and 41 social males (plus 2 neonatal cases), whereas skin biopsy on five true hermaphrodites has shown four with a female nuclear picture and only one with a male picture—Case 1 described above (M. L. Barr, 1954, personal communication).

Cases in the Literature

True hermaphroditism is probably less rare than the number of case reports might suggest. Several cases—for example, Photakis (1916) and Moriarty (1944)—were discovered by chance at post-mortem examination. Young (1937) collected 20 cases from the literature, to which McIver *et al.* (1944) added 11 others and two of their own. Greene *et al.* (1952) collected 48 cases and added one, and Fischer *et al.* (1952) collected 49 cases and added one. Two cases in the latter's list were, however, rejected by Greene (Stirling (1946)—"microscopy of testis unconvincing"; and Malgras and Gricouroff (1939) as ovarian tissue was deduced indirectly and not demonstrated histologically). Excluding these two cases, we are left with a net total of 53 cases. To these must be added six further cases: Broster *et al.* (1938), Berardinelli (1952), Brewer *et al.* (1952), Baker *et al.* (1953), Delore *et al.* (1953), and Laycock and Davies (1953). With the new cases reported here, the total number of acceptable cases is therefore 61.

Summary

Two cases of true lateral hermaphroditism are reported and the further progress of a third (bilateral) case is noted.

The first case presents the clinical problem of predicting the predominant psychological sex in adult life when true hermaphroditism is diagnosed in infancy. In this case the chromosomal sex, as determined by the method of Moore *et al.* (1953), was male, which agreed with the preponderance of male gonadal tissue. On the other hand, the third case cited has grown up a psychologically normal male, but the chromosomal sex is female. This suggests that the chromosomal determination of sex is of limited value in these cases and must

be considered in conjunction with all the available clinical evidence in deciding on the sex of upbringing.

The diagnosis of true hermaphroditism rests finally on the histology of the gonads, but a tentative diagnosis may be made on clinical grounds.

Fifty-nine previous cases of true hermaphroditism have been found in the literature. This small number is probably not a true indication of the real incidence of the condition.

I am indebted to Mr. L. R. Broster for permission to publish these cases under his care, and for his encouragement and help in the preparation of this paper. I have also to thank Dr. W. E. Evans and Professor M. L. Barr for reports on the sections.

REFERENCES

- Baker, W. J., Callahan, D. H., and Ragius, A. B. (1953). *J. Urol. (Baltimore)*, 69, 439.
 Berardinelli, W. (1952). *Acta endocr. (Kbh.)*, 9, 297.
 Brachetto-Brian, D., Grimaldi, F. E., and Costa, A. O. T. (1943). *Rev. Asoc. méd. argent.*, 57, 900.
 Brewer, J. I., Jones, H. O., and Culver, H. (1952). *J. Amer. med. Ass.*, 148, 431.
 Broster, L. R., Allen, C., Vines, H. W. C., Patterson, J., Greenwood, A. W., Marrian, G. F., and Butler, G. C. (1938). *The Adrenal Cortex and Intersexuality*, p. 42. Chapman and Hall, London.
 Crew, F. A. E. (1927). *The Genetics of Sexuality in Animals*, p. 26. Cambridge Univ. Press, Cambridge.
 Davis, D. M., and Scheffey, L. C. (1946). *J. Urol. (Baltimore)*, 56, 715.
 Delore, P., Noel, R., Perrin, J., Guinet, P., and Blanc, J. (1953). *Pédiatrie*, 8, 458.
 Engle, E. T., Lattimer, J. K., and Yeaw, R. C. (1946). *J. Urol. (Baltimore)*, 56, 731.
 Fischer, H. W., Lischer, C. E., and Byars, L. T. (1952). *Ann. Surg.*, 136, 864.
 Greene, R., Matthews, D., Hughesden, P. E., and Howard, A. (1952). *Brit. J. Surg.*, 40, 263.
 Hinman, F. (1935). *Principles and Practice of Urology*, p. 134. Saunders Philadelphia.
 Lattimer, J. K., Engle, E. T., and Yeaw, R. C. (1943). *J. Urol. (Baltimore)*, 50, 481.
 Laycock, H. T., and Davies, D. V. (1953). *Brit. J. Surg.*, 41, 79.
 McIver, R. B., Seabaugh, D. R., and Mangels, M. (1944). *J. Urol. (Baltimore)*, 52, 67.
 Malgras and Gricouroff (1939). *Mém. Acad. Chir. (Paris)*, 65, 272.
 Moore, K. L., Graham, M. A., and Barr, M. L. (1953). *Surg. Gynec. Obstet.*, 96, 641.
 Moriarty, J. D. (1944). *Amer. J. Path.*, 20, 799.
 Photakis, B. (1916). *Virchows Arch. path. Anat.*, 221, 107.
 Stirling, W. C. (1946). *J. Urol. (Baltimore)*, 56, 720.
 Young, H. H. (1937). *Genital Abnormalities, Hermaphroditism and Related Adrenal Diseases*, p. 159. Baillière, Tindall and Cox, London.

Medical Memoranda

Hair-grips as Foreign Bodies

The accidental swallowing of metallic foreign bodies by children is of everyday occurrence. The great majority pass through the gastro-intestinal tract without incident, being voided per anum in a few days, and, considering their size and shape, it is indeed remarkable how some manage to get through in tiny children without causing harm.

I would here like to draw attention to one variety of metallic foreign body commonly swallowed by young children—the familiar hair-grip. Hair-grips do not always pass through the gastro-intestinal tract with the same ease as the many other types of foreign body; sometimes they remain in the stomach or become impacted in the second part of the duodenum. This is particularly the case in very young children.

During the past ten years I have treated seven children with swallowed hair-grips and have had to resort to operation for the removal of the grip on four occasions. The four cases which came to operation comprised two girls and one boy, all under the age of 4 years, and one girl aged 7 years. The other three, all girls, were aged 4½, 6½, and 13 years, and in these the grip was passed normally in a short space of time. Fig. 1 is a photograph of the four sizes of grip in common use, and it is the third largest size which is usually swallowed.

In children, particularly those under the age of 4 years, the grip tends to remain in the stomach, or, if it passes the

pylorus, may become impacted vertically across the second part of the duodenum. Fig. 2 is an x-ray film showing the grip in such a position.

In two of the four cases which came to operation the grip was impacted across the duodenum, one child having swallowed the grip 17 days and the other 14 days before admission. The other two failed to pass the pylorus, one having been in the stomach five days and the other ten days before operation.

It is a simple matter to remove grips from the stomach, but nothing like so easy from the second part of the duodenum. In one of my two cases the grip was coaxed back into the stomach, and in the other it was coaxed onwards into the jejunum before removal, both being difficult manoeuvres. However, I feel it is better to coax the grip

FIG. 1.—Photograph (actual size) of the four hair-grips in common use. The third largest is the one usually swallowed.

FIG. 2.—X-ray photograph of a grip impacted vertically in the second part of the duodenum in a girl aged 7 years. The gas-filled shadows of the stomach and colon can be clearly seen.

into an accessible region for removal than to open up or attempt to mobilize the second part of the duodenum in a young infant.

Prevention is better than cure, and mothers should be warned not to use grips for dressing young children's hair, but instead to employ the old-fashioned hair-slide with a large ornamental ring attached. It is the practice of the nursing staff of the children's surgical ward at the Royal Bucks Hospital, Aylesbury, to remove all hair-grips from young children on admission.

In those who have swallowed a hair-grip, treatment with a bulky diet and "normacol" is at once started, and x-ray films are taken of the whole abdomen at forty-eight-hour intervals to check progress. All stools are examined for the passage of the foreign body. In many cases emetics have been administered prior to admission to hospital in the