- Reiss, M., Badrick, F. E., Halkerston, I. D. K., and Plaice, C. (1951). *Nature (Lond.)*, **168**, 206.
- Renold, A. E., Jenkins, D., Forsham, P. H., and Thorn, G. W. (1952). *J. clin. Endocr.*, **12**, 763.
- Samuels, L. T., Evans, G. T., and McKelvey, J. L. (1943). *Endocrinology*, **32**, 422.
- Sandberg, A. A., Nelson, D. H., Palmer, J. G., Samuels, L. T., and Tyler, F. H. (1953). *J. clin. Endocr.*, **13**, 629.
- Sayers, G. (1950). *Physiol. Rev.*, **30**, 241.
- , Burns, T. W., Tyler, F. H., Jager, B. V., Schwartz, T. B., Smith, E. L., Samuels, L. T., and Davenport, H. W. (1949). *J. clin. Endocr.*, **9**, 593.
- , and Sayers, M. A. (1948). *Recent Progr. Hormone Res.*, **2**, 81.
- Selye, H. (1946). *J. clin. Endocr.*, **6**, 117.
- Simpson, S. A., Tait, J. F., Wettstein, A., Neher, R., Euw, J. v., Schindler, O., and Reichsteia, T. (1954). *Experientia (Basel)*, **10**, 132.
- Stack-Dunne, M. P. (1953). *Ciba Foundation Colloquia on Endocrinology*, **5**, 133.
- Talbot, N. B., Butler, A. M., Berman, R. A., Rodriguez, P. M., and MacLachlan, E. A. (1943). *Amer. J. Dis. Child.*, **65**, 364.
- Tarantino, C. (1951). *Folia Endocr. (Pisa)*, **4**, 197.
- Thorn, G. W. (1954). Medical Research Society Lecture.
- , Forsham, P. H., Prunty, F. T. G., and Hills, A. G. (1948). *J. Amer. med. Ass.*, **137**, 1005.
- , Jenkins, D., and Laidlaw, J. C. (1953). *Recent Progr. Hormone Res.*, **8**, 171.
- Tobian, L. (1949). *J. clin. Endocr.*, **9**, 319.
- Tyler, F. H., Migeon, C., Florentin, A. A., and Samuels, L. T. (1954). *Ibid.*, **14**, 774.
- Van Cauwenberge, H., and Betz, H. (1952). *Lancet*, **1**, 1083.
- , and Heughebaert, C. (1951). *Ibid.*, **1**, 771.
- Venning, E. H. (1946). *Endocrinology*, **39**, 203.
- , Randall, J. P., and Gyorgy, P. (1949). *Ibid.*, **45**, 430.
- Wilkins, L., Lewis, R. A., Klein, R., Gardner, L. I., Crigler, J. F., Rosenberg, E., and Migeon, C. J. (1951). *J. clin. Endocr.*, **11**, 1.
- , Gardner, L. I., Crigler, J. F., Silverman, S. H., and Migeon, C. J. (1952). *Ibid.*, **12**, 257.
- Wilson, H., Glyn, J., Scull, E., McEwen, C., and Ziff, M. (1953). *Proc. Soc. ex Biol. (N.Y.)*, **83**, 648.
- Zaffaroni, A. (1950). *J. Amer. chem. Soc.*, **72**, 3828.
- (1953). *Recent Progr. Hormone Res.*, **8**, 51.

ENDOCRINE TREATMENT IN PSYCHIATRY*

BY

R. E. HEMPHILL, M.D., D.P.M.

Lecturer in Charge of Department of Psychiatry, University
of Bristol; Medical Superintendent, Bristol Mental
Hospitals Group

Endocrine therapy has a definite but restricted place in psychiatric practice; in some conditions its use is accepted, in others presumptive. With a few possible exceptions, such as cretinism, there are no psychiatric disorders in which an endocrine disturbance or deficiency has been proved to be an exclusive or major factor in causation. In contrast to the many papers published on research in the endocrinology of psychiatric illness in recent years, there are few about endocrine therapy and fewer that carry conviction.

Since the ductless glands are essential to adaptation and the preservation of physical and emotional stability, and can be disturbed by psychological upset, it is reasonable that psychiatrists should have looked to the field of endocrine treatment.

Although physiological disturbances and abnormalities of body constitution can be seen in any selection of chronic psychotics, the usual symptomatology of endocrine illnesses is rare in psychiatric disorders. Laboratory methods for investigating ductless gland function which have been evolved up to the present are probably not sufficiently precise or reliable to indicate and guide therapy in the absence of clinical signs of endocrine disease, and are not yet convenient for routine use. Thus endocrine treatment is still empirical, and the attempts to make it rational by laboratory methods have not yet been successful.

Endocrine preparations may be used because of their physiological and pharmacological effect or as replacement; probably to some extent both aspects enter into successful therapy. It is improbable that lack of one hormone is responsible for any psychiatric disorder, and a replacement therapy is unlikely to be found.

An endocrine disturbance is only one factor in an illness which can be regarded as a reaction between physical and psychological components; but it may be responsible for great difficulties in adjustment. Endocrine therapy aims at improving this factor, so contributing to the total therapeutic effort by benefiting the adaptive or reactive capacity of the patient. It must be preceded by careful psychiatric assessment and investigation, and accompanied by adequate psychotherapy.

Endocrinology in psychiatry can conveniently be considered from the following aspects: (1) conditions encountered in psychiatric practice for which therapy is no longer experimental; (2) psychiatric aspects of endocrine disorders; and (3) experimental endocrine treatment of the psychoses and psychoneuroses.

Conditions Encountered

The Menopause

The features of menopausal depression, irritability, or neurosis need only be mentioned briefly here. There are few women who do not experience psychological changes during the months or years in which ovarian function is failing. The degree to which a woman is disturbed seems to depend upon her capacity for emotional adjustment, general stability, and reaction to ill-health. Many women who suffer most at the menopause report that they have usually experienced mood swings and physical discomfort at some part of the menstrual cycle. It is not correct to assume that a certain kind of personality suffers at the menopause, or is immune; but unbalanced and especially aggressive females often become unreasonable almost to the point of psychosis both at the menstrual cycle and at the menopause. A gradual menopause produces more discomfort than an abrupt cessation. Vasomotor disturbances and hot flushes correspond closely with mood swings. It is believed that these are due to a dysbalance in the relationship between pituitary-stimulating hormones and the peripheral ovarian hormones that are now failing, which has been suggested by high prolactin excretion.

Treatment with synthetic or natural oestrones (stilboestrol, dienoestrol, or oestrone), or, in severe cases, oestroform by injection, in sufficient quantity to control hot flushes, is all that is required for the somatic disturbances, and makes psychotherapy easier. Stilboestrol is toxic to about 20% of women, especially the older; the dose is about 0.1 mg. daily. The usual dose of oestrone is from 0.2 to 1 mg. daily. A severe menopausal disturbance sometimes occurs years after artificial menopause by radium, or after hysterectomy if the ovaries remain; this is to be treated on the same lines as the normal menopause and is mentioned for this reason. Convulsive therapy—so effective in involutional melancholia—often aggravates menopausal depression and is contraindicated in the presence of hot flushes, because of its stimulating effect on anterior pituitary lobe function.

Involutional Melancholia

Unlike the menopausal reaction, this is a true psychotic depression, with ideas of guilt or of incurable illness, much agitation, and a marked tendency to suicide. It rarely follows a gradual menopause with vasomotor disturbances, and it usually occurs a year or more after sudden cessation of menses. The pre-psychotic personality is conscientious and inflexible, in contrast to the lability of the menopausal reaction. There is often an appearance of rapid physical ageing, and in some cases there seems to be an anterior pituitary insufficiency, which may be benefited by treatment

*From a Maudsley Bequest Lecture given in London on April 5, 1954.

with A.C.T.H. But, normally, E.C.T. is the treatment of choice. Experience suggests that oestrones are ineffective, or even contraindicated, as they may suppress anterior pituitary activity.

Pre-menstrual Tension

This term indicates a state of mental and physical tension occurring towards the latter half of the menstrual cycle and prominent about a week before menstruation. The current view is that it is due to water retention produced by abnormal elevation of the oestradiol/progesterone ratio (Rees, 1953; Malleon, 1953).

The diagnosis is made on the occurrence of irritability, emotionalism (even unreasonable aggression), mental discomfort, lack of concentration, coupled with physical discomfort, such as a sense of being blown out, a gain in weight, discomfort in the fit of clothes—due presumably to localized adiposity or water retention. There are occasionally attacks of migraine, or of epilepsy in epileptic subjects.

The usual treatment is ammonium nitrate or ammonium chloride in coated tablets, 0.5 to 1 g. thrice daily for 10 to 12 days before the onset of menses. If this is unsuccessful, progesterone injections, 25 mg. twice daily, or ethisterone tablets, 5 to 20 mg. orally, daily or twice daily. Methyltestosterone, 10 mg. daily, is said to be effective, but may cause virilization.

Assuming that the condition is due to endocrine dysbalance, normal regulation may be established in a few months, when treatment can be discontinued.

Abnormal Libido

Impotence.—This is the inability to perform or complete the sexual act in a male. It may be caused by conditions such as hypogonadism and intersexuality; but usually the external genitalia are normal and the impotence is of psychological origin. This is rarely benefited by endocrine therapy, except where it accompanies the male climacteric or depression, or where it is intermittent and has not existed for long. In long-standing cases of psychological impotence, the interest has been displaced from the sexual object to anxiety whether the act can be performed, causing extinction of the sexual stimulus. Where impotence has occurred in a setting of exhaustion or business worries and has not been prolonged, and where there has not been much controversy between husband and wife, testosterone may improve the sense of well-being, increase libido, and, by making one or two performances possible, improve self-confidence so that anxiety disappears. It is best administered by injection—25 mg. of testosterone propionate three times weekly for a month; oral administration of methyltestosterone is insufficient. Treatment should be discontinued after a fortnight if there has been no response. The patient should not be led to believe it will produce a cure, lest it should not be successful. Intensive psychotherapy is essential. It is unwise to attempt treatment with testosterone in long-standing cases, because of the worsening effects produced by disappointment, as it will certainly fail.

Frigidity.—The diagnosis of frigidity is obviously difficult to make, and the term is a relative one. However, loss of sexual desire or enjoyment is not uncommonly complained of by married women, especially after childbirth; and it has been claimed that treatment with small doses of either oestrone or testosterone improves the appreciation of physical sexual stimuli and will therefore reduce temporary frigidity. Testosterone may cause virilization, and for this reason some authors prefer oestrone. Inunction of external genitalia with oestrone or testosterone cream has been advocated. True frigidity of long standing—that is, inability to experience or elaborate psychically sexual stimuli—is a rare condition, and probably cannot be improved by hormones.

Homosexuality and Excessive Sexuality.—The direction of homosexual or heterosexual drives cannot be altered with sex hormones; but the force of sexual drive in males can be reduced by treatment with female sex hormones. There are a number of papers on the subject, and all the

authors are in general agreement about the indications and the rationale. Treatment with female sex hormones reduces the production of gonadotrophic hormones, producing a secondary effect on testicular function and secretion. The reduction of sex drive is comparable to that achieved by surgical castration, although some authors claim that endocrine treatment is more effective (Hamilton, 1943). Stilboestrol is the usual preparation, administered in increasing doses until the nipples become pigmented and the breasts sore; an alternative is ethynol oestradiol, 0.5 or 1 mg. daily. Eventually almost complete testicular atrophy will be produced, but not necessarily a total suppression of the abnormal sex drives. It is interesting that notice has been taken of this treatment at the recent trial of an extreme and persistent homosexual offender, in whom abnormal behaviour had continued in spite of the production of almost complete atrophy of the testes by stilboestrol. The judge took into consideration that the offender had therefore done everything possible that medical treatment could offer, although in his case it had not been completely successful. Persistent homosexual offenders should therefore be encouraged to persevere with this treatment, although in some cases a satisfactory control of sexual behaviour may not necessarily be achieved.

Male Climacteric

The gradual diminution in the production of male sex hormones in middle and later age is held to be responsible for loss of physical and mental powers and libido. The male climacteric as an entity has been well discussed by Spence (1954). The diagnosis of male climacteric reaction cannot often be made with confidence, because arteriosclerosis or other systemic diseases complicate the picture and there is no dramatic event like the cessation of menses to indicate endocrine change. Moodiness, irritability, tearfulness, inability to concentrate, paraesthesiae, and sometimes hot flushes, suggest the diagnosis. Treatment is with testosterone by injection, 25 mg. daily or thrice weekly until there is improvement, methyltestosterone 5 to 10 mg. three times a day being given as continuation therapy for a year or more. There is usually improvement in mental and physical activity, sexual performance, and altered sense of well-being. Better peripheral circulation and flushing of the face are favourable early signs.

Cretinism

A causal association between thyroid deficiency and failure of intellectual and physical development, known as cretinism, was observed early in the history of modern medicine. Some authors make a distinction between juvenile myxoedema and cretinism. The cretin suffers from athyreosis from intrauterine life or early infancy (Means, 1948). The clinical picture is well known, but the results of treatment are less certain, for good results cannot be obtained in infantile myxoedema unless the brain is free from congenital abnormality.

At one time it was believed that only early and cautious treatment was likely to be successful. Surveys by Lewis *et al.* (1937), and by Topper (1951), who gives a good bibliography, indicated that promptness and continuity of thyroid treatment did not altogether decide whether a child remained backward or not. Topper pointed out that, according to electroencephalographic studies, cerebral defect explains much of the mental deficiency of the treated cretin, and that when such a defect is present in infancy no amount of thyroid can improve the brain. In her large series, 39% remained defective in spite of treatment.

Treatment should be pushed to the point of tolerance, which can be found by trial and error. The small doses usually recommended in cretinism are based on requirements necessary for alteration of hair and skin, and do not produce acceleration of growth and intellectual development. While tolerance is being determined the patient should be kept in hospital. Means and Lerman (1935) pointed out that with a constant daily dose the maximum

effect is not produced for about three weeks, and it is not safe to increase the dose oftener than every three to four weeks. Toxic symptoms are rapid pulse, a small rise in temperature initial loss of weight, increased nervousness, and frequent bowel movements; and some authors believe that these occur at the limit of safety and can be tolerated for a while. Dangerous symptoms due to excessive dosage are extreme restlessness, twitching, diarrhoea, vomiting, continued loss of weight, very rapid pulse, high fever, and syncope. As much as 4 gr. (260 mg.) daily is given at 21 months without adverse effect, but average doses are: 6 to 18 months, $\frac{1}{2}$ to 2 $\frac{1}{2}$ gr. (30 to 160 mg.); 1 $\frac{1}{2}$ to 8 years, 1 to 2 gr. (65 to 130 mg.); 8 to adolescence, 1 $\frac{1}{2}$ to 3 gr. (100 to 200 mg.). It is worth noting that the dose tolerated is not proportionate to the age and weight, for a young cretin can tolerate as much as a 12-year-old child with hypothyroidism (Brown, 1951).

Psychiatric Disturbances in Endocrine Disorders

Although psychological disturbances are a feature of many endocrine disorders they rarely amount to recognized psychiatric syndromes. Classical psychoses are uncommon even in thyrotoxicosis. Treatment of some endocrine conditions may precipitate a psychotic episode.

Thyrotoxicosis often follows an acute emotional upset, and it has been contended that it occurs usually in frustrated and emotionally dependent individuals who wish to take care of others. Ham *et al.* (1951) have agreed with some others that symptoms of thyrotoxicosis can be understood as the patient's response to early experiences and that although the experiences differ in detail they fall into a pattern characteristic of the disease. In all the patients they studied there were extreme feelings of insecurity in early life, a strong sense of responsibility, but a tendency to suffer in silence; this is in some contrast to the anxious patient, who has no such tendency. From the psychosomatic point of view, some cases of thyrotoxicosis may be an end-result of an endocrine disturbance operating through emotional pathways. Thyroidectomy may be followed by a degree of myxoedema, producing the usual sluggishness. Where there has been a coexisting anxiety state, or the diagnosis should have been an anxiety neurosis and not thyrotoxicosis, sometimes the thyroid requirements of body and brain are not met. The patient will be less able to cope with the anxiety situation than before operation, and the picture will then be continuing anxiety with agitation or even restless delirium. Treatment with thyroid hormone is indicated.

Myxoedema usually produces apathy and sluggishness. The administration of thyroid hormone may activate a paranoid psychosis, as if, by increasing metabolism, an uneasy psychological balance had been disturbed. Withdrawal of thyroid is not necessarily followed by an improvement in the psychiatric state, and cases of chronic paranoid psychosis arising in this way are occasionally seen. Caution is necessary in treating myxoedema in patients of paranoid constitution. Myxoedema may be associated with confusion and depression, especially in later life (Asher, 1949). A picture of organic brain change, with clouding of consciousness, delirium, or agitation, may be due to myxoedema or its cardiac complications (Calvert *et al.*, 1954). In such states of hyperactivity myxoedema may be overlooked unless remembered. Many of these patients improve with thyroid hormone; but care is needed, for an increased metabolism in the aged or in organic brain disorder may produce accentuation of confusion, delirium, and death.

Addison's Disease.—In adrenal hypofunction the capacity for adaptation is reduced. Cleghorn (1953) has compared the psychological changes in Addison's disease with myxoedema, and agrees with Engel and Margolin (1942) that the psychiatric symptoms of apathy, negativism, depression, and irritability occur in more than 50% of cases; he also quotes Stoll (1953), who reported similar disturbances in 35% in the pre-Addisonian state and 66% in the developed con-

dition; according to Stoll, they all improved remarkably under treatment with cortisone.

Adrenogenital Syndrome.—Virilism in females due to adrenocortical hyperplasia or cortical tumours has been the subject of much study. Broster *et al.* (1938) and others have shown that in extreme cases a combination of aggressiveness and paranoid thinking are to be expected. It has been claimed that after operation a more normal adrenal function is accompanied by reduction in psychopathic elements. These are extreme conditions, and operation seems to be very seldom indicated. Nevertheless, this work suggests that further research in the relationship between hypercortical adrenalism and aggressive paranoid behaviour might be worth while.

Experimental Endocrine Treatment

Endocrine treatment in the psychoses is still in the experimental stage, since up to the present no conclusive evidence has been produced about its rationale and value. In spite of the views expressed by some workers there is nothing to suggest that a new system of psychiatric diagnosis or classification can be derived from endocrine investigations, and classical concepts cannot be disregarded, nor can laboratory methods replace clinical observation.

Most analytic methods so far evolved lack precision, and it is not certain what are the range and the significance of normal values in non-psychotic individuals, how much variation within the range normally occurs, and how the values are affected by age, sex, physical illness, diet, and state of nutrition.

Therapy directed on the basis of figures alone is notoriously unsatisfactory in general medicine, and is even more open to error in psychiatry. In a series of 150 recent and chronic psychotics, with some psychoneurotics, treated on laboratory indications at the Bristol Mental Hospitals, only 12 showed a significant improvement considered to be due to the hormone therapy. Psychiatric recovery or deterioration does not necessarily correspond with laboratory findings.

Nevertheless, in spite of the limitations of the methods, and the impossibility of defining scientifically the dimensions of psychiatric illnesses and degrees of improvement or deterioration, there is a good deal of evidence to justify research in the field of experimental endocrinology, with a hope of eventual therapy in some conditions.

Gjessing (1938) has shown that in the rare condition called periodic catatonia, alterations in nitrogen balance coincided with the catatonic episodes, and the attacks could be terminated or aborted by treatment with thyroxine and dried thyroid to discharge nitrogen stored. The value of Gjessing's treatment has been confirmed by other workers.

Many psychotics, especially schizophrenics, do not react normally to hormones. It has been shown that schizophrenics are, or soon become, insensitive to thyroid; and insulin resistance in schizophrenia is well known. The same seems to be true for cortisone and A.C.T.H.: neither depressives nor schizophrenics usually exhibit the euphoria and emotional changes described in normals. Hench *et al.* (1950) and others believed that the familiar euphoria was not due to relief of pain, but a positive factor accompanied by increased mental function. Some workers have assumed that psychological changes were most marked where there was some evidence of previous mental illness or psychological maladjustment; this is certainly not the case. Lewis and Fleminger (1954) have shown that of 12 patients with a history of recent mental illness who had been treated with cortisone or A.C.T.H. none developed a severe mental illness. In the Bristol Mental Hospitals, A.C.T.H. has been used extensively, and reactions comparable to those described in normals have never been seen. This suggests that in some psychiatric disorders the ductless gland system is abnormal; it may be insensitive or react in an abnormal way, or metabolic processes may destroy or neutralize the action of hormones.

A great amount of research in the endocrinology of schizophrenia, especially of the adrenal cortex, has been carried out at various centres. Evidence of failure to preserve adequate endocrine equilibrium and of abnormal adrenal responsivity has been advanced, but the treatment of schizophrenia with A.C.T.H. on the assumption of evidence of lack of adrenal stimulation, attempted by Glaser and Hoch (1951), by the Worcester group (Hope *et al.*, 1952), by Kobbernagel *et al.* (1953) in Scandinavia, and at the Bristol Mental Hospitals and elsewhere, has been practically unsuccessful.

Cortisone has been used, on the whole without much success, but Cohn *et al.* (1953) have reported that 6 out of 21 chronic patients who had been ill for about six years were discharged from hospital. Their patients were treated with between 300 and 800 mg. daily for four weeks. Toxic effects were observed in four at least, with oedema, retention of urine, and renal colic. It is possible that those who improved did so because of the destructive action of cortisone on the hypothalamus.

Castor *et al.* (1951) have shown that A.C.T.H. in rats produced degeneration of the paraventricular nuclei of the thalamus; cortisone did the same, but it also produced changes in the thalamic and hypothalamic nerve cells. These areas are usually affected by prefrontal leucotomy, and it is probable that Cohn's success was due not to the physiological but to the destructive effect of the cortisone.

Treatment with thyroid has been recommended in various conditions. Danziger and Kindwall (1953) have reported success in a number of cases of schizophrenia in which failure to remain in remission after other forms of treatment, or periodic fluctuations, seem to be common features. Their material may include cases of Gjessing's type. It is therefore worth while administering thyroid to the point of toxicity empirically in relapsing schizophrenia after E.C.T. and insulin-coma therapy have been unsuccessful. Thyroid is valuable in reducing apathy in schizophrenics after insulin coma (Mayer-Gross *et al.*, 1954).

Yet the role of the thyroid in psychiatric illness is still quite uncertain. Reiss (1954), using a radioactive technique, has reported that the thyroid activity was within the limits he regarded as normal in about 70% of psychiatric admissions. In a selected series of 450 cases, after treatment by various methods, including hormones, E.C.T. and insulin coma, the thyroid activity was reported as "normal" in 72%. This suggests that the thyroid state is of consequence in only a very small percentage of psychiatric cases, if at all, irrespective of whether the thyroid activity is regarded as causally related to the illness or indirectly as an indicator of the state of the endocrine equilibrium.

In manic-depressive psychosis A.C.T.H. has been used as an alternative to E.C.T., on the assumption that E.C.T. exercises its favourable effect by the mobilization of anterior pituitary hormones. But, although in a very few cases remissions have been seen after a single dose of 25 mg. of A.C.T.H., this hormone is practically valueless in depression and without effect in mania, as Cleghorn *et al.* (1950) and the Worcester group have reported.

Sands (1954) recommends dehydroisoandrosterone ("diandrone") for young inadequate psychopaths with timidity, physical immaturity, little energy, and perhaps homosexuality; to combat the residual disabilities after insulin coma therapy in schizophrenia; in stammering, on a basis of inadequacy; and to assist psychotherapy in some cases of neurosis. Strauss *et al.* (1952) originally gave three criteria: abnormal 17-ketosteroid excretion, a mild or slowly deteriorating mental illness, and the appearance of physical immaturity. They regarded the normal ketosteroids as 6 to 21 mg. per 24 hours, and the beta fraction 5 to 15%. Diandrone is contraindicated where there is evidence of aggressive traits. Dosage is 5 to 30 mg. orally. Overdosage is suggested by undue irritability and overactivity.

Summary

There is a place for endocrine treatment in psychiatry, and the psychiatrist is often confronted with complications of the menses, the menopause, disturbance of libido, homosexuality, male climacteric, and cretinism. Treatments for these are reviewed.

In the field of the psychoses there is some evidence of endocrine disturbances, the exact nature of which is not yet clear. Random cures of psychotic conditions are always to be expected, but they contribute little to the understanding of the physiology of mental disorders.

Assuming that functional derangements of endocrine balance occur spontaneously and are responsible for psychotic illnesses, spontaneous remissions are to be expected, which must make one extremely cautious in attributing recovery to a particular therapy. It is probable that there is a time lag between alteration in physiological state and the appearance of mental change, which would make a reliable correlation of physiological and clinical observations difficult.

REFERENCES

- Asher, R. (1949). *British Medical Journal*, 2, 555.
 Broster, L. R., Allen, C., Vines, H. W. C., Patterson, J., Greenwood, A. W., Marrian, C. F., and Butler, G. C. (1938). *The Adrenal Cortex and Intersexuality*. Chapman and Hall, London.
 Brown, E. E. (1951). *Northw. Med. (Seattle)*, 50, 22.
 Calvert, R. J., Smith, E., and Andrews, L. G. (1954). *British Medical Journal*, 2, 891.
 Castor, C. W., Baker, B. L., Ingle, D. J., and Li, C. H. (1951). *Proc. Soc. exp. Biol. (N.Y.)*, 76, 353.
 Cleghorn, R. A. (1953). *J. clin. Endocr.*, 13, 1291.
 —, Graham, B. F., Saffran, M., and Cameron, D. E. (1950). *Canad. med. Ass. J.*, 63, 329.
 Cohn, J. B., Steckler, G. A., and Rubinstein, J. (1953). *J. clin. exp. Psychopath.*, 14, 161.
 Danziger, L., and Kindwall, J. A. (1953). *Dis. nerv. Syst.*, 14, 3.
 Engel, G. L., and Margolin, S. G. (1942). *Arch. intern. Med.*, 70, 236.
 Gjessing, R. (1938). *J. ment. Sci.*, 84, 608.
 Glaser, G. H., and Hoch, P. H. (1951). *Arch. Neurol. Psychiat. (Chicago)*, 66, 697.
 Ham, G. C., Alexander, F., and Carmichael, H. T. (1951). *Psychosom. Med.*, 13, 18.
 Hamilton, J. B. (1943). *Proc. Soc. exp. Biol. (N.Y.)*, 54, 309.
 Hench, P. S., Kendall, E. C., Slocumb, C. H., and Polley, H. F. (1950). *Arch. intern. Med.*, 85, 545.
 Hope, J. M., Elmadjian, F., and Malamud, W. (1952). *J. clin. exp. Psychopath.*, 12, 267.
 Kobbernagel, F., *et al.* (1953). *Acta endocr. (Kbh.)*, 13, 162.
 Lewis, A., and Fleminger, J. J. (1954). *Lancet*, 1, 383.
 —, Samuel, N., and Galloway, J. (1937). *Ibid.*, 2, 5.
 Maleson, J. (1953). *Ibid.*, 2, 158.
 Mayer-Gross, W., Slater, E., and Roth, M. (1954). *Clinical Psychiatry*, p. 288. Cassell, London.
 Means, J. H. (1948). *The Thyroid and its Diseases*, 2nd ed., p. 274. Lippincott, Philadelphia.
 —, and Lerman, J. (1935). *Arch. intern. Med.*, 55, 1.
 Rees, L. (1953). *J. ment. Sci.*, 99, 62.
 Reiss, M. (1954). *Ibid.*, 100, 687.
 Sands, D. E. (1954). *Med. ill. (Lond.)*, 8, 311.
 Spence, A. W. (1954). *British Medical Journal*, 1, 1333.
 Stoll, W. A. (1953). *Die Psychiatrie des Morbus Addison*. Thieme, Stuttgart.
 Strauss, E. B., Sands, D. E., Robinson, A. M., Tindall, W. J., and Stevenson, W. A. H. (1952). *British Medical Journal*, 2, 64.
 Topper, A. (1951). *Amer. J. Dis. Child.*, 81, 233.

"For doctors in Nyasaland the mango season means a plague of 'mango fractures.' Every year about this time the hospitals, especially in Zomba, the capital, are crowded with piccaninnies with broken limbs, plaster-of-paris hallmarks of an over-ambitious pursuit of the delectable fruit. Recently in Zomba there were eight small African boys with fractured legs—all victims of the mango craze, and all very sorry for themselves. But the little devils will never learn so long as their elders fail to set a good example of careful climbing and cautious picking: in a village not far from Zomba two Africans, grandfathers both, broke their arms when falling from mango trees, which often reach a height of sixty and seventy feet. Mangoes, incidentally, which were probably introduced into the Protectorate by Arab slave traders, are the most widely cultivated tropical fruit. They say Buddha often rested in the shade of mango trees. He would be unwise to take the risk in Nyasaland."—*Manchester Guardian*.