from cloudy swelling to necrosis. *Medulla*: A similar degeneration of the straight tubules, many of which contained blood casts. The vessels showed some thickening, chiefly of the muscle coats.

Liver.—2,000 g. Flabby, yellow, and fatty. The cells were irregular in size and shape and were separated by spaces. No cirrhosis or congestion was seen. In the portal areas the vessels and ducts appeared thickened, with loose fibrous tissue rather irregularly disposed. Some lymphocytic infiltration was also present in these areas.

Pathological Data

Date	Throat Swab	Urine	Blood Urea	Serum Creatine	Serum Creatinine	Serum Bilirubin
6/4/54	Predominant growth <i>Str. viridans</i>	Acid. Creatine ++. Many white and red cells. No casts				
9/4/54		Creatine, 840 mg./24 hrs. Creatinine, 1,100 mg./24 hrs.	177 mg./100 ml.	8 mg./100 ml.	4.5 mg./100 ml.	
12/4/54			215 mg./100 ml.			8.25 mg./100 ml.

Discussion

Not only was this case particularly acute and severe but it also presented the unusual features of kidney and liver involvement. It is extremely difficult to evaluate the significance of these hepatic and renal changes. To say that the patient died from a hepato-renal syndrome although the polymyositis was contained is not really informative, neither is it fully in accordance with the facts. The term "hepato-renal syndrome," however poorly defined, would appear to include patients dying often in coma, usually jaundiced, and in whom renal and hepatic damage is demonstrated microscopically. In the kidney severe tubule necrosis, most marked in the convoluted tubules, and in the liver cellular damage, which appears to be further enhanced by the presence of excess bile which stains the liver cells, are the usual histological findings (Dible, 1953). In the case under discussion there was very marked renal damage amounting to a cortical necrosis. Microscopically there were changes in both cortex and medulla. The glomeruli were congested, while the convoluted tubules showed a wide range of degenerative change. The liver cells were not bile-stained and there was no evidence of a primary causative lesion. There seems, then, to be considerable evidence to support the view that the renal lesions in this case resulted from the disease process of dermatomyositis.

It is interesting to record the similarity between the nature of renal lesions occurring in other collagen disease, such as disseminated lupus erythematosus and polyarteritis nodosa, and that of the renal lesions here described. In the two former conditions it is a well-established fact that cortisone and A.C.T.H. may suppress all evidence of the disease except when renal lesions occur, in which case the processes are invariably progressive. A similar failure of these hormones would appear to have been responsible for the advanced renal disease in this case. Ragan (1950) reports improvement in two long-standing cases of dermatomyositis treated with A.C.T.H. In the most recent case report by Clein (1953) mention is made of the benefit derived from cortisone and A.C.T.H. He used both hormones with equal success. He also stressed the high polymorphonuclear leucocyte count, which was also present in the case under discussion.

Summary

A case of acute dermatomyositis with necropsy findings is reported. Renal lesions thought to have resulted from the disease process are described. These do not

seem to have been recorded previously. The response of the muscle and skin lesions to cortisone is noted, while the hormone appears not to have influenced the kidney damage.

It is a pleasure to acknowledge the help and advice of Dr. A. Nelson-Jones, under whose care the patient was admitted. Dr. E. James also gave much helpful advice. Dr. S. Marshall kindly supplied the detailed histological report and Dr. F. Brafield did the biochemical investigations. Miss H. Child was most helpful with the clerical work.

REFERENCES

- Clein, L. (1953). *British Medical Journal*, 2, 1357.
Dible, J. H. (1953). In *Recent Advances in Pathology*, edited by G. Hadfield, p. 302. London.
Ragan, C. (1950). *Proceedings of the First Clinical A.C.T.H. Conference*, p. 423. Churchill, London.
Wainger, C. K., and Lever, W. F. (1949). *Arch. Derm. Syph. (Chicago)*, 59, 196.

Medical Memoranda

Tropical Pulmonary Eosinophilia in a West African

This paper reports a case of tropical pulmonary eosinophilia in a Nigerian. So far as can be determined, this condition has not been previously reported as occurring in a West African.

CASE REPORT

The patient was a male Nigerian aged 45, whose occupation was a leprosy inspector.

Previous History.—In 1943 he had pneumonia affecting the left side of the chest and was in hospital for three weeks, being treated with sulphonamides. He was in the Army, in Burma, from 1944 to 1946, and in 1944 had pneumonia affecting both sides of the chest. He was ill for five weeks and was treated with penicillin and sulphonamides. In 1946 he was given oil of chenopodium for ascariasis. Later that year he was discharged from the Army in the Owerri province of Nigeria. He remained there until 1952, when he went to Calabar. He returned to Owerri province one month before he was seen with the current complaint.

Present History.—For six weeks he had complained of swellings in the skin on different parts of the body, each lasting a few days. Three weeks before being seen he had developed a dry cough, which was worse on lying down. The cough had improved slightly during the last week. For three weeks he had been short of breath on exertion—for example, on riding his bicycle on the flat. He also complained of a burning sensation all over the inside of his chest, and had been feeling generally off colour.

Examination showed that he was well nourished. Swellings were seen on right cheek, right calf, and right forearm which were probably Calabar swellings. Respiratory system: no impaired movement of the thoracic cage or impaired percussion note; medium crepitations present at both lung bases. Cardiovascular system: N.A.D. Gastro-intestinal system: N.A.D. No hepatomegaly or splenomegaly. Other signs: lymph nodes in the groins and axillae were palpable but unlikely to be pathological. The patient was afebrile.

Laboratory Investigations.—Haemoglobin, 14 g.; R.B.C., 5,380,000 per c.mm.; W.B.C., 37,800 per c.mm. (polymorphonuclear leucocytes 5%, eosinophils 88%, lymphocytes 7%). No microfilariae were seen in thick blood films made at 12 noon. No autohaemagglutination occurred either at room temperature or when the blood was put into a refrigerator. The stool did not contain any helminth ova. Sputum: no specimen available. X-ray not available.

The blood count remained of the same order over the next seven days, after which penicillin therapy was started with 300,000 units daily for three days. Unfortunately the patient did not receive a full course of penicillin, as the observer left Nigeria at that point. One week after the beginning of the penicillin therapy the total white cell count

had fallen from 37,800 to 24,700 per c.mm. and the eosinophils from 88 to 73%—that is, from 33,264 to 18,031 per c.mm. Two months later, without any further treatment, the white cell count had fallen to 9,400 per c.mm. and the eosinophils to 60% (5,640 per c.mm.). Four months after penicillin the white-cell count was 12,000 per c.mm. and the eosinophils 57% (6,840 per c.mm.). The patient stated that he felt better.

COMMENT

It is regretted that an x-ray examination of the chest could not be undertaken. However, in view of the clinical and haematological findings it would appear likely that the patient was suffering from tropical pulmonary eosinophilia. The possible presence of a *Loa loa* infection would produce a degree of eosinophilia, but hardly one of 88% (33,264 eosinophils per c.mm.). There was no evidence to suggest the presence of another helminth infection. Hunter (1946) reported a case of tropical pulmonary eosinophilia in a European aged 37 who had been resident in Nigeria for 12 years. Wilson (1947) reported the condition in three Africans in East Africa, but it is believed that the present paper reports the occurrence of the condition in a West African for the first time. It is interesting to speculate whether this patient acquired the disease in India some eight years previously and that it had only become manifest recently, or whether it had been acquired in Nigeria.

It is not possible to say with any assurance whether or not there had been any response to penicillin in this case, although there is a suggestion that the drug had some effect.

This case was seen at the Leprosy Research Unit, Uzuakoli, Nigeria. Thanks are due to Dr. J. Lowe for permission to investigate the case and to publish the findings, and to Sister O. Smith for her co-operation.

JAMES A. MCFADZEAN, M.D.,

National Institute for Medical Research, London.

REFERENCES

- Hunter, E. A. (1946). *British Medical Journal*, 1, 877.
Wilson, H. T. H. (1947). *Ibid.*, 1, 801.

Long-term Prognosis in Infantile Eczema

In 1953 Purdy published the results of an investigation into the after-history of cases of infantile eczema in Edinburgh. To amplify his findings a similar investigation into the cases of infantile eczema occurring in Dundee over the same period was carried out. Letters were sent to those patients who attended the skin department of Dundee Royal Infirmary with infantile eczema during 1931–7 inclusive. The questionnaire adopted was the same as that used by Purdy, but in addition questions were included regarding the onset and disappearance of asthma. In order to have results as closely related as possible to those of Purdy, the same criteria were followed in the selection of cases from the hospital records.

RESULTS

In a large number of cases it was impossible to locate by letter persons whose last known address was that of 15 to 22 years ago. Thus Table I shows that, of 136 questionnaires sent out, 92 (67.6%) were returned as "dead" letters.

TABLE I.—Result of Questioning

Total No. of persons sent letters	136 (94 males, 42 females)
"Dead" letters returned	92
Useful replies	37 (20 males, 17 females)
Useless replies (deaths at early age)	2
Replies not returned	5

In the 37 useful replies received, 12 patients claimed to have had no sequelae to their primary attack of infantile eczema. Table II gives the incidence of the generally accepted sequelae and compares the figures found in this study with those in Purdy's series. In assessing the final results of the Besnier's prurigo group all patients were examined personally unless they had been seen recently by a consultant of this department.

TABLE II.—Results of 37 Replies

	No.	Asthma	Asthma Denied; Wheezing Admitted	Hay-fever	Besnier's Prurigo	Migraine
Male	20	9	1	5	10	1
Female	17	4	4	5	5	3
Total	37	13 (35.1%)	5 (13.5%)	10 (27%)	15 (40.5%)	4 (10.8%)
Purdy's series	93	37 (39.8%)	10 (10.7%)	39 (41.9%)	23 (24.7%)	7 (7.5%)

The overall figure for asthma is 35.1%. If the five patients who failed to return the questionnaire are classed as having no asthma then a minimum figure of 30.9% is reached. This minimum is too low, as it assumes that none of the wheezing group of five patients and none of the "lost" five patients are asthmatic. If it is conceded that the wheezing group are asthmatic a minimum of 42.8% is reached, and excluding the "lost" five patients the figure is 48.6%. This range of roughly 43 to 49% is similar to Purdy's of 30 to 50%. Asthma developed in 13 patients in the first decade and disappeared in four males and one female in the second decade.

The figures for Besnier's prurigo show a range of 35.7 to 40.5%; the minimum figure presumes a complete absence of the complaint in the five patients who failed to return the questionnaire.

Table III shows a regrouping of the patients and agrees with Purdy that asthma without skin involvement is the most frequent sequela, with a range of 23.8 to 27%.

TABLE III.—Regrouping of Cases from Table II

	No.	Asthma (or Wheezing Attacks) but No Besnier's Prurigo	Besnier's Prurigo but No Asthma (or Wheezing)	Both Asthma (or Wheezing) and Besnier's Prurigo	Hay-fever but No Asthma (or Wheezing)
Male	20	5	5	5	1
Female	17	5	2	3	
Total	37	10 (27%)	7 (18.9%)	8 (21.6%)	1 (2.7%)
Purdy's series	93	27 (29%)	5 (5.4%)	20 (21.5%)	14 (15%)

Besnier's prurigo in association with asthma is next in frequency, with a range of 19 to 21.6%. Besnier's prurigo alone has a range of 16.6 to 18.9%, which is much higher than that observed in Purdy's series—namely, 4 to 5.4%.

Hay-fever was present in 27% cases in Table II, but in the regrouping in Table III only 2.7% claim to have hay-fever without asthma.

Migraine occurred in four patients, three of them females. I agree with Purdy that it is a very doubtful outcome of infantile eczema.

The answers to questions regarding nervous stability and intelligence were so varied that it was impossible to support or refute the widely held belief that sufferers from infantile eczema are compensated with a high intelligence.

COMMENT

Of the 37 (27%) useful replies received to a questionnaire similar to Purdy's, 25 describe at least one of the recognized sequelae of infantile eczema. Of these 25, 10 had asthma, 8 Besnier's prurigo and asthma, and 7 Besnier's prurigo alone. Although the number of useful replies in this investigation was disappointingly low, it was decided to publish the results so that they may be added to Purdy's findings, and to any subsequent figures recorded from other centres, in order that the ultimate total number may be high enough to yield an accurate picture of the sequelae of infantile eczema.

I wish to thank Dr. J. Kinnear for permission to use his cases and for his helpful advice.

O. A. FINN, M.D.,

Senior Registrar, Department of Dermatology,
Royal Infirmary, Dundee.

REFERENCE

- Purdy, M. J. (1953). *British Medical Journal*, 1, 1366.