in its intensity. His blood pressure was 108/70. The lungs were clear. He could now be classified as grade I.

Comment

The diagnosis of dextrocardia with mitral stenosis was well established from physical findings, electrocardiographic studies, and angiocardiology. The aetiology of mitral stenosis was only presumptive before operation, but was established beyond doubt from histological study of the atrial wall.

Dextrocardia of the mirror type with situs inversus is probably not as rare as is commonly believed; but unless the condition is associated with other congenital cardiac anomalies or cardiac disease no symptoms are produced. These symptoms, when they do appear, are those of the associated abnormality and not due to the dextrocardia *per se*. The signs in the heart are, of course, found on the right chest instead of on the left.

It is claimed by some workers that the condition is common in white persons. It is worth noting, however, that the case reported prior to this was in a negro (Bleyer and Saphir, 1953), and our case is in an Indian.

Summary

A case of pure mitral stenosis due to rheumatic infection in a true dextrocardiac (with situs inversus) is described.

The successful performance of valvotomy in such a case is reported.

REFERENCES

- Baker, C., Brock, R. C., Campbell, M., and Wood, P. (1952). *British Medical Journal*, 1, 1043.
Bleyer, J. M., and Saphir, W. (1953). *Amer. Heart J.*, 46, 772.
Goldberger, E. (1950). *Unipolar Lead Electrocardiography*, 2nd ed. Lea and Febiger, Philadelphia.
Mounsey, P. (1953). *Brit. Heart J.*, 15, 135.

IDIOPATHIC PULMONARY HAEMOSIDEROSIS IN A YOUNG ADULT

BY

N. A. J. HAMER, M.B., M.R.C.P.

Captain, R.A.M.C.

(From the British Military Hospital, Singapore)

Idiopathic pulmonary haemosiderosis is being recognized with increasing frequency in recent years. Most of the cases described are in children, and the mortality rate is high. Considered in retrospect, the clinical picture is remarkably uniform, but early diagnosis may be difficult. Several cases of milder forms of the condition in young adults have been reported. In these the outlook is generally good and long remissions may occur. A further adult case is presented in this article.

Case Report

A National Service man aged 19 was first admitted to hospital in October, 1948, when aged 13, with a three-weeks history of intermittent frontal headache, pallor, and weakness. He described about six similar episodes since an attack of chicken-pox a few months previously. There was some dyspnoea on exertion. For four weeks he had been coughing up moderate amounts of brownish-red sputum. Oral iron therapy had had no apparent effect on the pallor. He had four sisters and nine brothers, all well except a brother with pulmonary tuberculosis. There had been no previous illness of significance.

On examination he was quiet and intelligent, and gave a clear story. He was very pale and listless. There were

no signs of cardiac failure or jaundice, and no haemorrhagic manifestations. His tongue was normal. There were no enlarged lymph nodes and the spleen could not be felt, but a sharp rubbery liver edge was palpable 1 in. (2.5 cm.) below the costal margin. The pulse rate was 110 per minute and the blood pressure 120/80 mm. Hg. The heart and chest were normal, as was the central nervous system.

He was transfused immediately after admission to hospital. The haemoglobin level after transfusion was 9 g.%. This rose to 11 g.% after two weeks, and remained at this level until discharge one week later. No iron therapy was given at this time. The E.S.R. and the leucocyte count were normal. No urobilinogen was found in the urine.

The patient did not attend for observation, but was re-admitted with severe anaemia six months after leaving hospital. Dark-red sputum was produced during most of the time between the admissions to hospital. There were no new physical signs. The haemoglobin value was 4.1 g.%. The red cells showed anisocytosis and were poorly filled with haemoglobin. The icteric index was 8 units, and no urobilinogen was found in the urine. A red-cell fragility test showed haemolysis from 0.56% to 0.32% saline, almost the same as a control.

After a week, treatment with oral iron was started and the haemoglobin level rose rapidly, reaching 12.4 g.% after four weeks, when the patient was discharged. During this period the chest was x-rayed, but no abnormality was reported. He remained completely free of symptoms from July, 1949, until December, 1953, a period of 4½ years.

In December, 1953, the cough with reddish-brown sputum returned. This attack lasted six weeks. There were several minor recurrences during the next few months. He began his national service in October, 1953, and at the end of April, 1954, he arrived in Singapore; the haemoptyses appeared again a few days later. After three weeks he reported sick and a radiograph of the chest showed miliary mottling of the lung fields.

On admission to hospital his only symptom was a mild frontal headache on exertion. Apart from a slight sternal depression no abnormalities were found on examination. The mucosae were of good colour and there was no cyanosis or jaundice. The mouth and tongue were normal. There was no finger-clubbing. No lymph nodes were enlarged, and the spleen and liver were not palpable.

Investigations.—Several blobs of dark-brown sputum with a little mucus were produced each day. On microscopy numerous macrophages containing haemosiderin granules were seen, and a very few red blood cells. No acid-fast bacilli were found on direct examination or culture of six specimens. X-ray examination of the chest showed a normal cardiac shadow, confirmed on screening. There was diffuse mottling throughout the lung fields, maximal in the middle zones. There was no suggestion of a clear centre in any of the markings. Blood picture: Hb, 14.5 g.%; red cells, 5,000,000 per c.mm.; P.C.V., 45%; E.S.R., 36 mm. in the first hour (Wintrobe). Absolute values: M.C.V., 90 cubic μ ; M.C.H., 29 $\gamma\gamma$; M.C.H.C., 32%. Reticulocyte count, 2.5%; leucocyte count, normal; platelet count, 200,000 per c.mm.; bleeding-time, 2 minutes; clotting-time, 4 minutes; prothrombin time, 15 seconds (equal to control). Capillary fragility test, negative. The indirect Coombs test and trypsin cell test showed no antibodies. Cold agglutination was negative at 1 in 4. Red-cell fragility showed haemolysis from 0.52% to 0.28% saline. Blood W.R. and Kahn test were negative; serum bilirubin, 0.5 mg. per 100 ml. on two occasions; serum proteins, 7.5 g.% (albumin 5.5 g.%, globulin 2 g.%). An E.C.G. was normal. The urine was normal on repeated tests, and no urobilinogen was present. The Mantoux test was negative to 1 in 10,000, but positive to 1 in 1,000 O.T. X-ray films of the skull, hands, long bones, and abdomen were normal.

During three months' observation there has been little change in the clinical condition. Daily haemoptyses continue, but the haemoglobin has remained between 13.3 and

14.6 g.%. The reticulocyte count has varied from 2.5 to 0.7%. No iron has been given during this period.

On the first few days in hospital there was a mild evening pyrexia up to 100° F. (37.8° C.), but this quickly disappeared. Slight headache recurs from time to time.

Aetiology

The basic pathological process in idiopathic pulmonary haemosiderosis is repeated intrapulmonary haemorrhage, resulting in deposition of haemosiderin in the lungs.

Pulmonary haemosiderosis also occurs after prolonged pulmonary congestion in mitral stenosis and in chronic left ventricular failure (Lendrum, 1950), and the pathological effects on the lungs are similar to the idiopathic forms. According to Hastings-James (1952), haemosiderosis may follow any form of pulmonary haemorrhage repeated over long periods. In the idiopathic form, however, fresh blood in the sputum is most unusual. It is probable that in these cases haemorrhage occurs diffusely into the alveoli themselves, haemosiderin is formed, and is engulfed by macrophages before the material is produced as sputum. The mechanism differs from the secondary types of pulmonary haemosiderosis in which there is focal bleeding followed by widespread absorption of the pigment.

Three main causative factors have been postulated as the possible fundamental lesions in idiopathic pulmonary haemosiderosis:

Defects in Pulmonary Arterioles and Elastic Tissue.—The extensive changes in the elastic tissue of the lungs seen in most cases at necropsy suggested that a defect in the elastic fibres of the pulmonary arterioles might be the primary lesion in pulmonary haemosiderosis. However, the necropsy assessment of elastic tissue change is difficult owing to the varying effect of distension (Nancekievill, 1949), and it is possible that some of the appearances seen were due to distortion. In some cases with only a short history before death there is no evidence of defect in the elastic tissues. On the whole it appears that these changes are the result, rather than the cause, of haemosiderosis, and the degree of elastic-tissue damage is roughly proportional to the duration and intensity of the disease.

Functional Changes in Pulmonary Capillaries.—Even those cases with a short history and no change in the elastic tissues show capillary stasis at necropsy. It is possible that dilatation of the capillaries in the pulmonary interstitial tissue and changes in the permeability of the capillary walls may lead to diapedesis of red blood cells and subsequent haemosiderosis. Recurrent haemorrhages will initiate fibrosis and destruction of elastic tissue.

Haemolytic Anaemia.—The haematological picture is of recurrent loss of red blood cells in the absence of external haemorrhage, associated with a compensatory increase in erythropoiesis. There is no evidence of decreased stability of the red blood cells or of failure of the blood-clotting mechanism. Cold auto-agglutinins are present in some cases in the acute attack. Wyllie *et al.* (1948) suggest that this is a secondary antibody response to red blood cells destroyed within the body. A positive Coombs test has also been found on occasion. Few cases have been examined in detail for circulating haemolysins, but in one case the haemolytic activity of the serum at body temperature was increased (Wiesmann *et al.*, 1953), and in another the life span of the patient's red blood cells on transfusion was shorter than normal (Jonsson *et al.*, 1951). If a haemolytic process is responsible for idiopathic pulmonary haemosiderosis, it is difficult to see why the lesions should be restricted to the lungs, unless the physiological conditions in the pulmonary capillaries increase the activity of the haemolysin.

From the present evidence it seems most likely that the fundamental lesion in idiopathic pulmonary haemosiderosis is a change in permeability of the pulmonary capillaries. This results in intrapulmonary bleeding, which produces fibrosis and elastic-tissue damage. Erythrocyte destruction within the body may cause the production of antibodies.

Discussion

The occurrence of idiopathic pulmonary haemosiderosis in adult life is rare. Kerley (1951) gives a brief description of five adult cases, and Cliff (1953) describes the case of a man who first had symptoms at the age of 20. After two acute attacks he was under observation for three years, and during this time had no symptoms or anaemia, but haemoptyses continued. This closely resembles the present case.

It seems likely, as Kerley (1951) suggests, that there is a mild form of the disease which persists for many years and may be found in young adults. These subjects may have typical recurrent attacks of severe anaemia in the later years of childhood or young adult life, and later settle down to a steady state with recurrent minor haemoptyses but no anaemia. Similar cases will probably be discovered more frequently with the increasing use of mass miniature radiography, and idiopathic pulmonary haemosiderosis must be considered in the differential diagnosis of miliary x-ray shadows in the chest of a symptomless young adult. Radiological changes may persist for several years after the anaemia and haemoptyses have disappeared. A diagnosis of Lederer's anaemia was made in the present case during the first acute attack. Some cases previously included in this group, which remained well after transfusion, may well be in fact idiopathic pulmonary haemosiderosis.

The diagnosis in the early stages is often difficult, as there may be no radiological changes at first. Walton and Williams (1951) found none for 18 months in their case, and in the patient described in the present article a chest x-ray film was reported as normal during the acute phase of the disease. Young people with anaemia of obscure origin should have sputum or gastric washings examined for macrophages containing haemosiderin. If x-ray changes are present lung puncture must be considered, to confirm the diagnosis. This is not without hazard, as the widespread destruction of elastic tissue that occurs greatly decreases the retractility of the lung, and may possibly prolong bleeding if a large vessel is damaged. It was thought to be inadvisable in the present case.

Summary

A case of idiopathic pulmonary haemosiderosis is described, and the aetiology of the condition is discussed.

The existence of a milder form of idiopathic pulmonary haemosiderosis in young adults is brought to notice.

It is suggested that the disease be more often considered in the differential diagnosis of miliary shadows in the chest x-ray film and of obscure anaemia in young people.

My thanks are due to Brigadier D. Bluett, O.B.E., for permission to publish this article.

REFERENCES

- Cliff, J. M. (1953). *Brit. J. Radiol.*, **26**, 302.
Hastings-James, R. (1952). *J. Fac. Radiol.*, **4**, 44.
Jonsson, B., Vahlquist, B., and Agner, K. (1951). *Blood*, **6**, 665.
Kerley, P. (1951). *Textbook of X-ray Diagnosis*, 2nd ed., edited by S. C. Shanks and P. Kerley, **2**, 178. London.
Lendrum, A. C. (1950). *J. Path. Bact.*, **62**, 555.
Nancekievill, L. (1949). *British Medical Journal*, **1**, 431.
Walton, M., and Williams, A. A. (1951). *Ibid.*, **2**, 390.
Wiesmann, W., Wolvius, D., and Verloop, M. C. (1953). *Acta med. scand.*, **146**, 341.
Wyllie, W. G., Sheldon, W., Bodian, M., and Barlow, A. (1948). *Quart. J. Med.*, **17**, 25.

The Ministry of Health has sent a memorandum to hospital authorities (H.M.(55)35) explaining that notifications of patients diagnosed in hospital as suffering from tuberculosis should usually be sent to the medical officer of health of the district in which the patient normally lives. Where, in the case of long-stay patients, the hospital itself can reasonably be regarded as the patient's place of residence, the notification should be sent to the medical officer of health of the district in which the hospital is situated.