EMERGENCIES IN GENERAL PRACTICE

TREATMENT OF SEVERE ASTHMA

BY

A. MARGARET C. MACPHERSON, M.D., F.R.C.P.*Physician, Elizabeth Garrett Anderson Hospital for Women*

The development of the first really severe attack of asthma is an alarming event for the patient, for the patient's friends, and, at times, for the doctor. Usually there has been some previous evidence of an asthmatic tendency which may have amounted to little more than discomfort. Then an attack of tightness in the chest lasts longer than usual, increases in severity, and the patient's efforts become concentrated simply in trying to breathe out the air which overfills his lungs. Any attempts his relatives or friends make to help are of little avail and their distress at standing by and watching his gasping respirations only too often adds to the patient's alarm. It is in this atmosphere that the doctor finds himself. Successful treatment depends to a great extent on the tacit assumption that the attack can and will be treated successfully. It is rare for an asthmatic attack to be fatal, unless it occurs in elderly patients, or is complicated by other conditions, such as spontaneous pneumothorax, or unless wrongly treated by respiratory sedatives.

Much depends on the way in which the situation is handled. The doctor must be prepared to give up time to the patient's treatment, and time spent in initial treatment is time saved in the long run. The patient will usually have found the position which gives greatest relief, either sitting forward in bed or sitting up in a chair.

Antispasmodic Treatment

Treatment of the bronchospasm should be begun at once. It is rare that adrenaline is unsuccessful, provided it is given correctly, and it is important to give it with care when first used. Once the patient appreciates its use he will look forward with confidence to its benefits during subsequent attacks; if it is used inadequately it may need much persuasion and perseverance on the next occasion to overcome his prejudice against it.

A dose of 5 minims (0.3 ml.) of adrenaline, 1 in 1,000, is given hypodermically. If this dose does not give relief after 10 minutes, adrenaline should be injected at the rate of 1 minim (0.06 ml.) a minute, keeping the needle *in situ*, until the spasm is relieved. In this way large doses of adrenaline can be given without ill effects.

In those cases in which adrenaline is ineffective, aminophylline can be given. It is best administered by intravenous injection, the contents of one ampoule containing 0.25 g. in 10 ml. of water being injected slowly. This can be repeated after four hours, and if necessary the dose can be doubled provided the total of 20 ml. is given slowly. Intravenous treatment may usefully be supplemented by giving the aminophylline in a rectal suppository. This is a convenient way of maintaining some antispasmodic action over a prolonged period and is particularly useful for administration at night.

Antihistamine drugs are usually of little help in relieving severe asthmatic attacks.

Sedatives

During the acute stage sedative treatment is usually inadvisable. As the attack passes off, small doses of sedative can be given to calm the patient and to help ward off another attack. Phenobarbitone in $\frac{1}{2}$ -gr. (32-mg.) doses is usually effective. Opium derivatives and large doses of barbiturates

are contraindicated. The patient's life may depend on his respiratory efforts, and depression of the respiratory centre, particularly by the use of morphine, may prove fatal.

Oxygen

If there is marked cyanosis, oxygen will often help the patient. It should be given with a B.L.B. mask or "polymask." The administration of oxygen, however, may distress the patient more than benefit him, as the application of anything over the mouth or nose may increase the sense of suffocation, and the very suggestion of oxygen may be associated in the patient's mind with the terminal stages of severe illness.

A.C.T.H. and Cortisone

A.C.T.H. and cortisone are on occasions of definite use in treating asthma, but, with our present knowledge, they should be used with caution. A.C.T.H. and cortisone are given in doses of 100 g. in divided doses six-hourly during the first 24 hours and continued, with reduced dosage if possible, for three or four days; cortisone can be given by mouth. There is no doubt that in many instances this treatment will give definite relief. If, however, the treatment has to be repeated, its results, using the same dose, may be disappointing and an increased amount will be required to produce the same effect. Large doses and more continued treatment are not without danger to the patient. At the present time it is probably advisable to withhold these remedies, giving them only when there is urgent need. They are usually called for in treating elderly patients with emphysema who are in great distress and in danger of developing cardiac failure, and also in cases in which the acute attack is prolonged and is developing into status asthmaticus.

Status Asthmaticus

If the attack becomes prolonged, with no remissions, and extends over a period of several days and nights, the condition then is indeed a difficult one to treat. It is advisable whenever possible to remove the patient from home surroundings, preferably into a hospital. The atmosphere of a hospital ward, with efficient nursing and the knowledge that there is always medical help at hand if required, will do much to quieten the patient. Continuous treatment with aminophylline can be carried out if required by slow intravenous drip, using 0.5-1.5 g. of aminophylline in a litre. At this stage the patient is often suffering from some degree of dehydration, since the continued breathlessness has made drinking and eating difficult or almost impossible. By giving aminophylline intravenously, the necessary fluid can be supplied, and it may be useful to use 5% glucose-saline as the vehicle for the drug. The patient should also be encouraged to take sips of water by mouth. When cyanosis is marked, an oxygen tent may be necessary. Cortisone or A.C.T.H. may be required and should be given if other treatments have been tried and failed, or, as already indicated, should be given to patients with severe emphysema or cardiac failure.

Treatment after the Attack

As soon as the attack has been controlled, it becomes necessary to take steps to prevent another attack developing. A small dose of sedative such as phenobarbitone, $\frac{1}{2}$ gr. (32 mg.) two or three times during the day will often quieten the patient. The first evidence of recurrence of wheezing or tightness in the chest should be treated immediately. An

injection of adrenaline, 3 to 5 minims (0.2 to 0.3 ml.) should be given at once. The earlier this is done, the more likely it is to ward off a fresh attack. If it is not possible for the doctor to give the injection immediately, it is often advisable for suitable patients to be taught how to give the injection to themselves. When this is not feasible, ephedrine, $\frac{1}{2}$ gr. (32 mg.), or isoprenaline, 20 mg., can be taken in tablet form; isoprenaline tablets being dissolved slowly under the tongue. Atomizers may be used to give adrenaline or isoprenaline by inhalation. Many of the proprietary preparations for the treatment of asthma contain one or other of these drugs together with phenobarbitone. Aminophylline suppositories given in the early part of the night may help to prevent an attack in the early hours of the morning. Some patients find antihistamine preparations beneficial, if taken as soon as the first symptoms of spasm are noticed. Expectorant mixtures containing potassium iodide and ammonium carbonate relieve those patients who have difficulty in coughing up sputum after an attack and are of particular use for patients who have chronic bronchitis complicated by asthma.

It is now recognized that one of the most successful treatments in controlling the incidence of attacks and in controlling an attack in its early stage is breathing exercises. If the patient conscientiously practises those exercises which encourage proper diaphragmatic breathing and full expiration, he will be able to reduce the number of attacks and also can often abort an attack. The breathing exercises can be combined with postural exercises when there is evidence of localized chronic infection in the bronchial tree.

Prevention of Pulmonary Infection

Care should be taken to detect pneumonic lesions during the period after an asthmatic attack, particularly in those patients who suffer from chronic bronchitis. During an asthmatic attack, bronchi or bronchioles may become plugged with thick mucous secretion, with resultant segments of collapsed lung beyond. Usually re-expansion takes place, but at times the collapsed portion of the lung becomes infected and prompt treatment with antibiotics is indicated if this complication is suspected.

Next article on Emergencies in General Practice.—
"Post-partum Maternal Collapse," by Miss Josephine Barnes.

Refresher Course Book.—Copies of the second volume of collected articles from the Refresher Course for General Practitioners published in the *Journal* are still available at 25s. (postage 1s.) each. The volume contains 60 articles, all of which were revised by their authors for publication in the book. The first volume is now sold out and out of print. Copies of the second volume may be obtained from the Publishing Manager, B.M.A. House, Tavistock Square, London, W.C.1, or through any bookseller.

PRACTICAL ADVICE ON HEARING-AIDS

In a lecture at the Institute of Laryngology and Otology, London University, on April 29 Mr. J. CHALMERS BALLANTYNE, assistant director of the audiology unit of the Royal National Throat, Nose, and Ear Hospital, discussed the indications for advising a hearing-aid, the choice of a suitable instrument, and the way to get the best results from it.

The ability of a deaf person to hear with an aid, said Mr. Ballantyne, was determined very largely by the type, as well as the degree, of his deafness, and it was essential to have a clear understanding of the differences between the two broad types of deafness—perceptive (or "nerve") deafness and conductive (or "middle-ear") deafness. The hearing apparatus had two parts, a perceiving apparatus and a conducting apparatus. The perceiving apparatus consisted

of the end-organ of hearing itself—that is, the organ of Corti—and the acoustic nerve and its central connexions. The conducting apparatus consisted of the external auditory canal, the middle ear, and the Eustachian tube. It also included the fluids of the inner ear. Its function was to conduct sound-waves from their source to the end-organ, amplifying them to some extent as it did so. Perceptive deafness resulted from any lesion in the perceiving apparatus, while conductive deafness was due to lesions in the conducting apparatus. The two types of deafness could nearly always be distinguished from one another by a careful history, a thorough clinical examination, and a single tuning-fork.

Indications for the Use of Hearing-aids

There were very few deaf people who could not derive some benefit from using an aid, continued Mr. Ballantyne, and it was much better to start using one too early than too late. When deafness was purely conductive in nature, as in early cases of otosclerosis and in suppurative disease of the middle ear, the perceiving apparatus was normal. In theory, all that was required to allow such a patient to hear normally was to amplify sounds to such a degree that they would reach the normal perceiving apparatus at a normal degree of loudness. In practice also, the necessary amplification could usually be provided by an electrical hearing-aid.

When deafness was purely perceptive, as in the deafness following meningitis or the deafness of old age, the problem was much more complex. Amplified sounds reached the abnormal perceiving apparatus with the same degree of loudness as in the normal ear, sometimes even louder. But they were distorted sounds, often uncomfortable and sometimes painful. And it was characteristic of perceptive deafness that, beyond a certain point, the ability to understand speech became reduced and not enhanced by further amplification. On the other hand, many patients suffering from perceptive deafness could derive real benefit from an electrical aid.

Construction of Electrical Aids

Since the second world war the design and construction of electrical hearing-aids had improved enormously. Aids consisted of three parts: a microphone, an amplifier, and a receiver. The microphone converted sound energy into electrical impulses; the amplifier amplified the sounds; and the receiver converted the electrical impulses back into sound energy.

The Amplifier.—Most electrical aids used a system of thermionic valves, the power required by this system being provided by two batteries. A low-tension battery warmed the filaments in the valves, and a high-tension battery provided the power for the amplification. Many commercial firms are now substituting the so-called "transistor" valve for the thermionic valve.

The Receiver.—There were three different types of receiver: the insert receiver, the flat receiver, and the bone-conduction receiver. The insert receiver fitted into the ear itself and was by far the most commonly prescribed. It should be made to measure for each individual user. It could be used even in the presence of active middle-ear discharge, provided that the ear was cleaned regularly by simple aural toilet and the insert kept free from discharge. An external otitis might result in a few cases, when the use of an insert must be discontinued, at least temporarily. The flat receiver was worn with a headband. It was large, uncomfortable, and now rarely prescribed. Leakage of sound was almost invariable, with consequent "howling." The bone-conductor receiver also caused considerable discomfort by the presence of a headband, but it might be indicated in some cases of aural discharge with external otitis. In a very limited number of cases of conductive deafness, not exceeding 2% of all hearing-aid users, it might give a very good performance. One commercial firm now produced a bone-conductor receiver carried on a spectacle frame.