limitation of movements are the prime features and in which the symptoms of acute inflammation have subsided.

Of 50 acute cases treated (see Table), 36 obtained complete relief of symptoms from the injection of hydrocortisone alone within 7 to 14 days. Only three cases failed to improve at all.

Treatment of Periarthritis of the Shoulder with Hydrocortisone

	No. of Cases	Complete Recovery	Partial Improvement	No Change
Acute lesions	50	36	11	3
Chronic "	23	13	5	5

In the chronic group the results were not nearly so good, presumably as the inflammation had subsided and the fibrinolytic action was less powerful. Nevertheless 13 out of 23 cases progressed to a complete recovery with hydrocortisone injections and exercises over a period of four to six weeks.

Conclusions

In the treatment of the various acute and chronic soft-tissue lesions within and around the shoulder-joint hydrocortisone has led to a new method of approach to what was previously a most difficult therapeutic problem.

The majority of both local and generalized acute shoulder lesions showed a dramatic response to hydrocortisone injections into the various components of the rotator cuff.

Although we have had a lower percentage of cases with full relief of symptoms than Robecchi and Capra (1953), we nevertheless feel that the use of the hormone is indicated, at least for a trial period, in every acute case.

The results in the more chronic shoulder lesions, where fibrosis is a prominent feature, have not been so good. This would indeed be expected from consideration of the pathology. In association with routine physiotherapy, however, improvement of the condition has been obtained in a sufficiently high percentage of cases to warrant further trial of the method. Other forms of treatment are so notoriously unsuccessful in the experience of the majority of authors that we feel that hydrocortisone provides a marked advance in what Cyriax and Troisier call the "apparently endless search for effective treatment of 'freezing arthritis.'"

Summary

The pathology of periarthritis of the shoulder-joint and the theoretical effects of hydrocortisone in its various stages are discussed.

The technique of treatment by the injection of hydrocortisone, hyaluronidase, and procaine hydrochloride is described.

Of the acute cases treated by this method 72% progressed to complete recovery in 7 to 14 days and a further 22% were improved.

Of the "frozen shoulder" type of case 56% regained full function in four to six weeks, following hydrocortisone injections and shoulder exercises; 22% were improved.

We consider that the use of hydrocortisone is a notable advance in the treatment of soft-tissue shoulder lesions.

Our thanks are due to Miss M. Waldron, of Guy's Hospital Department of Illustration, for Figs. 1 and 2.

REFERENCES

- Codman, E. A. (1909). *Publ. Mass. gen. Hosp.*, 2, 521.
 — (1934). *The Shoulder*. Boston.
 Crisp, E. J., and Kendall, P. H. (1955). *Lancet*, 1, 476.
 Cyriax, J., and Troisier, O. (1953). *British Medical Journal*, 2, 966.
 Duplay, S. (1872). *Arch. gén. Méd.*, 2, 513.
 Neviaser, J. S. (1945). *J. Bone Jt Surg.*, 27, 211.
 Robecchi, A., and Capra, R. (1953). *Rev. Rhum.*, 20, 757.
 Simmonds, F. A. (1949). *J. Bone Jt Surg.*, 31B, 426.
 Steindler, A., and Marxer, J. L. (1946). *Traumatic Deformities and Disabilities of the Upper Extremity*. Springfield, Ill.
 Wilson, C. L. (1943). *Arch. Surg. (Chicago)*, 46, 307.

LOCAL HYDROCORTISONE IN DE QUERVAIN'S DISEASE

BY

B. G. B. CHRISTIE, M.B., B.S.

Registrar, Department of Physical Medicine and Rheumatism, the Middlesex Hospital

In 1895 Fritz de Quervain, a Swiss surgeon, described five cases of stenosing tenosynovitis at the radial styloid, the prominent symptom of which was pain in the region of the wrist-joint. Subsequent writers have observed that the pain may be referred up the arm as high as the elbow, or downwards to the dorsum of the thumb. The pain appears to be due to movements within the thickened and stenosed synovial sheath, containing the tendons of abductor pollicis longus and extensor pollicis brevis, as it passes under the extensor retinaculum of the wrist in the first dorsal compartment. The thickening of the sheath in some cases is so marked as to be easily visible, and in the majority is palpable. The condition is common, and often the cause of considerable disability to the housewife and worker. Analogous stenosis has been described in all the remaining five tunnels beneath the extensor retinaculum, in most of the sheaths of the flexor tendons, and in the tibialis anterior and peroneal tendons.

Most authorities agree that stenosing tenosynovitis is often due to mechanical factors or trauma, and that histologically the signs of inflammation in the sheaths or tendons are rare.

Since the original description many methods of treatment have been suggested, operation holding pride of place and having been suggested by de Quervain himself. Surgical treatment entails incision of the tendon sheath, or, if this is greatly thickened, excision of a portion. Other more conservative methods of treatment still in use are the inunction of various substances, immobilization of the wrist and thumb in a plaster or plastic splint (Thompson *et al.*, 1951), and the application of physiotherapeutic measures. Injection of the sheath with fluids and local analgesics had also been tried. Although strongly advocated by Ellis (1951), inunction rarely seems to give a lasting cure, and the same applies to physiotherapy. Immobilization often relieves the pain, only to be followed by a recurrence of the symptoms when activity is resumed. According to Wiles (1949) it is doubtful whether many cases recover without operation.

The recent reports of the value of local hydrocortisone in "tennis elbow" and soft-tissue lesions seemed to merit the controlled trial of its use in the present condition. Howard *et al.* (1953) describe its use in three cases of de Quervain's disease and 11 cases of tenosynovitis, and Cyriax and Troisier (1953) report a favourable result in one case of tenosynovitis of flexor carpi radialis following two injections of hydrocortisone. In view of these findings it was felt to be justifiable to treat a number of patients by this method and compare the results with cases treated by injection of local analgesic alone.

Material

The series consists of 20 consecutive cases of de Quervain's disease referred to the Department of Physical Medicine and Rheumatism of the Middlesex Hospital.

Three criteria were held as essential before such a diagnosis was made. These were: (1) Tenderness on palpating the tendons of abductor pollicis longus and extensor pollicis brevis as they lie along the lower outer border of the radial styloid. (2) Pain, radiating up the forearm and down the thumb, on active abduction and extension of the thumb and aggravated by the resisted movement. Passive movements were invariably painless. (3) Pain on flexing the thumb into the palm and closing the remaining fingers around it. This is further aggravated by forced ulnar deviation of the hand at the wrist (Finkelstein's (1930) test). All cases in the present series conformed to these criteria.

In none of the 20 cases reviewed was there any obvious predisposing factor, although in three the symptoms were aggravated by occupational factors (tailor's cutter, sempstress, and poster artist) involving frequent repetitive movements of the thumb. Sixteen of the cases (80%) occurred in females. The right and left hands were affected approximately equally, although all the patients were right-handed. The age incidence varied from 18 to 74 years, there being rather more patients in the younger than in the older age groups. The symptoms had been present for periods varying from four weeks to three years (average six months).

It is interesting to note that crepitus was not elicited in any of these cases in the region of the tendon sheath, either on palpation or on active or passive movements. The thickening of the tendon sheath was visible only as a tumour in one patient, in whom it was very marked. This completely disappeared over a period of five weeks following one hydrocortisone injection with complete return of function, the patient returning to her work as a sempstress entailing continuous use of the thumb.

The patients were divided into two groups: (1) Group A, receiving 1 ml. (25 mg.) of a suspension of hydrocortisone acetate into the tendon sheath following infiltration with 1 ml. of 2% procaine hydrochloride; and (2) Group B, injected with 1 ml. of 2% procaine hydrochloride alone and observed over the ensuing period—when relief was no longer experienced these patients were treated in the same manner as Group A. There are therefore 20 cases in Group A and 10 in Group B, the latter acting as the control series.

Method

The method adopted in all these injections, whether of procaine or hydrocortisone acetate, was standardized as follows. With the patient seated facing the operator, the point of the needle was introduced through the skin about 1 cm. distal to the tip of the radial styloid in the line of the tendon of abductor pollicis longus. An intradermal bleb of local analgesic was raised, and through this the needle was passed, it being directed upwards over the radial styloid so as to enter the common sheath of the tendons concerned. The contents of the syringe were then discharged, with the production of a longitudinal "sausage-shaped" tumour, which was clearly defined and taken as evidence that the injected fluid had actually entered the tendon sheath. In those cases that received both procaine and hydrocortisone injections at the same attendance the procaine was given first, being immediately followed by the hydrocortisone, a second needle and syringe being used.

It was found easier to insinuate the needle into the tendon sheath if the patient adopted a position of moderate ulnar deviation. In all cases the skin was first swabbed with surgical spirit and a small dressing applied after the injection. Patients were instructed to use their wrists normally and informed that no special precautions were necessary.

Results

The patients treated with procaine and hydrocortisone injections (Group A) can be further subdivided into those treated in this manner initially and those treated following procaine injections alone. In seven out of the first ten

patients there was complete relief of symptoms for one hundred days or more following the combined injection. For the purposes of the report this was regarded as a clinical cure. The three remaining patients relapsed at intervals after injection, but were relieved by a further injection. One of these, because of his regular relapses, has been referred for surgery, and it seems that the other two may have to be referred in a similar manner.

The patients in Group B showed relief of symptoms varying from one hour to one to two days, gradually waning, so that after five days there was a complete relapse to the original state in all cases. The average duration of the relief was about two days. Although the pain disappeared in all cases immediately after the injection it was often accompanied by a dead feeling of the wrist, which was not experienced when hydrocortisone was used. These patients were subsequently added to Group A.

Of these latter patients when treated with hydrocortisone and procaine injections, one relapsed during the eighth week and is now symptom-free following a second injection. The remaining nine patients have no pain at periods varying from five and a half to fifteen weeks after the injection.

During the treatment of these cases no serious complications were encountered. In two cases there was a complaint of dull aching pain in the wrist following hydrocortisone injection, but this subsided in a few days without treatment. One case developed a superficial thrombophlebitis, which responded to strapping for ten days.

Discussion

The recent use of hydrocortisone in other soft-tissue lesions prompted this survey of its use in de Quervain's disease. The results of this small series are encouraging and seem to merit further trial of the method. It appears that hydrocortisone has a place in the treatment of this troublesome condition and may be used as an alternative to surgery in many cases. With supplies of hydrocortisone becoming more plentiful this simple method of treatment may become more widespread, thereby making it unnecessary for most patients to undergo operation. In those cases in this series in which treatment has succeeded, full function has been restored and all pain relieved almost immediately after the injection, although in one or two there was a time interval of several days before full relief was experienced. It is difficult to retain patients for long periods of observation when complete relief of symptoms has ensued. Nevertheless, although relief for one hundred days has been regarded as a cure for the purpose of this series, it is known that several patients have had no relapse for periods approaching six months. It is suggested that cases of de Quervain's disease should first be treated by local hydrocortisone injection, and that those who relapse within a short period be referred for operation in the usual way. It will also prove valuable in patients not wishing to undergo surgery and has the great advantage of leaving no scar. If further experience substantiates these findings it would appear to be the treatment of choice in de Quervain's disease and in tenosynovitis occurring at other sites.

Summary

De Quervain's disease is described and various methods of treatment are reviewed. Attention is drawn to the fact that analogous stenosis occurs in many other sites.

Twenty cases have been treated with local injection of 1 ml. (25 mg.) of a suspension of hydrocortisone acetate and 1 ml. of 2% procaine hydrochloride solution. Ten controls were treated with an injection of 1 ml. of 2% procaine alone.

The figures show an advantage in favour of local injection of hydrocortisone, 70% of the patients having remained free of symptoms for periods exceeding three months following one injection.

The results of this small series are encouraging, and seem to merit the further trial of hydrocortisone in de Quervain's disease and in tenosynovitis occurring at other sites.

I wish to thank Dr. A. C. Boyle for permission to quote these cases and for his interest and many helpful criticisms at all stages of the survey; Mr. Philip Wiles and Mr. P. H. Newman for kindly referring cases; and Roussel Laboratories for generously supplying all the hydrocortisone used.

REFERENCES

- Cyriax, J., and Troisier, O. (1953). *British Medical Journal*, 2, 966.
De Quervain, F. (1895). *Korres-Bl. schweiz. Aertz.*, 25, 389.
Ellis, M. (1951). *British Medical Journal*, 2, 777.
Finkelstein, H. (1930). *J. Bone Jt Surg.*, 12, 509.
Howard, L. D., jun., et al. (1953). *Ibid.*, 35, 526.
Thompson, A. R., Plewes, L. W., and Shaw, E. G. (1951). *Brit. J. industr. Med.*, 8, 150.
Wiles, P. (1949). *Essentials of Orthopaedics*, 1st ed., p. 321. Churchill, London.

EFFECTS OF MODIFIED E.C.T. ON THE ELECTROCARDIOGRAM

BY

RONALD GREEN, M.B., F.F.A.R.C.S.

AND

ARNOLD WOODS, B.M., M.R.C.P.

From St. George's Hospital, London

There have been several investigations into electrocardiographic changes during electric convulsion therapy (E.C.T.). Most of them were carried out before the use of muscle relaxants, and consequently the records could be taken only after the end of the convulsion. The purpose of this paper is to record our findings in patients whose convulsions were attenuated by muscle relaxants to such an extent that the electrocardiogram (E.C.G.) could be recorded during the convulsion, thereby providing further information regarding the safety of the procedure.

Method

Relaxation was achieved by the use of suxamethonium chloride, but, as the slightest movement of the limbs during stimulation interfered with the tracings and made their interpretation difficult or even impossible, doses far in excess of those normally used for routine treatment had to be employed. Four milligrams of suxamethonium per stone (0.6 mg. per kg.) of body weight gives adequate attenuation of the fit for routine treatments in the majority of patients, but for our purpose it was necessary to administer up to 10 mg./stone (1.6 mg. per kg.) to obtain the degree of relaxation required.

Every patient was fasted for at least four hours and was given atropine, 1/100 gr. (0.65 mg.) hypodermically half an hour before treatment.

After explaining to the patient the nature of the investigation, a resting E.C.G. was taken. (At the beginning of the investigation the three standard leads were recorded on a three-channel Elmquist; later, a direct-writing Cambridge electrified cardiograph was used and Lead II recorded.) A "sleep dose" of intravenous thiopentone was administered slowly, 0.25 g. seldom being exceeded. It was followed immediately by a large dose of suxamethonium from a different syringe, but through the same needle. As soon as apnoea developed the lungs were inflated with oxygen through a face-mask and bag. A second recording was then taken to assess the effect of the relaxant. As the changes noted were slight and were regarded as of no significance, the relaxant was discontinued after 74 records had been taken.

The electric convulsion shock was given with a Plexicon electroplexy unit, while a further tracing was recorded. Except for a short period during the application of the electrodes and the insertion of the Multitone mouth gag as described by Back (1952), which at the same time permits the maintenance of a good air-way, the lungs were inflated with oxygen from the beginning of the apnoea until adequate respirations returned.

Material

We recorded 214 E.C.G.s from 105 patients who received E.C.T., 57 of the patients being investigated on more than one occasion. In 14 records, in spite of the use of the relaxant, there was interference gross enough to make interpretation difficult or even impossible; this occurred mainly in the early cases, before it was definitely known how much suxamethonium was needed to ensure complete inertia. In 50 records there was slight interference, which did not seriously affect the reading of the tracing, whilst 150 were good.

There were 37 males and 68 females in the series, and their ages ranged from 16 to 78. Just over 50% of the female patients were in the age group 30-50, and almost twice as many were under 50 as over 50, whereas amongst the males the cases were almost equally divided between those under 50 and those over 50, with the majority also about equally divided between the 30-50 and 50-70 age groups (see Table).

Age Incidence

	Under 30	30-50	51-70	Over 70	Total
Male	4	15	16	2	37
Female .. .	10	35	19	4	68
Total .. .	14	50	35	6	105

Most of these patients had normal hearts, but in 10 a cardiac abnormality had been recognized clinically or was suspected from the appearance of the control E.C.G. Two men, aged 57 and 77, gave a past history of cardiac infarction and had abnormal E.C.G.s, one showing a right bundle-branch block pattern. In a woman aged 61 there was also a right bundle-branch block picture but no history of cardiac pain and no other electrocardiographic abnormality. In three women, aged 50, 55, and 66, the control E.C.G. suggested old posterior cardiac infarction. Two women, aged 34 and 58, had clinical evidence of mitral stenosis, although not of a severe degree. In a man aged 60 there was evidence of left ventricular hypertrophy, and complete heart block was found in the control record of a woman aged 31.

Results

Effect of Suxamethonium

In the first 74 records the effect of the suxamethonium was studied. The rate became faster in 41, slower in 13, and remained unchanged in 20; in no case was the change in rate marked. There was slight depression of the ST segment or flattening of the T wave in 5 cases, possibly due to temporary inadequate oxygenation immediately following diaphragmatic paralysis. There were no premature beats and no significant conduction changes.

Effect of the Shock

Rate.—In the control records the pulse rate was under 100 in 48% and under 130 in 87%, although a sinus tachycardia of 160 was encountered in 4 cases and of 170 in one case. A marked tachycardia developed within a few seconds of the shock being given. This usually lasted about 30 to 45 seconds, and in 65% of the cases the rate was over 130. In 37% it was between 140 and 160 and in 10% it was over 160, the most rapid rate recorded being 180. In nearly all instances the rate had returned to pre-convulsion figures within two minutes. In a few cases the sinus tachycardia was preceded by a sinus bradycardia lasting two to three