ORIGINAL ARTICLE

Reference chart for relative weight change to detect hypernatraemic dehydration

Paula van Dommelen, Jacobus P van Wouwe, Jacqueline M Breuning-Boers, Stef van Buuren, Paul H Verkerk

Arch Dis Child 2007;**92**:490–494. doi: 10.1136/adc.2006.104331

See end of article for authors' affiliations

Correspondence to: Paula van Dommelen, TNO Quality of Life, PO Box 2215, 2301 CE Leiden, The Netherlands; P.vanDommelen@pg.tno.nl

Accepted 7 July 2006
Published Online First
31 July 2006**Objective:** The validity of the rule of thumb that infants may have a weight loss of 10% in the first days after birth is unknown. We assessed the validity of this and other rules to detect breast-fed infants with hypernatraemic dehydration.**Design:** A reference chart for relative weight change was constructed by the LMS method. The reference group was obtained by a retrospective cohort study.**Participants:** 1544 healthy, exclusively breast-fed infants with 3075 weight measurements born in the Netherlands and 83 cases of breast-fed infants with hypernatraemic dehydration obtained from literature.**Results:** The rule of thumb had a sensitivity of 90.4%, a specificity of 98.3% and a positive predictive value of 3.7%. Referring infants if their weight change is below -2.5 SDS (0.6th centile) in the reference chart in the first week of life and using the rule of thumb in the second week had a sensitivity of 85.5%, a specificity of 99.4% and a positive predictive value of 9.2%.**Conclusions:** The rule of thumb is likely to produce too many false positive results, assuming that for screening purposes the specificity needs to be high. A chart for relative weight change can be helpful to detect infants with hypernatraemic dehydration.

Exclusive breast feeding up to the sixth month of life is important for optimum infant development and growth as breast milk contains all the necessary nutrients in ideal proportions.¹ Breast feeding protects against infections and allergies, and plays a major role in mother–infant bonding.²

In the Netherlands, 78% of mothers initiated breast feeding in the period 2001–2003. After 1 month 51% and after 4 months 25% of infants were fed primarily on human milk.³ The WHO and UNICEF started the “Baby Friendly Hospital Initiative” to promote breast feeding.⁴ In the Netherlands, this program is mainly focused on improvement of support and encouragement of breast feeding in general health care.

Almost all mothers are capable of breast feeding their infant successfully. However, in some cases initial milk supply is insufficient because of a poor start to milk production or transfer. If the infant's needs are not met for several days, dramatic weight loss and an increase in serum sodium concentration occur and the infant develops hypernatraemic dehydration.^{5–7} Hypernatraemic dehydration may cause serious complications, such as fits, disseminated intravascular coagulation and multiple cerebrovascular accidents, and may even result in death.^{8–12}

A retrospective, population-based study reported an incidence rate of hypernatraemic dehydration of 7.1 per 10 000 breast-fed infants.⁶ This is probably the minimum incidence as cases may have been missed because they occurred in infants before initial discharge from hospital^{13 14} or because appropriate investigations were not performed. The clinical day of presentation of hypernatraemic dehydration is usually around 10 days of age.⁶

In clinical practice, weighing is an essential part of the assessment of an infant's growth and hydration status. However, there is no evidence-based consensus for “normal” and “abnormal” early relative weight change (RWC). Several studies reported the normal (50th centile) or extreme (1st, 2.5th or 5th) centile RWC for exclusively breast-fed infants.^{6 15} However, these centiles are not precisely described with respect

to day of measurement nor shown on standard growth charts. Several authors propose different rules of thumbs for identifying “abnormal” RWC.^{16–20} It is suggested that many midwives use the rule of thumb that infants may have a weight loss of 10% ($= -10\%$ RWC) and should regain birth weight by 10–14 days of life.⁵ To our knowledge, no evidence-based referral rule is available to detect infants with hypernatraemic dehydration.

This study describes a reference chart for breast-fed infants between postnatal days 2 and 11. This chart, together with reports of hypernatraemic dehydration obtained from the literature, will be used to define an evidence-based referral rule. The centiles of the chart can be used as a test to detect infants with hypernatraemic dehydration. The test is considered positive if a breast-fed infant's relative weight decreases below a chosen centile and negative if it stays above. Sensitivity, specificity and positive predictive value (PPV) will be used to optimise this rule. This test will be compared to the rule of thumb that infants may have a maximal weight loss of 10%.

METHODS

Population

We selected a representative reference group of healthy, exclusively breast-fed infants and a group of breast-fed infants diagnosed with hypernatraemic dehydration. The reference group was obtained from a retrospective cohort study initiated in three primary care midwife practices in the Netherlands (metropolitan Amsterdam South-East, rural Heerhugowaard and the country town of Veenendaal). In the Netherlands, a midwife either assists the delivery at home or in an outpatient clinic, or is involved in follow-up care after hospital delivery by a gynaecologist. We selected 1544 infants born in 2002 with a weight measurement (in grams) at birth and at least one

Abbreviations: PPV, positive predictive value; RWC, relative weight change; SDS, standard deviation score

Table 1 Characteristics of healthy, breast-fed reference infants

Characteristics	Healthy, breast-fed infants (n = 1544)
Maternal age in years	30 (4.7)
Girls (%)	49
Gestation in weeks (n = 1543)	39.5 (1.4)
Preterm <37 weeks (%)	2.0
Parity (%)	
First	45
Second	36
Third or more	19
Delivery (%)	
Spontaneous	80
Caesarean section	10
By vacuum extraction or forceps	10
Birth weight in kg	3.44 (0.46)

Data are means (SD) or percentages.

weight measurement between postnatal days 2 and 11. The infants were weighed at home by a midwife with a calibrated electronic scale. In this study, the midwife was instructed to weigh the infant routinely.

Infants hospitalised with hypernatraemic dehydration were identified by a literature search. Articles written in Dutch, English, French or German published between 1970 and 2005 that describe infants with hypernatraemic dehydration were obtained using the search program PubMed with the MESH terms "dehydration" and "breastfeeding". References in these articles were used to increase the number of articles describing infants with hypernatraemic dehydration. We assumed that an infant had hypernatraemic dehydration when the author(s) of the article diagnosed the infant as such. In 47 articles we identified 129 cases of breast-fed infants with hypernatraemic dehydration with a weight measurement at birth and day of presentation or a calculated RWC at day of presentation. A total of 83 literature cases had a day of presentation between 2 and 11 days of life and these were used in this study.^{6 9 11 14 19 21-42} Serum sodium concentration was known for 80 literature cases. All infants were born at term.

Statistical analysis

RWC was calculated as the difference in weight at day of presentation ($w(t)$) and birth weight ($w(t_0)$) divided by birth weight in percentage, or in formula: $100\% \cdot (w(t) - w(t_0)) / w(t_0)$. Day of birth was represented by day 0. A reference chart for relative weight was obtained by the LMS method.⁴³ The LMS method summarises the distribution of relative weight as it changes according to age by three curves representing the Box-Cox power (L-curve), the median (M-curve) and the coefficient of variation (S-curve). The L-, M-, and S-curves were used to convert data into standard normally distributed data. Such a data point is called a z score or standard deviation score (SDS). Normality of SDS was tested by so called "worm plots" for different age groups.⁴⁴ A log power-transformation was applied to age in the LMS method. Since the LMS method works only with positive values, an amount of 25% was added to relative weight and afterwards subtracted from the centiles. Each infant had multiple weight measurements. All weights were included in the analysis and were treated as independent as we did not find an association between the number of measurements and birth weight (t test, $t = 1.14$, $p = 0.26$).

The centiles of the curve and the 10% weight loss were used as a test. The specificity of the 10% rule was calculated as the mean of the percentiles of the reference chart that have a 10% weight loss for each day. To calculate PPV, we assumed that the

Table 2 Number of measurements between 2 and 11 days of life in healthy, breast-fed infants and infants with hypernatraemic dehydration

Characteristics	Healthy, breast-fed infants	Infants with hypernatraemic dehydration
Number of infants	1544	83
Number of measurements on		
Day 2	9	0
Day 3	505	9
Day 4	263	9
Day 5	618	4
Day 6	128	15
Day 7	287	10
Day 8	272	11
Day 9	864	6
Day 10	93	16
Day 11	36	3

incidence of hypernatraemic dehydration is 7.1 per 10 000 breast-fed infants.⁶

Calculations for the LMS method were performed with LMS Light version 1.16 (Institute of Child Health, London, UK) compiled on 15 April 2002. All other analyses were performed with S-plus version 6.2 (Insightful, Seattle, WA, USA).

RESULTS

The characteristics of the reference infants are given in table 1. The number of measurements in reference infants and in those with hypernatraemic dehydration are shown in table 2.

RWC was not normally distributed (Shapiro-Wilk normality test: $W = 0.975$, $p < 0.01$). To obtain normally distributed SDS for RWC, we used the LMS method with a Box-Cox power transformation of approximately 0.5. Normality of SDS was tested by worm plots of different age groups. The shape of the worm plots was reasonably flat, indicating that the data follow the assumed distribution in this age period.

Figure 1 shows a reference chart with standard deviation lines of the RWC of healthy, breast-fed infants as well as the RWC of 83 infants with hypernatraemic dehydration on the day of presentation. The rule of thumb of 10% weight loss is also indicated on the chart. The standard deviation lines or percentiles on this chart show which percentages of infants have the same RWC. For example, if a 5 day old infant weighs 3315 g and has a birth weight of 3750 g, then the calculated RWC is $100\% \cdot (3315 - 3750) / 3750 = -11.6\%$. Notice that -11.6% RWC at day 5 on the chart corresponds to -2.6 SDS or the 0.5th percentile. This means that only 0.5% of 5 day old infants have a RWC less than this infant. To avoid the user calculating weight as a percentage, we converted the -2.5 SDS RWC centile to weights by age for a given birth weight. This converted -2.5 SDS centile is shown on fig 2 for different birth weights. The infant in the previous example has a birth weight of 3750 g. The -2.5 SDS centile for this infant is shown by the fourth line from the top, starting at 3750 at day 0. Follow this line until you reach day 5 and notice that 3315 g at day 5 is just below the line.

Maximal negative RWC for healthy, breast-fed infants is at 3 days after birth, with a mean RWC of -6.0% (95% CI: -5.7% to -6.2%). The mean increases by approximately 1% per day from -6% at day 3 to 0% at day 8. However, even after 11 days about a third of these infants have not yet regained their birth weight; in contrast with healthy, breast-fed infants, the mean of these patients is consistently declining. The mean RWC for the infants with hypernatraemic dehydration is -18.5% (95%

Figure 1 Reference chart with standard deviation lines of relative weight change ($= 100\% \times (\text{weight} - \text{birth weight}) / \text{birth weight}$) for healthy, breast-fed infants as well as relative weight change in 83 infants with hypernatraemic dehydration on day of presentation (day of birth is day 0) and showing the rule of thumb of 10% weight loss.

CI: -17.0% to -19.9%). The mean decreases by approximately 2% per day from -10% at day 3 to -25% at day 10.

Notice that there were no cases of hypernatraemic dehydration before day 3, probably due to the fact that it takes some time before insufficient breast feeding leads to weight loss. We therefore applied the rules from 3 days up until 11 days after birth.

Table 3 shows sensitivity, specificity and PPV for several referral rules: the rule of thumb (10% test), the SDS rules and a combination of the -2.5 SDS test in the first week (3–6 days after birth) and the 10% test after the first week. All sensitivities for these tests were above 85% with less than 3% false positives. The sensitivity of the 10% test was similar to the that of the -2 SDS rule and specificity was slightly higher in the first week, although not significantly so ($p > 0.05$). Combining the -2.5 SDS test in the first week with the 10% rule after the first week results in a sensitivity of 85.5% and a specificity of 99.4%; this is similar to the -2.5 SDS test for the first 2 weeks. This specificity is significantly higher ($p < 0.05$) than that for the -2 SDS rule.

Cases with a positive -2 or -2.5 SDS test had a significantly higher mean serum sodium concentration (163 mM) compared to cases with a negative -2 SDS test (149 mM) ($t = 2.6$, $df = 78$, $p = 0.01$) and with a negative -2.5 SDS test (151 mM) ($t = 3.0$, $df = 78$, $p = 0.004$). Of the cases with a positive -2.5 SDS test, 89% had a concentration of >149 mM, so the test detects the more severe cases of dehydration. Of the cases with a concentration of >149 mM, 91% had a positive -2.5 SDS test and 97% a positive -2 SDS test, and of the cases with a concentration of >159 mM, all cases had a positive -2.5 SDS test (and therefore a positive -2 SDS test).

Eight cases of hypernatraemic dehydration had a very small RWC. Three cases had a RWC between -2 SDS and -1 SDS and five cases had a weight change above -1 SDS.^{19 35 36 42} Clinical information was given in some studies; only mild and transient symptoms in these infants were reported. Serum sodium concentration was reported for six cases: four cases had a concentration below 149 mM and two above 149 mM (both 157 mM).

Figure 2 The -2.5 SDS relative weight change centile converted to weights by age for a given birth weight.

DISCUSSION

We developed a reference chart for breast-fed infants between postnatal days 2 and 11. This chart, together with cases of hypernatraemic dehydration obtained from the literature, was used to define an evidence-based referral rule. As far as we know, this is the first reference chart for RWC and the first evidence-based investigation of referral rules to detect infants with hypernatraemic dehydration. Our results show that a reference chart for RWC can be helpful to detect infants with hypernatraemic dehydration.

The RWC chart shows that the mean maximal weight loss occurs 3 days after birth and is 6% for a healthy, breast-fed infant. This is in agreement with several other studies which reported that breast-fed infants may lose up to 6%^{45–47} or 7%^{15 19 48 49} of their birth weight during the first week of life. The American Academy of Pediatrics and others also reported that normal weight loss reaches its peak at 3–5 days after birth.⁵⁰ Livingstone and the American Academy of Pediatrics Work Group on Breastfeeding suggested that a weight loss of greater than 7% of birth weight indicates possible breast feeding problems.^{19 50} Others suggested that a weight loss of 8% or more warrants further investigation.^{16–18}

Most authors reported that many midwives use the rule of thumb that infants may lose up to 10% of birth weight. Our results show that most infants with hypernatraemic dehydration have a weight loss of 10%. However, referral to a hospital of all infants with a weight loss of $>10\%$ would probably lead to many false positive results in the first week of life, assuming that for screening purposes the specificity needs to be sufficiently high. Therefore, we suggest applying the 0.6th centile (-2.5 SDS) as a criterion for referral to a hospital in the first week of life or using a weight loss of $>10\%$ after the first week of life. At the hospital, further diagnostic biochemical testing should be carried out. As it takes some time before insufficient breast feeding leads to weight loss, clinical differentiation between normal infants and those with hypernatraemic dehydration is not really possible in the first 2 days after birth. Infants with a weight loss of $>10\%$ (or <-2 SDS)

Table 3 Sensitivity, specificity and positive predictive value (PPV) for several referral rules for between postnatal days 3 and 11

Test	Sensitivity (%)	Specificity (%)	PPV (%)
10%	90.4	98.3	3.7
-2.5 SDS	85.5	99.4	9.2
-2 SDS	90.4	97.7	2.7
-2.5 SDS 3–6 days and 10% 7–11 days	85.5	99.4	9.2

Figure 3 Flowchart to detect dehydrated infants or infants at risk of dehydration. *Monitor the infant when mild clinical symptoms are present and refer the infant to hospital when severe clinical symptoms are present.

in the first week after day 2, should be monitored closely and require more intensive evaluation of breast feeding and possible intervention to correct problems with breast feeding. Furthermore, referral may also be warranted in infants with other clinical symptoms of dehydration even if weight loss is not particularly high. Clinicians should combine RWC values with examination of the infant, knowledge of feeding patterns, and number of wet diapers and frequency and quality of stools. We suggest using the flowchart in fig 3.

In addition to the 10% weight loss, another rule of thumb among midwives is that infants regain their birth weight by 10–14 days. The chart in this study shows that 50% of infants have regained their birth weight 8 days after birth, which is also consistent with other reports.^{6, 15} This study also shows that even after 11 days, about a third of infants have not yet regained their birth weight. We also expect that at day 14 a high percentage of infants will not have regained their birth weight. Therefore, we assume that this rule will lead to many false positive results. Macdonald *et al*¹⁵ suggested a revised intervention criterion: offer additional breast feeding support to those losing 10% of their birth weight but still consider this as normal and only consider weight loss above 12.5% or failure to regain birth weight by 21 days as being abnormal and requiring medical assessment. We applied the 12.5% weight loss rule to our data with infants from birth to 11 days old and found a sensitivity of 83.1% and a specificity of 99.9%. This rule has a better specificity (+0.5%) at the cost of a lower sensitivity (−2.4%) compared to the −2.5 SDS rule. With the 12.5% weight loss rule, 2.4% of the cases are missed. We think that a decrease in sensitivity of 2.4% is high and we therefore recommend using the proposed flow chart. However, one could consider using the 12.5% weight loss rule at day 3 as the −2.5 SDS line almost reaches 12.5% at day 3.

In our study we used information from cases with hypernatraemic dehydration reported in the literature. We expected that this information is biased towards the more severe cases of hypernatraemic dehydration, since severe cases are more likely to be reported than mild cases. Recently Moritz *et al*¹¹ found that only 17% of cases of hypernatraemic dehydration had non-metabolic complications. Therefore, the sensitivity and PPV in this study are likely to be lower for all infants with hypernatraemic dehydration. On the other hand, PPV may also be an underestimate as this value was based on a

minimum incidence rate of hypernatraemic dehydration. It would be very interesting in the future to test and possibly optimise our proposed referral rules using new cases with dehydration.

There is evidence that the degree of weight loss in babies born in a particular environment may be associated with the way that environment is managed.^{52, 53} In populations with “baby friendly” care, the prevalence of hypernatraemic dehydration may be lower than in populations with care that is less baby friendly. We assumed that the prevalence of hypernatraemic dehydration is 7.1 per 10 000 breast-fed infants. Based on this prevalence we calculated the PPV of several referral criteria. Since PPV is dependent on prevalence, in populations with a lower prevalence (perhaps due to baby friendly care) the PPV may be lower, whereas in populations with a higher prevalence the PPV of the same referral criteria will be higher.

We assumed that RWC expressed as a percentage is uncorrelated with birth weight. This means that a heavy child and a light child have the same distribution of RWC. However, this may not be true, as the degree, timing and variability of RWC may be quite different in small infants compared to large infants. We therefore tested the relationship between birth weight and RWC corrected for age using a linear mixed-effects model (residual variance = 1.53, AIC = 15 864). We found that an infant with a birth weight of 2.5 kg has on average a 1% greater RWC than an infant with a birth weight of 4.5 kg. As this is a relatively small difference for a large difference in birth weight, we decided to use the methodology unconditional on birth weight. The latter approach is also more convenient in practice than, for instance, various RWC curves for different categories of birth weight.

In this study, the weights of the infants were obtained in a research setting. The midwife was instructed to weigh the infant routinely. This means that the number of measurements should not depend on the status of the infant. To determine if this is indeed the case, using standard two-sample t tests we tested the dependence of the number of measurements and the status of the infants by testing the difference in RWC each day between the infants whose weight was being measured for the first time (besides their birth weight) and those who were being reweighed. We refitted the LMS method without the cases which were possibly reweighed because of a high RWC, and

What is already known on this topic?

- Several authors have proposed different rules of thumbs for identifying abnormal relative weight change (RWC).
- It is suggested that many midwives use the rule of thumb that infants may have a weight loss of 10% (= -10% RWC) but should regain their birth weight by 10–14 days.

What this study adds

- As far as we know, we have conducted the first evidence-based investigation of referral rules for infants with hypernatraemic dehydration and have developed the first reference chart for relative weight change.

found that the difference between the median RWC in the newly constructed growth chart and the reference chart based on all infants was negligible ($\leq -0.2\%$).

We conclude that the rule of thumb that infants may have a relative weight loss of 10% is excellent after the first week of life. However, in the first week of life this rule will produce too many false positive results. A chart for RWC can be helpful to detect infants with hypernatraemic dehydration.

ACKNOWLEDGEMENTS

We are grateful to Mieke Huisenga, Wendelien Roepke, Marleen Snablie and the midwife practices for obtaining the reference group data. We thank Caren Lanting and Jan Maarten Wit for their suggestions for improvement.

Authors' affiliations

Paula van Dommelen, Jacobus P van Wouwe, Jacqueline M Breuning-Boers, Stef van Buuren, Paul H Verkerk, TNO Quality of Life, Leiden, the Netherlands

Competing interests: None declared.

REFERENCES

- 1 World Health Organization. *The optimal duration of exclusive breastfeeding. Report of an expert consultation*. Geneva: World Health Organization, 2001.
- 2 Lawrence RA, Howard CR. Given the benefits of breastfeeding, are there any contraindications? *Clin Perinatol* 1999;26:479–90, viii.
- 3 Lanting CI, Van Wouwe JP, Reijneveld SA. Infant milk feeding practices in the Netherlands and associated factors. *Acta Paediatr* 2005;94:935–42.
- 4 Naylor AJ. Baby-Friendly Hospital Initiative. Protecting, promoting, and supporting breastfeeding in the twenty-first century. *Pediatr Clin North Am* 2001;48:475–83.
- 5 Laing IA, Wong CM. Hypernatraemia in the first few days: is the incidence rising? *Arch Dis Child Fetal Neonatal Ed* 2002;87:F158–62.
- 6 Oddie S, Richmond S, Coulthard M. Hypernatraemic dehydration and breast feeding: a population study. *Arch Dis Child* 2001;85:318–20.
- 7 Clarke TA, Markarian M, Griswold W, et al. Hypernatremic dehydration resulting from inadequate breast-feeding. *Pediatrics* 1979;63:931–2.
- 8 Arboit JM, Gildengers E. Breast-feeding and hypernatremia. *J Pediatr* 1980;97:335–6.
- 9 Rowland TW, Zori RT, Laffleur WR, et al. Malnutrition and hypernatremic dehydration in breast-fed infants. *JAMA* 1982;247:1016–17.
- 10 Clarke AJ, Sibert JR. Hypernatraemic dehydration and necrotizing enterocolitis. *Postgrad Med J* 1985;61:65–6.
- 11 Cooper WO, Atherton HD, Kahana M, et al. Increased incidence of severe breastfeeding malnutrition and hypernatremia in a metropolitan area. *Pediatrics* 1995;96:957–60.
- 12 Kaplan JA, Siegler RW, Schmunk GA. Fatal hypernatremic dehydration in exclusively breast-fed newborn infants due to maternal lactation failure. *Am J Forensic Med Pathol* 1998;19:19–22.
- 13 Chaudhary R, Twaddle S, Levi R, et al. Hypernatremic dehydration in breast-fed infants. *Paediatrics Today* 1998;6:85–7.
- 14 Ng PC, Chan HB, Fok TF, et al. Early onset of hypernatraemic dehydration and fever in exclusively breast-fed infants. *J Paediatr Child Health* 1999;25:585–7.
- 15 Macdonald PD, Ross SR, Grant L, et al. Neonatal weight loss in breast and formula fed infants. *Arch Dis Child Fetal Neonatal Ed* 2003;88:F472–6.
- 16 Neifert MR. Clinical aspects of lactation. Promoting breastfeeding success. *Clin Perinatol* 1999;26:281–306, v–vi.
- 17 Neifert MR. Prevention of breastfeeding tragedies. *Pediatr Clin North Am* 2001;48:273–97.
- 18 Simoes EA, Pereira SM. The growth of exclusively breastfed infants. *Ann Trop Paediatr* 1986;6:17–21.
- 19 Livingstone VH, Willis CE, Abdel-Wareth LO, et al. Neonatal hypernatremic dehydration associated with breast-feeding malnutrition: a retrospective survey. *CMAJ* 2000;162:647–52.
- 20 DeMarzo S, Seacat J, Neifert M. Initial weight loss and return to birth weight criteria for breast-fed infants: challenging the 'rule of thumb'. *Am J Dis Children* 1991;145:402.
- 21 Bajpai A. Hypernatremic dehydration in a neonate. *Indian Pediatr* 2002;39:599–600.
- 22 Becker PG, Conard JA. Breast-feeding failure. *Indiana Med* 1990;83:648–50.
- 23 Boumahni B, Pyaraly S, Randrianaly H, et al. Hypernatremic dehydration and breastfeeding. *Arch Pediatr* 2001;8:731–3.
- 24 Canet E, Canet J, Perrain MF. Quel est votre diagnostic? *Concours Med* 1984;106:1689–91.
- 25 Chilton LA. Prevention and management of hypernatremic dehydration in breast-fed infants. *West J Med* 1995;163:74–6.
- 26 Gebbara BM, Everett KO. Dural sinus thrombosis complicating hypernatremic dehydration in a breastfed neonate. *Clin Pediatr (Phila)* 2001;40:45–8.
- 27 Harding D, Moxham J, Cairns P. Weighing alone will not prevent hypernatraemic dehydration. *Arch Dis Child Fetal Neonatal Ed* 2003;88:F349.
- 28 Hilliard TN, Marsh MJ, Malcolm P, et al. Radiological case of the month. Sagittal sinus thrombosis in hypernatremic dehydration. *Arch Pediatr Adolesc Med* 1998;152:1147.
- 29 Jaramillo I, Lopez G, Hernandez H. Hypernatremic dehydration and death in an infant. *Pediatr Emerg Care* 2003;19:62–3.
- 30 Kini N, Zahn S, Werlin SL. Hypernatremic dehydration in breast-fed infants. *Wis Med J* 1995;94:143–5.
- 31 Macdonald PD, Grant L, Ross SR. Hypernatraemia in the first few days: a tragic case. *Arch Dis Child Fetal Neonatal Ed* 2003;88:F350.
- 32 Marino R, Gourji S, Rosenfeld W. Neonatal metabolic casebook. Hypernatremia and breast feeding. *J Perinatol* 1989;9:451–3.
- 33 Mercier JC, Outin S, Paradis K, et al. [Breast feeding and hypernatremic dehydration. 3 case studies]. *Arch Fr Pediatr* 1986;43:465–70.
- 34 Molteni KH. Initial management of hypernatremic dehydration in the breastfed infant. *Clin Pediatr (Phila)* 1994;33:731–40.
- 35 Niestijl AL, Sauer PJ. Breast feeding during the first few days after birth: sometimes insufficient. *Ned Tijdschr Geneesk* 2003;147:2405–7.
- 36 Pascale JA, Brittan L, Lenfestey CC, et al. Breastfeeding, dehydration, and shorter maternity stays. *Neonatal Netw* 1996;15:37–43.
- 37 Rand SE, Kolberg A. Neonatal hypernatremic dehydration secondary to lactation failure. *J Am Board Fam Pract* 2001;14:155–8.
- 38 Roddey OF, Martin ES, Swetenburg RL. Critical weight loss and malnutrition in breast-fed infants. *Am J Dis Child* 1981;135:597–9.
- 39 Rosenfeld W, deRomana GL, Kleinman R, et al. Improving the clinical management of hypernatremic dehydration. Observations from a study of 67 infants with this disorder. *Clin Pediatr (Phila)* 1977;16:411–17.
- 40 Sofer S, Ben Ezer D, Dagan R. Early severe dehydration in young breast-fed newborn infants. *Isr J Med Sci* 1993;29:85–9.
- 41 van der Heide PA, Toet MC, van Diemen-Steenvoorde JA, et al. Hypertonic dehydration in 'silent' malnutrition of breast-fed infants. *Ned Tijdschr Geneesk* 1998;142:993–5.
- 42 Willis CE, Livingstone V. Infant insufficient milk syndrome associated with maternal postpartum hemorrhage. *J Hum Lact* 1995;11:123–6.
- 43 Cole TJ, Green PJ. Smoothing reference centile curves: the LMS method and penalized likelihood. *Stat Med* 1992;11:1305–19.
- 44 van Buuren S, Fredriks M. Worm plot: a simple diagnostic device for modelling growth reference curves. *Stat Med* 2001;20:1259–77.
- 45 Maisels MJ, Gifford K. Breast-feeding, weight loss and jaundice. *J Pediatr* 1983;102:117–18.
- 46 Marchini G, Fried G, Ostlund E, et al. Plasma leptin in infants: relations to birth weight and weight loss. *Pediatrics* 1998;101:429–32.
- 47 Marchini G, Stock S. Thirst and vasopressin secretion counteract dehydration in newborn infants. *J Pediatr* 1997;130:736–9.
- 48 Maisels MJ, Gifford K, Antle CE, et al. Jaundice in the healthy newborn infant: a new approach to an old problem. *Pediatrics* 1988;81:505–11.
- 49 Thureen PJ, Hay WW Jr. Advice for selected breast-feeding issues. *Compr Ther* 1996;22:802–5.
- 50 American Academy of Pediatrics Section on Breastfeeding. Breastfeeding and the use of human milk. *Pediatrics* 2005;115:496–506.
- 51 Moritz ML, Manole MD, Bogen DL, et al. Breastfeeding-associated hypernatremia: are we missing the diagnosis? *Pediatrics* 2005;116:343–7.
- 52 Mikiel-Kostyra K, Mazur J. Hospital policies and their influence on newborn body weight. *Acta Paediatr* 1999;88:72–5.
- 53 Avoa A, Fischer PR. The influence of perinatal instruction about breast-feeding on neonatal weight loss. *Pediatrics* 1990;86:313–15.