patients who received lactose, 4 of the 26 patients who received sulphatriad, and 1 of the 26 who received sulphanilamide developed complications. There is obviously little difference between the effects of the treatments in this respect.

Discussion

We do not propose to enlarge upon the epidemiological aspects in the present report. This study must be regarded as a "pilot" one undertaken in an endeavour to prospect the possibilities of such a form of investigation. Nevertheless it is interesting that our results show certain similarities to those obtained by the Commission on Respiratory Diseases. It must be remembered that their clinical material was all adult and confined to cases in which the temperature exceeded 100° F. (37.8° C.) and which thus qualified for admission to hospital. From the results of throat swabs and the demonstration of a rise in serum antibodies they concluded that roughly half of their cases were streptococcal. They also found some dissimilarity between the clinical pattern of the streptococcal and non-streptococcal cases in that chills, headache, and malaise were commoner in the former, whereas cough, nasal obstruction, and hoarseness were commoner in the latter.

As part of the American study the clinicians were asked to define the probable bacteriological cause before receiving the result of the throat swab, and it was found that the non-streptococcal cases were fairly accurately classified. There were, however, numerous errors among the cases finally proved to be streptococcal. Our own experience was that errors were frequent in both groups. Thus each of us saw patients in whom the clinical diagnosis of streptococcal tonsillitis appeared "obvious," but from whom a normal bacteriological flora was obtained from two throat swabs and a post-nasal swab; and others in whom a mildly inflamed throat produced almost a pure culture of haemolytic streptococci. The bacteriological examination of all such cases in general practice is not a practicable proposition; and yet precise differentiation of the cases without it is really impossible. Despite our observations, which suggest that a differentiation can be made between the different groups comprising the series, there seems to be no single clinical criterion which can be applied to diagnose "streptococcal" tonsillitis. In such circumstances it is obvious that the term must be used with considerable caution.

The results of the therapeutic trial do not support the thesis that sulphonamides exert a beneficial effect in the treatment of the "acute throat." Those who place more faith in the virtue of clinical impressions will no doubt be unconvinced by an array of "statistics." But it is well to remember that the average duration of fever and pain in those cases treated with lactose was exceedingly short—namely, 1.07 and 1.95 days respectively. In view of this, added to the fact that in about half of the cases a streptococcal aetiology is unlikely, it seems that to use drugs which have definite and known dangers is unnecessary, besides being expensive.

We are indebted to Dr. W. R. Thrower, of May and Baker Ltd., for supplying us with tablets of uniform size and appearance. Professor A. Bradford Hill and his colleagues undertook the statistical analysis and provided the random numbers, and to them we express our thanks.

REFERENCES

- Anderson, T. (1949). *British Medical Journal*, 2, 860.
Commission on Acute Respiratory Diseases (1944). *J. Amer. med. Ass.*, 125, 1163.

LEAD-POISONING IN CHILDREN

REPORT OF FIVE CASES, WITH SPECIAL REFERENCE TO PICA

BY

N. F. ELLIOTT BURROWS, M.A., B.M., D.C.H.
Senior Resident Medical Officer, Belgrave Hospital for Children

**JOHN RENDLE-SHORT, M.A., M.B., M.R.C.P.
D.C.H.**

Late House-physician, Hospital for Sick Children, Great Ormond Street

AND

**DENIS HANNA, M.B., M.R.C.P., M.R.A.C.P.
D.C.H.**

Late House-physician, Hospital for Sick Children, Great Ormond Street

Lead-poisoning in infancy and childhood is rare in Great Britain, although in other countries, notably Australia, America, and Japan, many cases have been described.

In a survey of the literature of this country for the past 60 years we have found only seven other reports of plumbism in infants and children. Stephenson (1898) reported poisoning in an infant from the use of white lead as toilet powder, and Cadman (1902) and Findlay (1935) each described a case due to lead nipple-shields. Rodgers, Peck, and Jupp (1934) described the case of a boy aged 2 years who had chewed toy soldiers, and Gartside (1948) reported two cases in a family from burning battery casings as fuel. Clark (1950) has reported a case in a breast-fed infant due to the lead contained in the family water supply.

We have recently seen three cases, and two others have been found in the records of the Hospital for Sick Children, Great Ormond Street. The five cases are reported below.

Sources of Lead

The source of the lead is broadly determined by the age of the patient. (1) In infancy there are dangers such as the application of powder containing lead to the breasts or to the infant's skin, and the use of lead nipple-shields. The use of lead powder by mothers was found to be the common cause of lead encephalopathy in Japanese infants. (2) In children up to about 5 years the ingestion of lead paint or metallic lead due to perverted appetite is the most likely cause, although peculiar circumstances have resulted in large outbreaks in Queensland, Australia. Climatic conditions there favoured the powdering of paint, which was readily transferred to the children's mouths (B.M.A. Queensland Branch Report, 1922). (3) In children of school age perverted appetite is not often a factor, but they may encounter a great diversity of hazards, as in adults. The majority of 40 cases due to burning battery casings as fuel were in older children (Williams *et al.*, 1933).

Pica.—Four of the five cases here described exhibited pica, or perverted appetite. All children pass through a temporary phase of putting objects in their mouths, but pica refers to a morbid craving, of longer duration, for such undesirable articles as earth, sand, coal, wall-paper, cloth, hair, paint, and even wood-lice. The condition may result from gastro-intestinal disturbances, but mental deficiency, neurosis, and simple habit are often

responsible. Ruddock (1924) was probably the first to point out the relationship of lead-poisoning to pica when he stated that some children affected with the former possessed a morbid craving to gnaw painted objects such as window-sills, white furniture, crib railings, porch railings, and other articles. To this list may be added toy soldiers, farm animals, etc. McKhann (1926) found pica to be the commonest forerunner of plumbism, and the condition has since been stressed by Stetson (1947). The case reported by Rodgers *et al.* (1934) in this country also suffered from pica.

Clinical Features

The manifestations of chronic lead-poisoning in children differ from those in adults in several important respects. In adults the main features are gastro-intestinal symptoms—chiefly colic and constipation—anaemia, lead line on the gums, and peripheral palsy involving the muscles most used. In children the emphasis is on encephalopathy, which occurs in a high percentage of cases; eye signs are common, while the lead line and muscle palsies are rare.

The manifestations of encephalopathy are probably secondary to an intense cerebral oedema, accompanied by a raised intracranial tension, which in young patients may produce a bulging fontanelle, separation of the sutures, and a cracked-pot note on percussing the skull.

The clinical features in children have been well described by Blackfan (1917), McKhann (1932), and Ford (1945).

Laboratory and Other Investigations

The blood shows a moderate anaemia, reticulocytosis, punctate basophilia, and an excess of normoblasts; the white count is normal, although a leucocytosis may be present in acute poisoning or with superadded infection. In severe cases the serum bilirubin may be elevated owing to excessive haemolysis.

The urine commonly shows a trace of protein, and glycosuria is often present; ketone bodies may also be found.

The cerebrospinal fluid is often abnormal (Weller and Christensen, 1925). In the presence of encephalopathy the pressure is almost always high; the protein is increased and there is a moderate pleocytosis. The sugar may also be raised, although this is not pathognomonic, as it may occur in convulsions due to other causes.

Lead estimations in the blood and urine may give valuable confirmatory evidence. Special precautions must be taken to ensure that the syringes and containers are lead-free.

Radiography is of the greatest value, as changes are practically always present (Park *et al.*, 1931; Caffey, 1931; Vogt, 1932). Radio-opaque bands, due to the deposition of lead in abnormally numerous and condensed trabeculae, are seen at the growing ends of the long bones and the growing margins of flat bones. Multiple bands or lines are sometimes seen, but their presence does not necessarily indicate intermittent absorption of lead. If the deposition of lead ceases, an area of bone of normal density appears on the epiphysal side of the band, and as the bone grows this band becomes wider and less well defined. Since lead is deposited only in actively growing bone, this finding is not present in adults.

Straight x-ray films of the abdomen may reveal specks of radio-opaque material due to the presence of metallic lead or lead salts in the gut.

In the presence of high intracranial tension a radiograph of the skull may show separation of the sutures.

Differential Diagnosis

In children this can be considered under two heads—before and after encephalopathy has developed.

Before the cerebral signs appear, vomiting, abdominal pain, and constipation are the principal symptoms, and, if associated with pica, lead-poisoning should always be excluded. This triad of symptoms may suggest a surgical emergency. The label "cyclical vomiting" is always dangerous, as it tends to lull the physician into a belief that an adequate diagnosis has been made.

After encephalopathy has developed, a number of conditions may be simulated. (1) Diabetes mellitus, suggested by abdominal pain and vomiting, drowsiness, or coma associated with sugar and acetone in the urine. In lead-poisoning, however, urine is not passed in excess and the blood sugar is normal. (2) An intracranial space-occupying lesion, especially the common medulloblastoma of the cerebellum. Localizing signs may be absent, and papilloedema may be found in both conditions. However, convulsions and twitching are unusual with sub-tentorial tumours, and very frequent in lead encephalopathy. The cerebrospinal fluid may be similar in both. Good (1941) has recommended radiography of the long bones to demonstrate the presence or absence of lead lines in every case of raised intracranial pressure if choked disks are present. (3) Tuberculous meningitis should always be excluded in any child with doubtful cerebral signs. (4) "Encephalitis" is often an unsatisfactory diagnosis, and it is possible that some cases so labelled may be due to plumbism.

Treatment

After absorption, lead is deposited in the bones and, to a lesser extent, in the organs, especially the liver and brain. It is excreted by the kidneys, which appear to have a definite renal threshold for it. Lead deposited in the bones is innocuous. There are therefore two lines of treatment available: the lead may be immobilized in the bones and thereby rendered harmless, or mobilized to increase its excretion by the kidneys.

There is a close relationship between calcium metabolism and the behaviour of lead in the body. Calcium has a specific action in aiding lead deposition in the bones, and should be given in large quantities, as milk (3 pints (1.7 litres) daily) and calcium lactate by mouth. Vitamin D also exercises a beneficial effect by fixing lead phosphate in the bones. Calcium gluconate intravenously (10 ml. of 10% solution) should be used in crises such as convulsions, coma, and severe colic, when the acute symptoms are often dramatically relieved.

"De-leading"—the removal of lead from the bones and its elimination from the body—is very dangerous in children owing to the risk of encephalopathy. Most authorities think that it should never be undertaken. Both acidosis and alkalosis aid the removal of lead from the bones, and so must be avoided.

The cerebral oedema of lead encephalopathy is very resistant to treatment. The following methods have been advised: hypertonic magnesium sulphate per rectum or intramuscularly, hypertonic glucose intra-

venously, repeated lumbar punctures, and even surgical decompression of the brain. The rapid withdrawal of cerebrospinal fluid by the lumbar route is not without danger, as medullary compression, fits, or coma may be precipitated.

Convulsions must be controlled by phenobarbitone or paraldehyde intramuscularly.

Dimercaprol (B.A.L.) has been used with success in the treatment of persons poisoned by various metals, notably mercury and gold, but the evidence of its benefit in human cases of lead-poisoning is largely experimental and controversial.

Braun *et al.* (1946), experimenting on animals, found an increased mortality in a group of rabbits treated with dimercaprol compared with an untreated control group. Their findings were confirmed by Germuth and Eagle (1948). Weatherall (1948), however, working with mice, obtained opposite results.

In a group of men exposed to the hazard of industrial lead, Ryder *et al.* (1947) found that dimercaprol brought about a rapid fall in the concentration of lead in the blood and a considerable increase in urinary excretion of lead.

Telfer (1947) reported a case of chronic lead-poisoning treated with dimercaprol in which an increased urinary excretion of lead occurred with beneficial results. R. H. S. Thompson (1950—personal communication) considers dimercaprol is worthy of trial and suggests a dosage of 1.5 mg. per kg. four times a day for one week. This should be doubled in cases of lead encephalopathy.

Prognosis

Lead-poisoning in children carries a grave prognosis. The younger the child the higher the mortality. Encephalopathy is of particularly grave significance. Focal or generalized sequelae involving the central nervous system are common. McKhann and Vogt (1933) record that of 45 cases of encephalopathy 11 died and 12 had permanent sequelae—epilepsy, tremors, mental retardation, blindness, and speech defect; and Byers and Lord (1943) state that mental changes leading to unsatisfactory progress at school were present in 19 out of 20 children who had previously been normal. Pneumoencephalography revealed cortical atrophy in six patients who suffered permanent cerebral damage (Levinson and Zeldes, 1939).

Case 1

A boy aged 7 was first seen in September, 1948, at the Hospital for Sick Children, Great Ormond Street, with a three-years history of repeated attacks of vomiting, abdominal pain, and constipation lasting from one day to three weeks. He was of normal intelligence. A provisional diagnosis of periodic syndrome (cyclical vomiting) was made, and he was admitted to hospital. A blood count showed a mild anaemia and punctate basophilia. A radiograph of the abdomen revealed specks of opaque material, but in a subsequent film these had disappeared. It was now discovered that he suffered from pica and would eat anything. He particularly enjoyed chewing lead soldiers. The diagnosis was changed to " ? plumbism." He was seen in the out-patient department after discharge, when his mother stated that he had eaten no more lead soldiers. A blood count was normal and the attacks were therefore again considered to be due to periodic syndrome.

In August, 1949, he developed slight jaundice, and the vomiting, abdominal pain, and colic became much worse. He was readmitted in November, 1949.

On examination he was seen to be a quiet, co-operative, thin child, with dry skin and icterus of the conjunctivae.

Pallor and blue line on the gums were not present. The abdomen was scaphoid, not tender. He had slight tremor of the hands, otherwise no abnormality was noted in the central nervous system. The fundi were within normal limits. For the first six days he continued to vomit, and was very quiet. He was constipated, complained of abdominal pain, and would not eat.

Investigations.—A blood count showed: Hb, 9.4 g. % (67%); red cells, 3,600,000 per c.mm.; punctate basophilia, 1.3%; serum bilirubin, 1.5 mg. per 100 ml.; fasting blood sugar, 125 mg. per 100 ml. Examination of the urine revealed: acid, acetone —, faint trace of protein, orange reduction with Benedict's reagent. Liver-function tests were normal. A radiograph of the abdomen again revealed specks, presumably of metal in the gut. Ring shadows were clearly visible parallel to the iliac crests (Fig. 1). Radiographs of the long bones showed bands of increased density near the epiphyses (Fig. 2).

FIG. 1.—Case 1. Radiograph showing metal particles in the gut and ring shadows parallel to the iliac crests.

FIG. 2.—Case 1. Radiograph of long bones showing bands of increased density near the epiphyses.

Progress and Treatment.—The child was started on a high-calcium diet, vitamin D, and alkalis. Lumbar puncture on the sixth day gave a pressure of 100 mm., and the fluid contained 1 cell per c.mm. and 70 mg. of protein and 93 mg. of sugar per 100 ml. About two hours later he lapsed into unconsciousness and remained semicomatose for three days. Vigorous treatment for the encephalopathy was carried out. He was given magnesium sulphate by mouth, 50% glucose intravenously, and 25% magnesium sulphate intramuscularly. On the seventh day the blood pressure was 165/120 and the cerebrospinal-fluid pressure had risen to 190 mm. The child started twitching all over, but much more on the left side than on the right. One major convulsion occurred. Two days after the return of consciousness he developed wrist-drop on the left side which lasted one week. He remained well and his blood pressure slowly fell to 105/65. Ten days later he suddenly lapsed into coma again, with twitching on the left side of his body. The pulse was full, there was marked vasodilatation, and the blood pressure was 115/10. This second attack of encephalopathy lasted one day only. Clinically he made an uneventful recovery and was discharged to his home.

Special Reports.—**Electroencephalogram** (Dr. W. A. Cobb, National Hospital, Queen Square): "Very little normal activity can be made out. All areas show moderate-voltage slow rhythms, maximal on the right side. No change on over-breathing. A grossly abnormal record." **Psychological report** (Dr. Mildred Creak and Miss Durward): "Result indicates low average intelligence. Considerable difficulty with all items involving visual-motor co-ordination. Chronological age 7 years 5 months, mental age 6 years 10 months. Intelligence quotient, 92." **Lead estimations:** Blood lead, 0.3 µg. per 100 ml. (normal 0.01–0.03 µg.); urine lead, 0.285 mg. per 24 hours (normal not more than 0.2 mg. per 24 hours).

Subsequent History.—After being home a few days the child started to vomit and to have "vacant attacks," during which his head and eyes deviated to the left. After one of these a left facial paralysis was noticed. He was readmitted, and the attacks were controlled with intravenous calcium gluconate and phenobarbitone by mouth. Later, a second psychological report states: "There is mental slowness, slight slurring of speech, and spurts of aggression. He behaves like someone who is very slightly drunk."

The report on a further E.E.G. stated: "In the right fronto-parietal region there is a focus of very slow, rather rhythmic waves of quite high amplitude. They frequently give a hint of a repetitive complex-form analogous to slow spike-and-wave complexes, though this is never clearly defined."

Case 2

A girl aged 2½ was brought to the out-patient department of the Belgrave Hospital for Children in September, 1949, on account of vomiting and loss of weight. According to the mother, the child's condition had slowly worsened during the previous 12 months. She had got paler, had lost weight, and had seemed increasingly miserable—so much so that her whole personality seemed to have altered. Her condition had markedly deteriorated in the previous two months; she had become anorexic and constipated, and had vomited on many occasions. It was learnt that she had insatiable pica, and that she chewed almost anything.

On examination she was found to be of normal height for her age, although about 5 to 6 lb. (2.3 to 2.7 kg.) under weight. She was anaemic, markedly hypotonic, and extremely miserable, screaming at the slightest interference. No other abnormalities were found on physical examination.

Investigations.—Mantoux-negative at 1:10,000. A blood count on admission showed: Hb, 40%; red cells, 2,860,000; marked anisocytosis and polychromasia; punctate basophilia, 2.5%; normoblasts, 0.5%; white cells, 23,700; polymorphonuclear cells, 56%. Six weeks later it was: Hb, 84%; red cells, 4,700,000; white cells, 9,300. The cerebrospinal fluid 12 days after admission showed: clear colour-

less fluid with pressure varying between 120 and 160 mm.; cells, 2 per c.mm.; protein, 25 mg. per 100 ml.; chlorides, 740 mg. per 100 ml.; sugar normal.

Progress and Further Investigations.—The child remained pathologically miserable in the ward, and would scream whenever anyone looked at her or went near her. As the first blood report mentioned punctate basophilia, it was decided to take radiographs of the long bones: the report from the radiologist (Dr. A. M. Rackow) stated that the radiological appearances were in favour of a heavy-metal poisoning. As the most likely heavy metal was lead, a search of the house where the child lived was undertaken, and it was noted that the skirting-boards and window frames had been gnawed down to the bare wood, while the painted handle of a doll's "pram" had been bitten down to the metal. On the boards and window-frames there were apparently two layers of paint—a brown one superficially and a blue one deeper. Samples of the paints were taken from these sources and from the pram handle. The following results were obtained: brown paint, no lead; deeper skirting-board and window-frame paint, 27% lead; pram-handle paint, 7% lead. Blood-lead estimation showed 0.08 µg. per 100 ml.

E.E.G. Report (Dr. W. A. Cobb).—"The E.E.G. was largely unsuccessful owing to excessive movement. So far as can be judged, however, there is no abnormality."

The child's haemoglobin started to rise on "colliron," and her behaviour improved slightly. She was discharged in December, 1949, and a follow-up revealed that she had stopped chewing the lead paintwork but was still very miserable.

Case 3

A baby girl aged 1 year 7 months, weighing 19 lb. 5 oz. (8.86 kg.), was admitted to the Hospital for Sick Children, Great Ormond Street, in December, 1949. She had been irritable, listless, and crying frequently since an attack of measles seven months previously. Vomiting had been present for six weeks and had increased recently. Constipation had been marked for three weeks. Her appetite was poor and she was steadily losing weight. During the night she had had many screaming attacks, which were relieved by nursing her in front of an open fire.

On examination she was seen to be pale, apathetic, and moderately wasted, with a feeble grizzly cry. Her throat, ears, lungs, heart, abdomen, fundi, and cranial nerves were normal. Normal movements were present in the limbs, but hypotonia was marked. Tendon reflexes were absent in her arms and ankles and diminished in her knees. Plantar responses were flexor. Her skull was not enlarged; the fontanelle was closed. Attempts to obtain a blood-pressure reading were unsuccessful.

A blood count revealed a hypochromic anaemia with a moderate punctate basophilia. The presence of the latter suggested lead-poisoning. The parents were again interviewed, and it was found that for about eight months the child had chewed anything she could lay her hands on—including painted woodwork. With this further evidence in favour of lead-poisoning, the long bones and abdomen were x-rayed and a blood-lead estimation was performed. Radiographs of the wrists and knees revealed a band of increased density at the ends of the femora, tibiae, radii, and some of the metacarpals. Radiologically the abdomen was normal.

The child's condition deteriorated. She continued to vomit, became drowsy and dehydrated, and developed a marked bilateral internal strabismus. There was no lead line on the gums or evidence of wrist-drop or foot-drop. After lumbar puncture (pressure in excess of 300 mg. of water, the only abnormality being 60 mg. of protein) she had a series of convulsions and died 18 hours later.

Necropsy (Dr. Vickers).—**Brain:** Slight cellular infiltration of pia arachnoid with perivascular oedema. No evidence of encephalitis or vascular lesion. **Liver:** Fatty change throughout with many hepatic-cell nuclei showing fine

eosinophilic acid-fast inclusions of the type found in lead-poisoning. *Kidney*: The most pronounced degenerative changes in the cortex were acidophilic; acid-fast nuclear inclusions were present in greatest numbers. *Oesophagus*: Just proximal to the cardiac orifice of the stomach the oesophagus showed a perforation 2.5 cm. in length, the lesion having the characteristics of an acute oesophageal ulcer (Wyatt and Khoo, 1949). The stomach and intestines contained altered blood in large quantity. No other ulceration was present in the gastro-intestinal tract. *Lead estimations*: Blood lead less than 0.03 μ g. per 100 ml.; bone lead, 89 parts per million; liver lead, 53 parts per million.

Case 4

The patient, a breast-fed male baby aged 5½ months, had previously been quite healthy. For 10 days before admission he had been irritable, with vomiting and loose stools. Five days before admission he had had a convulsion. He was admitted to another hospital and then transferred to the Hospital for Sick Children, Great Ormond Street.

On examination he was pale, limp, and irritable, with slight meningismus. An otitis media and otitis externa were present. A blood count showed: red cells, 3,300,000 per c.mm.; Hb, 65%; reticulocytes, 8.4%; punctate basophilia present; fragility range within normal limits; white cells, 23,200 per c.mm. (polymorphs 40%, lymphocytes 52%, monocytes 8%). The cerebrospinal fluid showed: pressure normal; cells, 3 per c.mm.; protein, 224 mg. per 100 ml.; sugar, 59 mg. per 100 ml.; chlorides, 720 mg. per 100 ml.

It was now learnt that the mother had been using a lead nipple-shield for the past 10 weeks. A radiograph of the

FIG. 3.—Case 4. Radiograph showing opaque lead lines 1 in. (2.5 cm.) from the epiphysis and widening of the bone at this point.

long bones revealed typical bands of increased density at the ends of the shafts. The urinary excretion of lead in 24 hours was 0.03 mg.

The child was given a blood transfusion for the anaemia and penicillin for the otitis. He became livelier, gained weight, and was sent home. Later he had to be readmitted for a discharging ear and diarrhoea and vomiting. Intravenous therapy was required to counteract

the dehydration. Further radiographs four months later showed that the bands were then situated about 1 in. (2.5 cm.) from the epiphysis, and had caused a widening of the bone at this point (Fig. 3).

Follow-up.—When seen at the age of 3 the patient was a normal healthy child. The blood picture was normal, but radiographs still showed very faint lines of increased density in the shafts.

Case 5

A boy aged 5 was admitted to the Hospital for Sick Children, Great Ormond Street, in August, 1931. He had been unwell for the past three weeks, with vomiting, pains in the back of his head, pallor, and constipation following an initial bout of diarrhoea. For an indefinite period he had been nibbling paint off the windows at home.

On examination he was thin and pale, and complained of abdominal pain. No abnormal physical signs were found apart from hepatic enlargement. A blood count showed: red cells, 3,375,000 per c.mm.; Hb, 58%; white cells, 4,700 per c.mm.; 6 normoblasts per 100 white cells. Marked basophilic stippling was present. Examination of the urine revealed no albumin or deposit; a trace of lead was present. Analysis of a sample of paint similar to that ingested showed

lead to be present. Radiographs of the long bones were not taken.

The child was discharged from hospital, but we have not been able to follow him up.

Discussion

Several interesting points emerge from the study of these cases. In Case 1 the temporary wrist-drop was considered to be Todd's paralysis rather than a true lead palsy. This is supported by the predominantly unilateral nature of the fits and the electroencephalogram showing maximal changes from the corresponding hemisphere.

Encephalography was carried out in Cases 1 and 2. It is hoped that the E.E.G. may prove to be of value in determining the prognosis in lead encephalopathy. It is likely that abnormal E.E.G. changes will be present in those subsequently showing mental retardation.

Glycosuria was observed in two of our cases. Goettsch and Mason (1940) reported glycosuria in 8 out of 39 cases. They discussed the possibility of its being pancreatic or cerebral in origin associated with hyperglycaemia, or renal in origin associated with a normal or low blood sugar. The evidence points to the latter view.

Case 3 showed typical necropsy findings of lead-poisoning. The cause of death was haemorrhage from acute ulceration of the oesophagus. Although ulceration of the alimentary tract may occur in acute lead-poisoning, it is not a feature of chronic lead-poisoning. However, it has been described in association with cerebral oedema.

Blood-lead estimations were performed in three cases: the level was abnormally high in two, and within normal limits in one. Urinary lead estimations were also done in two cases.

Present-day manufacturers of nursery equipment in this country are fortunately aware of the danger of lead paint to the child. Modern paints seldom contain lead; but old paint, with its high lead content, constitutes a great danger. In two of our cases old paint appeared to be the source of the lead.

We have described a further case of poisoning due to a lead nipple-shield. Mackay and Hunter (1949) have strongly condemned the use of lead nipple-shields.

Summary and Conclusions

Five cases of lead-poisoning in children are presented. Four were associated with pica and one was caused by the use of a lead nipple-shield.

The serious prognosis of the condition is stressed. One of our patients died and two appear to be mentally backward.

The sources of lead in such cases can be broadly divided according to the ages of the patients. (1) In infancy: due to lead nipple-shields or the use of lead-containing powders, etc.; (2) 1–5 years: due to children with pica eating lead along with other noxious substances; most commonly this is in the form of lead paint; and (3) school-children: due to accidents similar to those met with in adult life.

The clinical features of lead-poisoning are discussed. The manifestations in childhood differ from those in adults in that encephalopathy, with irritability, fits, and coma, is very common. Wrist-drop and the lead line on the gums, however, are rare.

The diagnosis is often first suspected from the finding of punctate basophilia in a blood film. Although not diagnostic, punctate basophilia should lead to further

investigations, of which x-ray examination of the long bones is the most important. The possible value of E.E.G. recordings is discussed.

Treatment should aim at immobilizing the lead in the bones. "De-leading" should never be attempted in children.

Lead-poisoning in children could be entirely prevented. Lead paint should never be used in places that children can reach. Lead nipple-shields should be prohibited.

The use of dimercaprol in the treatment of lead-poisoning is discussed and is thought to be worthy of trial, especially in cases of lead encephalopathy.

Lastly, it is stressed that the diagnosis of pica should always bring with it the question, "Is this child liable to ingest poisons by its habit of dirt-eating?" Probably the most important of such poisons is lead.

We gratefully acknowledge the help of Drs. Doyne Bell, R. Lightwood, D. Paterson, B. E. Schlesinger, and W. G. Wyllie, under whom these children were admitted. We would also like to thank Dr. Mildred Creak and Miss Durward for the psychological reports on Case 1; Dr. W. A. Cobb for the electroencephalograms on Cases 1 and 2; Dr. J. H. Ryffel and Dr. D. Graham Harvey for the lead estimations; and Mr. D. Martin for the radiographs. Our thanks are also due to Dr. D. Hunter for helpful advice and criticism.

REFERENCES

- Blackfan, K. D. (1917). *Amer. J. med. Sci.*, **153**, 877.
 B.M.A. Queensland Branch Rep. of Council (1922). *Med. J. Aust.*, **1**, 148.
 Braun, H. A., Lusky, L. M., and Calvery, H. O. (1946). *J. Pharmacol.*, **87**, Suppl., p. 119.
 Byers, R. K., and Lord, E. E. (1943). *Amer. J. Dis. Child.*, **66**, 471.
 Cadman, H. C. (1902). *Lancet*, **2**, 1458.
 Caffey, J. (1931). *Radiology*, **17**, 957.
 Clark, Norman S. (1950). *Arch. Dis. Childh.*, **25**, 297.
 Findlay, L. (1935). *Postgrad. med. J.*, **11**, 251.
 Ford, F. R. (1945). *Diseases of the Nervous System in Infancy, Childhood, and Adolescence*, p. 661. Springfield, Ill.
 Gartside, V. O. B. (1948). *Med. Offr.*, **80**, 241.
 Germuth, F. G., and Eagle, H. (1948). *J. Pharmacol.*, **92**, 397.
 Goettsch, E., and Mason, H. H. (1940). *Amer. J. Dis. Child.*, **59**, 119.
 Good, P. (1941). *Amer. J. Ophthalm.*, **24**, 794.
 Levinson, A., and Zeldes, M. (1939). *Arch. Pediat.*, **56**, 738.
 Mackay, H. M. M., and Hunter, D. (1949). *Lancet*, **2**, 674.
 McKhann, C. F. (1926). *Amer. J. Dis. Child.*, **32**, 386.
 — (1932). *Arch. Neurol. Psychiat.*, Chicago, **27**, 294.
 — and Vogt, E. C. (1933). *J. Amer. med. Ass.*, **101**, 1131.
 Park, E. A., Jackson, D., and Kajdi, L. (1931). *Amer. J. Dis. Child.*, **41**, 485.
 Rodgers, T. S., Peck, J. R. S., and Jupe, M. H. (1934). *Lancet*, **2**, 129.
 Ruddock, J. C. (1924). *J. Amer. med. Ass.*, **82**, 1682.
 Ryder, H. W., Cholak, J., and Kehoe, R. A. (1947). *Science*, **106**, 63.
 Stephenson, F. W. M. (1898). *Lancet*, **2**, 1473.
 Stetson, C. A. (1947). *J. Maine med. Ass.*, **38**, 10.
 Telfer, J. G. (1947). *J. Amer. med. Ass.*, **135**, 835.
 Vogt, E. C. (1932). *Ibid.*, **98**, 125.
 Weatherall, M. (1948). *Brit. J. Pharmacol.*, **3**, 137.
 Weller, C. V., and Christensen, A. D. (1925). *Arch. Neurol. Psychiat.*, Chicago, **14**, 327.
 Williams, H., Schulze, W. H., Rothchild, H. B., Brown, A. S., and Smith, F. R. (1933). *J. Amer. med. Ass.*, **100**, 1485.
 Wyatt, J. P., and Khoo, P. N. (1949). *Arch. Path.*, **47**, 110.

The Abortion Law Reform Association has just issued a leaflet of advice on how to seek termination of pregnancy. The patient should consult her general practitioner, putting before him the physical and mental grounds for her request. The general practitioner, if he considers the case a reasonable one, will then take a second opinion; and, should this opinion also be favourable, he will arrange for a competent surgeon to perform the operation. The recent judgments in the Bourne and the Bergmann-Ferguson cases give legal precedents to show that it is legal for a patient to ask her doctor for a termination of pregnancy; it is legal for a doctor to consider such a request, and, if he endorses it, to take a second opinion; and it is legal for a surgeon to act on such opinions and to operate, if these opinions are based on preserving the life or the health of the woman and are given in good faith.

KERATODERMA PALMARIS ET PLANTARIS CONGENITALIS

BY

DUNCAN MACAULAY, M.B., Ch.B., B.Sc., M.R.C.P.
D.C.H.

Senior Registrar (Paediatrics), Derby Area No. 1
Hospital Management Committee

The condition known by the above name (there are numerous synonyms) is apparently well known to dermatologists, and a considerable number of reported cases are on record. In 1933 Michael contended that no further cases should be published unless they showed features of particular interest. The following case seems to be the first instance in the white race in which the condition has been traced through seven generations. The family tree is also one of the largest on record.

Case Report

A male child aged 14 months was admitted to the Children's Hospital, Derby, in January, 1950, with an acute respiratory infection. He was the second child of parents who were not related. Apart from his skin

FIG. 1.—Photograph of patient's hands.

FIG. 2.—Photograph of patient's feet.