REFRESHER COURSE FOR GENERAL PRACTITIONERS

INSOMNIA

BY

E. B. STRAUSS, M.A., D.M., F.R.C.P.

Physician for Psychological Medicine, St. Bartholomew's Hospital; Lecturer in Psychological Medicine, St. Bartholomew's Medical College (University of London)

The work of Pavlov suggests that sleep is identical with general cortical inhibition, but this conception requires reconciling with the presence of a sleep centre, which almost certainly exists in the region of the hypothalamus. Such a sleep centre may be considered as initiating inhibitory impulses or, through its influence on adjacent metabolic centres, as producing a biochemical state favourable to the spread of general cortical inhibition. Anything, therefore, that interferes with the spread of cortical inhibition can give rise to sleeplessness. From this it follows that insomnia is not a nosological entity but a symptom.

The factors that can interfere with the physiological spread of cortical inhibition, and thereby give rise to insomnia, lend themselves to classification under two main heads, physical and psychic.

Physical Causes

Pain, Skin Irritation, Dyspnoea, Cough, Fever, and the Like.—It goes without saying that the proper way to treat insomnia arising from any of these causes is to eliminate the irritant stimuli. It must not be forgotten, however, that many patients who sleep badly are unaware of the cause. For example, some patients, without realizing it, are (for them) either too hot or too cold in bed; some people have a "natural" preference for a hard or a soft mattress, and are sleeping on the wrong sort; some use too few or too many pillows.

Toxic and Chemical Factors.—Insomnia may accompany any toxic illness or metabolic disorder. This applies especially to typhus, meningococcal meningitis, acute epidemic encephalitis, influenza, and pneumonia. In encephalitis, it would seem, the sleep centre itself is directly affected. The insomnia associated with true exhaustive states, especially when resulting from motor restlessness—for example, the hyperkinesia of chorea gravis—is probably toxic in origin. It is likely that the insomnia which is often associated with arteriosclerosis is of a dual nature: (a) the deficient elimination of metabolites by the kidneys leads to the accumulation of toxins in the blood stream; and (b) the hypochondriacal preoccupations which are so often met with in this condition give rise to insomnia of psychic origin.

The usual chemical substances responsible for insomnia are, of course, tea, coffee, tobacco, and, nowadays, amphetamine and allied preparations. People who are unduly sensitive to caffeine or the products of tobacco smoking would be wise to eschew tea, coffee, and smoking altogether rather than to content themselves with refraining only in the evenings, as is sometimes advised. Except in cases in which amphetamine is given specifically to lessen the depth of sleep, as in enuresis, the drug should not be given after 2 p.m.

Food Habits.—Some people whose gastric contents are voided slowly and who take heavy meals at night

cannot get to sleep on account of barely perceptible gastric discomfort. Others, especially those who have been told by their doctors that a loaded stomach at night gives rise to insomnia, go to the opposite extreme and are kept awake by hunger, or, if they manage to get to sleep, awaken in the early hours of the morning and cannot get to sleep again. Such persons should have a glass of milk and a few biscuits by the bedside to take as soon as they wake up.

Psychic Factors

Worry and "Overactive" Mind.—Worries due to economic difficulties, domestic strife, troublesome love affairs, and the like can clearly give rise to insomnia. If the general practitioner is prepared to act as a sympathetic audience for half an hour or so (on the principle that "troubles shared are troubles halved") instead of reaching immediately for his prescription pad the moment the word sleeplessness is uttered, many troublesome cases of so-called insomnia will be avoided.

There are many people whose minds become more and more active as evening approaches. These people often find that their minds continue to function automatically as soon as they get into bed. Much as it may go against the grain, they must reduce this mental activity to a minimum for at least an hour before retiring to rest. Such people often choose mistakenly a book which is too interesting, and hence too stimulating, in an attempt to "read themselves to sleep."

Agrypnophobia.—Perhaps this is the most common cause of insomnia. By this term is meant a morbidly anxious preoccupation with the idea of sleeplessness and its supposed "inevitable" evil consequences. Baudouin's "law of reversed effort" (if the will and the imagination are at cross-purposes, imagination wins the contest) is well illustrated by this type of insomnia. By suitable psychotherapeutic techniques the practitioner can help these patients to acquire an attitude of emotional indifference to whether they sleep soundly or not (see *Modern Treatment in General Practice*, 1944, p. 178).

Another phobia often associated with agrypnophobia is the fear of hypnotic drugs. The patient may be terrified of becoming dependent on such remedies and becoming a "drug addict." If hypnotic drugs are given at all, as they may well have to be in cases of this kind, they should be prescribed in full strength, as the anxiety arising out of fear of the drugs has to be overcome as an additional factor; a bad night in spite of hypnotic drugs (prescribed in insufficient quantity) adds to the general agrypnophobia.

Insomnia Complicating the Psychoneuroses.—Insomnia often complicates the psychoneuroses. Although it may not be one of the symptoms of which the patient chiefly complains, it is common for patients undergoing

psychotherapy to sleep even less well than before starting treatment. This is understandable, as patients under treatment are being forced to come to terms in full consciousness with psychic elements which, on account of their unacceptability to the person as a whole, had previously achieved incognito status through repression and other psychic mechanisms. Frequent bad nights interfere with psychotherapy unnecessarily; moreover, the insomnia always disappears when the general neurotic state has been coped with. It is therefore a wise practice to treat such incidental insomnia symptomatically by means of drugs.

Insomnia as a Conversion Symptom.—Insomnia may be the "chosen" symptom in conversion hysteria, especially in cases in which the factor of "secondary gain" is significant. Insomnia of this sort responds best to analytical psychotherapy of one kind or another.

Insomnia in the Psychoses.—It is probably the affective over-stimulation occurring in the psychoses (especially in manic-depressive psychosis and schizophrenia) which gives rise to insomnia in these conditions. As it is impossible to change this pattern solely by psychotherapy, the insomnia must be treated symptomatically by drugs and other appropriate remedies.

Insomnia in Children.—Insomnia in young children may represent a natural resentment of having their sleeping and waking hours forced on them by adults. Every intelligent mother or children's nurse knows how to reduce imperceptibly the child's mental activity as bed-time approaches and how to make going to bed itself a pleasant, if unexciting, procedure and something to be looked forward to. Some children deliberately keep themselves awake because they are afraid of nightmares, sleep-walking, or enuresis. Often a talk to the parents on common-sense lines will deal with such cases, but sometimes psychotherapy is required. Children who fear the dark should be allowed a night-light.

Treatment

As throughout medicine, diagnosis must precede treatment. It must be remembered from the start that, unless there is a lesion affecting the sleep centre itself, insomnia is a symptom, not a disease *sui generis*. Before treatment is undertaken it is essential to be assured that the patient really suffers from insomnia: apart from any question of exaggeration, there is great individual variation in the amount of sleep required by different people; again, elderly people appear to need rather less sleep than do younger persons.

Diagnosis should include a thorough physical examination and the taking of a full medical and psychiatric history. According to the principle of multiple aetiology, all morbid conditions can be regarded from the following standpoints: somatic, psychic, social and environmental, and constitutional and genetic. The physician must try to assess the relative importance of each one of these factors in arriving at his diagnosis. This discipline in itself will indicate what treatment is necessary, it being assumed that, whenever possible, it is better to treat the cause of the symptom than the symptom itself.

Psychogenic insomnia (including agrypniaphobia) may call for skilled psychotherapy, but many cases of insomnia due to worry or intellectual or emotional strain respond well to general measures combined with reassurance and persuasion from the patient's usual doctor.

Hypnotic Drugs

A few of the hypnotic drugs in common use will be considered in detail. However, before doing so it is desirable to lay down certain general rules about their prescription.

(1) Give a sufficiently large dose to be sure of producing an effect.

(2) Leave the anxious type of patient with as little discretion as possible in the matter of dosage and administration.

(3) Whenever possible, give the drug in the form of a mixture. This enables the almost imperceptible reduction of the dose, with or without the patient's knowledge. Moreover, aqueous mixtures are usually less irritating to the gastric mucosa than concentrates.

(4) Whenever there is even the slightest suicidal risk, hypnotic drugs should be handed over to a responsible relative or nurse for administration, and kept in a locked cupboard away from the patient's reach.

Withdrawal of hypnotics can be effected in various ways: by prescribing a hypnotic for use on alternate nights, and later for use only if there have been two unsatisfactory nights in succession; by ordering that it shall be taken only if the patient finds that he does not fall asleep within two hours of going to bed; by reducing the dose, unknown to the patient, by the use of cachets in which increasing amounts of sugar of milk are combined with diminishing amounts of sedative; or by taking advantage of a change of residence, as during a holiday, in order finally to dispense with sleeping draughts altogether.

The Barbiturates

The most efficient and, generally speaking, the most useful member of the barbiturate group is soluble barbitone. It is best given as follows:

Barbitone soluble	..	..	10 gr. (0.65 g.)
Concentrated infusion			
of orange	..	..	½ dr. (1.75 ml.)
Water	..	..	to ½ oz. (14 ml.)

2-4 dr. (7-14 ml.) in water at night when required.

Phenobarbitone, which is so commonly prescribed for insomnia, is a very poor and unreliable hypnotic. If a patient complains of a hang-over effect of soluble barbitone, there is no harm in prescribing:

Amphetamine sulphate tablet 5 mg.
2 tablets every morning and 1½ tablets at mid-day.

There are innumerable proprietary preparations of barbiturates, and all of them have their advocates. It is useful to be able to vary hypnotic drugs at frequent intervals, if only to prevent a patient from establishing a (largely psychological) dependence on any one preparation. *Pentobarbitone sodium*, which is put up in 1½-gr. (0.1-g.) capsules (dose, one or two capsules), is more quickly broken down in the liver and eliminated than most other barbiturates. Its hang-over effect is consequently less. It should always be preceded by sod. bicarb. 20 gr. (1.3 g.), as the presence of acid in the stomach prevents its solution and absorption.

Other Sedatives

Carbromal and *bromvaletone* are brominated urea preparations which are said to be less powerful and less toxic than barbiturates. They are especially useful when the main difficulty is in getting off to sleep. The dosage is 5-10 gr. (0.32-0.65 g.).

Chloral and bromide mixture is useful as an occasional hypnotic. Chloral acts chiefly on the basal cerebral structures, and the bromine salts mainly on the cortex. This combination makes its administration desirable when the patient is agitated.

Chloral hydrate	}	..	each 20 gr. (1.3 g.)
Sodium bromide			
Syrup of orange	..	..	1 dr. (3.5 ml.)
Chloroform water	..	..	to ½ oz. (14 ml.)

2-4 dr. (7-14 ml.) in water at night when required.

Paraldehyde is another valuable occasional hypnotic. It acts rapidly and, as a rule, efficiently, but has the disadvantage of causing the breath to smell of it for long after. In spite of this, it may occasionally become a drug of addiction. It should be given in a dose of at least 2 dr. (7 ml.) (a smaller dose may simply excite), flavoured with an equal quantity of syrup of orange, and the dose made up to 1 oz. (28.4 ml.) with infusion of orange. It may also be given in milk or iced water. In cases of acute mental excitement it may be necessary to repeat the dose in four hours. Paraldehyde may also be given by the rectum (3 to 4 dr. (10.5-14 ml.) in 6 oz. (170 ml.) of starch mucilage). Care should be taken to use a fresh preparation of the drug, as it undergoes oxidation on keeping, with the production of glacial acetic acid, which has been known to cause ulceration of the rectum. In order to prevent smarting, some petroleum jelly should be smeared round the anal orifice before the injection.

Reports of Societies

RENAL FUNCTION

At a meeting of the Renal Association, held at the Ciba Foundation on January 31, the place of Professor E. C. Amoroso, who was to have spoken on some aspects of the foetal kidney, was taken by Dr. ALISON MILES, who gave an account of her work on the kidney in the newborn.

The Foetal Kidney

It was a well-established fact, Dr. Miles said, that infant renal function differed markedly from adult. Renal function in the foetus was less well known. In the later stages of foetal life there was evidence that urine was normally produced, but the birth of infants with renal agenesis which were otherwise normal showed that the foetal kidney was not an essential organ. Many characteristics of foetal renal physiology had been elucidated, but so far anatomical explanations were largely lacking. In general, stillborn infants weighing more than 2.5 kg. had the full number of glomeruli. The first glomeruli to be formed were the innermost; in the newborn child the outer glomeruli were small, with relatively simple capillary loops and a thick visceral layer of epithelium. Except in premature infants, the difference in function of the neonatal kidney could not be attributed solely to smaller numbers of glomeruli; the renal tubules were relatively well developed at birth.

The renal vascular system in the newborn did not seem to have been very closely studied. Dr. Miles gave a preliminary description of the distribution of Indian ink in the kidneys of foetal and neonatal animals which had been injected *in vivo*. It was probable that the pattern of foetal and neonatal circulation was fundamentally similar to that

of the adult, but filtration from the outer glomeruli was probably less adequate. Function changes after birth seemed to be connected with the removal of the placenta, and possibly with the rise in blood pressure coincident with birth.

Amino-acid Excretion

Dr. CHARLES DENT discussed some aspects of tubular function, such as the excretion of uric acid and of phosphate, acidosis of renal origin, and other conditions. Touching on the amino-acids, he pointed out that they were often spoken of as if they were one thing, but actually there were 20 or more of them; sometimes they seemed to behave as individuals and at other times as a group. The pattern of excretion in the urine varied greatly from person to person, although it was constant in a given person. In general the variations were harmless, but there were patterns associated with severe pathological changes in the kidney, liver, brain, and other organs. So far as was known, the minor variations which occurred in normal people were not pathological and might be described as a kind of biochemical individuality, but in some of the analyses which he showed, giving variants on the normal pattern, there were pathological characteristics. The interesting point was that nearly all of the variations of urinary amino-acid pattern occurred in the presence of normal blood levels of all the individual amino-acids. The pattern variations were therefore imposed by variations in kidney tubule function. A further point was that nearly all the tubule dysfunctions, whether for amino-acids, phosphate, uric acid, etc., were hereditary. Dr. Dent also discussed the relationship between urinary pH and NH₃ production, and touched on other aspects of tubular function.

Antidiuretic Hormone

Professor E. B. VERNEY added a short contribution on the neurohypophysis and the kidney and described experiments which were being undertaken to determine the site of the pituitary osmoreceptors. He also discussed the site of action of the antidiuretic hormone. Probably it acted on the distal tubules, causing a reabsorption of water there and, under certain conditions, a diminished reabsorption of chloride.

MUSCLE-RELAXANT DRUGS

A meeting of the Section of Anaesthetics of the Royal Society of Medicine was held on February 2, with the president of the Section, Dr. W. ALEXANDER LOW, in the chair. Dr. H. O. J. COLLIER, chief pharmacologist of Allen and Hanburys Ltd., read a paper on the "Problems of Developing Muscle-relaxant Drugs in the Laboratory."

Desirable Properties

Dr. Collier said that still better muscle-relaxant drugs were needed; they should be more active clinically, and easier to standardize and produce than those at present available. The properties required of them by anaesthetists fell into five groups. First, the specificity of the muscle-relaxant action: the drug should have few side-effects, such as the release of histamine or the blocking of the sympathetic and parasympathetic ganglia. Secondly, the effect on respiration: the anaesthetist might wish the drug to spare the respiratory mechanism, as in lower abdominal operations, or on the contrary to arrest it, as in thoracic surgery. Thirdly, the duration of action: if used only for intubation, for example, a short-acting drug was needed, whereas medium- or long-acting drugs were wanted for surgical operations. Fourthly, compatibility with anaesthetic drugs: there should be no serious antagonism between the anaesthetic and the relaxant, and the relaxant drug might