REFRESHER COURSE FOR GENERAL PRACTITIONERS

SCARLET FEVER

BY

A. JOE, D.S.C., M.D., F.R.C.P.Ed.

Medical Superintendent, Edinburgh City Hospital

During the past few years, when scarlet fever has ceased to be an important cause of death or of serious morbidity, it has become conventional to regard the disease as of little consequence and to dismiss the apparatus of notification and hospital isolation, still widely practised, as a waste of time and resources. This has not resulted from any lack of appreciation of the importance of streptococcal infections, but rather from an awareness of the pathological potential of all such infections, so that paradoxically there has arisen a tendency to exalt the importance of almost all streptococcal infections except scarlet fever, and to dismiss the latter as merely differing from streptococcal tonsillitis in that a rash is present.

Epidemiologically, of course, there is no difference, and, from the standpoint of practical affairs, it is regrettable that in the past there has been too much emphasis on scarlet fever to the exclusion of other streptococcal infections, but is it rational to dismiss the rash so lightly? After all, it is the physical expression of a generalized toxæmia, which, added to the dangerous possibilities of a streptococcal invasion of the respiratory tract, cannot diminish in any way the sum total of the disease process. Again, while we may be thankful that at present scarlet fever is a mild infection we must remember that this was not always so, and Greenwood has pointed out that "scarlet fever is the most beautiful example within the range of epidemiology—an even more striking example than that of smallpox—of change of type, a change which has been effected certainly twice and probably three times in this country within the limits of modern history."

We are quite in the dark about the factor or factors which have operated to bring about these changes. It is tempting to look to our relatively superior environmental and economic conditions as accounting for our present favourable experience, but this cannot be the whole story, as in the first 25 years of the nineteenth century scarlet fever also showed itself in mild epidemic form, although this was a period in which conditions were economically very bad. In view of past history, therefore, and in the absence of knowledge of the factors determining major epidemiological changes, he would be bold who would predict that severe scarlet fever will never occur again. However, in such an event we shall be in possession of remedies much more potent than in the past to deal with the situation.

Definition

It has also been questioned whether the term "scarlet fever" has any claim to be retained in the clinician's vocabulary, and whether it would not be better to refer to all forms of streptococcal infection by some such designation as streptococcal fever. In my view this would result in some loss of precision, and, so long as we realize that scarlet fever can no longer be regarded

as an epidemiological entity, there is no reason why we should not continue to use the term as a convenient description for a well-recognized combination of signs and symptoms which repeats itself often enough to be regarded as a clinical entity. This need not prevent us from recognizing that, as in most other diseases, from time to time one or other common feature may be absent in individual cases, thus to some extent altering the clinical picture.

For practical purposes, therefore, we would define scarlet fever as one of the clinical manifestations of streptococcal infection characterized in the acute febrile stage by faucial inflammation and a generalized erythematous rash, and later by the occurrence of desquamation and of certain complications such as acute otitis media and adenitis caused by lymphatic spread of the streptococcus, and by others such as arthritis occurring at the end of the first week and nephritis which appears at about the end of the third week after onset and the aetiology of which is still undetermined.

Aetiology

To give a reasonable account of present-day views on scarlet fever we must go back to the investigations of the Dicks in 1923 and Dochez in 1924. Their work pointed almost conclusively to the haemolytic streptococcus as the causal organism of the disease, and, although well-marked biological differences were later found to occur within the group roughly defined by this term, no specific relationship could be shown to exist between any particular strain and scarlet fever. Thus Lancefield in 1933, by a precipitation method, showed that streptococci from most cases of scarlet fever fell within Group A, which, however, contained streptococci from many other forms of human streptococcal infection.

Again, by specific agglutination reactions, Griffith in 1935 subdivided the streptococci responsible for human disease into about 40 serological types and regarded these as constituting a bacterial species which he named *Streptococcus pyogenes*. His method of typing, when applied to tracing the spread of streptococcal infection among members of a community, indicated that such common clinical conditions as scarlet fever, streptococcal sore throat, erysipelas, and puerperal sepsis might be caused by the same serological strain, and in view of these findings epidemiologists had to recast their ideas on the subject of scarlet fever. The disease could no longer be regarded as forming an epidemiological entity in the sense that a patent infection derived from a case of scarlet fever would necessarily be another case of scarlet fever; it could be regarded only as one clinical manifestation of epidemic streptococcal disease which might result in a variety of clinical forms to each of which the clinician, for descriptive purposes, might apply a separate term.

Pathology

The discovery of the haemolytic streptococcus as the cause of scarlet fever also provided a basis for the present-day conception of the pathogenesis of the disease. The streptococcus is now regarded as having two main pathological activities—a pyogenic and a toxigenic. When recognizable infection occurs pyogenic activities produce inflammation at the site of implantation, usually the fauces and nasopharynx, while as a result of the toxin-producing activities an exotoxin is liberated into the blood stream and causes a general toxæmia which shows itself by the occurrence of the erythematous rash.

At any stage from the earliest part of the acute eruptive phase until convalescence is well advanced the pyogenic properties of the organism may display themselves and by lymphatic spread set up such septic complications as acute otitis media and cervical adenitis. To account for such complications as rheumatism and nephritis, complications which may follow all streptococcal infections of the upper respiratory tract, various explanations have been advanced, some believing that they may be the result of sensitization following infection, and others ascribing them to the release of, as yet, undifferentiated toxins. So far as pyogenic complications are concerned, however, this is not the whole story, as these, especially in the multiple-bedded wards of hospitals, may not be the pathological expression of the activities of the original invading organism.

Allison and Brown in 1937, employing the method of typing introduced by Griffith, found that when complications occurred in patients treated in large wards, at the time of development of the complication the streptococcus isolated was often of a serological type differing from that grown at the onset of the attack of scarlet fever. From this they concluded that, as a large ward was likely to contain patients with a variety of serological types of streptococcus, complications resulted from the interchange of organisms which was found to occur in such a community. This, of course, would imply a very narrow specificity in the immunity resulting from infection by individual types. With regard to antitoxic immunity acquired from exposure to the soluble exotoxin, or erythrogenic toxin as it is often called owing to its property of evoking the scarlatinal rash, although it has been suggested that there are certain differences within the group of erythrogenic toxins, practical experience shows that these cannot be great, if they exist at all.

Dick and Schultz-Charlton Reactions

Two immunity reactions in the human subject are described in connexion with scarlet fever—the Dick and the Schultz-Charlton tests. It may be thought that a consideration of these is somewhat academic, but for completeness a brief account of each is given.

The Dick test for susceptibility to scarlet fever is performed by the intradermal injection of 0.2 ml. of a suitable dilution of scarlet-fever toxin, which is the filtrate derived from a broth culture of a strain of haemolytic streptococcus known to be a good toxin producer. It is usually carried out on the flexor surface of the left forearm and controlled in the other arm by a similar injection of toxin which has been inactivated by heat. In susceptibles a circumscribed patch of erythema, 10–40 mm. in diameter, appears from six hours onwards and reaches its maximum

intensity in 18 to 24 hours. In immunes no reaction occurs at the site of injection. Pseudo-reactions due to the non-specific constituents of the test material are not numerous.

The test has given interesting results when performed on a large scale in the general population, positive reactors appearing from the fifth to sixth month of life onwards, reaching a maximum of 70 to 80% in the first and second years, and thereafter declining into adult life, approximately 20% being Dick-positive at the age of 20. Theoretically the Dick test should be of assistance in diagnosis, as all cases should be Dick-positive in the earliest stages, and a reversal to negative during convalescence would be absolute confirmation of the diagnosis. In practice, however, the value of the test in diagnosis is limited by the fact that all cases of scarlet fever do not give a positive result in the early stages, and a fair proportion remain positive throughout convalescence. At the present time the main application of the test is in deciding whether active immunization should be conferred on persons exposed to special risk of scarlet fever—for example, entrants to the nursing service of fever hospitals.

The Schultz-Charlton test, often described as the blanching or extinction phenomenon, is performed by the intradermal injection of 0.2 ml. of diluted scarlet-fever antitoxin into an area of skin where the rash is well marked—for example, on the abdomen. In the great majority of rashes in the early stage a localized disappearance of the rash is found over the area of skin infiltrated by the serum in from 6 to 18 hours after performing the test. The reaction is very convincing usually and is specific for scarlet fever; as it is relatively easy to perform, it might be held to be valuable in diagnosis, but in practice there are definite limits to its usefulness. For example, it is best seen in typical rashes in which little assistance in diagnosis is required. On the other hand, in later faded rashes, in faint transient rashes, or in coarse equivocal rashes the results are often disappointing, and even in hospital the test is not much employed nowadays.

It might be mentioned here that for diagnosis in difficult cases throat swabs examined for the presence of haemolytic streptococci are useful, not so much from the point of view of confirming the diagnosis as for their value in excluding the diagnosis of scarlet fever. One or two negative cultures for the haemolytic streptococcus obtained within the first few days of a febrile illness would be very strong presumptive evidence that the condition was not scarlet fever.

Treatment

Home Isolation

In past times, when a high proportion of cases of scarlet fever were isolated in hospital, the main interest of the practitioner was in diagnosis, but over a period of years several factors have combined to make the home isolation of cases of the disease a procedure to be encouraged where circumstances are satisfactory. In the prevalent mild epidemic type, severe cases are unusual and hospital authorities are only too anxious to reserve their diminished number of beds for more severe forms of infectious disease; it has been realized for years that hospital isolation of cases has done little to reduce the general incidence of scarlet fever; and, finally, what has been said about the pathogenesis of complications is a strong argument in favour of retaining

the patient in isolation in his own home. The fact also that we have potent specific remedies, the administration of which calls for only the simplest technical resources, makes the disease one which can be satisfactorily handled by the practitioner.

The patient must have a room to himself, and the following should be set apart for his exclusive use: crockery and cutlery, toilet requirements such as soap, towels, toothbrush, and hairbrush, a thermometer, and a supply of old books and toys which can be burnt afterwards. In the room an overall should be provided, and, near the door, a wash-hand basin with soap, nail-brush, and some dilute disinfectant for the hands if desired. Simple masks consisting of a few thicknesses of gauze, fixed by tapes over the mouth and nose, are also essential. The nurse and medical attendant must always put on a mask and overall when entering the room, and before leaving they should place the mask apart so that it can be included in the next batch of domestic washing, remove the overall, and scrub the hands with soap and water. Articles used in treatment or nursing should be sterilized at once—for example, crockery should be boiled or scalded after food. Rags or paper handkerchiefs should be provided to receive nasal or other discharges and burnt at once. After excreta have been deposited in the water-closet, chamber pots should be wiped out with a black fluid disinfectant. Bed and body linen can be dealt with in the domestic wash tub, and in this connexion it should be remembered that they must not on any account be sent to a public laundry.

General Management

The period of isolation recommended by various authorities for uncomplicated cases varies from two to four weeks; I now adopt a period of three weeks in hospital work, and practical experience teaches that clinical recovery from infection is a safe guide for the termination of isolation. That clinical recovery has been established presupposes a conscientious examination of the upper respiratory tract, and patients with catarrhal or suppurative discharges, even in minor form, should remain in isolation until these have cleared up. If these prove intractable admission to the isolation hospital should be sought so that specialist ear and throat attention may be directed to the eradication of the underlying pathological condition. Bacteriological standards of freedom from infectivity, unless exceptional circumstances call for them, are not generally recommended.

Mild cases should be kept in bed for seven to ten days, but in sharper attacks the patient may need to remain in bed for two to three weeks. During the acute stage the diet should be fluid, although many patients can take a light diet from the beginning, and when the temperature has fallen a full mixed diet should be given. In the early stages headache and restlessness are usually controlled by aspirin. Routine testing of the urine is wise, especially in the third week of illness, and the practitioner should not overlook routine examination of the heart.

Specific Treatment

Under this heading we must consider the sulphonamides, penicillin, and scarlet-fever antitoxin. The results of the administration of sulphonamides have been unspectacular, if not disappointing, and it is doubtful if there are many who employ these drugs

in scarlet fever, especially since penicillin became available.

The action of penicillin is directly on the streptococcus, and daily bacteriological examination of throat and nose swabs for haemolytic streptococci during the course of its administration indicates that the organism disappears from cultures from the third to the fifth day of treatment. Clinically this is paralleled by the subsidence of signs and symptoms of throat inflammation. The specific toxæmia is apparently unaffected and the temperature may not come down for two or three days, whilst the rash pursues its usual course. Although it is unlikely that the streptococcus is completely eliminated, the advantage of reducing the numbers of the organism is obvious and the possibility of complications substantially minimized. When penicillin is used in the treatment of such complications as cervical adenitis, acute otitis media, and rhinitis the clinical evidence of benefit is striking. This is especially the case with otitis media, in which the further complication of mastoiditis is now almost unknown.

The effects of the injection of scarlet-fever antitoxin in the early eruptive phase of scarlet fever are well marked and consistent. The temperature is brought down by crisis in 12 to 24 hours, the rash fades early, and subjective improvement of the patient's general condition is rapidly obtained. Subsequent desquamation may be scanty or retarded in appearance, and in some cases suppressed. Antitoxin treatment, however, is no complete answer to the problem of complications, and, as the disease at present retains its importance mainly on account of these, this somewhat limits the value of the treatment.

The practical recommendation which follows from this very brief summary of the action of penicillin and scarlet-fever antitoxin is that we are prepared to advocate the administration of penicillin to all, or practically all, cases of scarlet fever at the onset of illness. It is our practice to give injections of 100,000 or 200,000 units of penicillin in aqueous solution twice daily, or an injection of the equivalent total dose of procaine penicillin once daily, the treatment being continued for five days. Complications such as acute otitis media, adenitis, rhinitis, and other respiratory infections occurring in the course of the illness require penicillin treatment on the same lines. When the initial toxæmia is sharp, as shown by a temperature of 102° F. (38.9° C.) or over and a bright rash, it is wise to employ antitoxin treatment in addition. One injection of 3,000 to 6,000 U.S.A. units of scarlet-fever antitoxin given intramuscularly at the start of treatment will usually give the required result. Obviously only a minority of cases of the present mild epidemic type of scarlet fever will require antitoxin as well as penicillin treatment.

As on all occasions on which therapeutic sera are administered, the practitioner should be on the look-out for serum sickness which I have thought to be rather more prevalent over the last year or two than in the past after injections of scarlet fever antitoxin. The common form of serum sickness after intramuscular injection is that occurring at an interval of 7–10 days after injection and which is shown by pyrexia, urticarial rash, and, occasionally, by localized oedema of the face and extremities. Should this be troublesome an injection of 1 ml. of adrenaline, 1 in 1,000, followed by a course of one of the antihistamine drugs for two or three days should relieve the patient. Before deciding to give antitoxin the practitioner should inquire into

the history of previous serum injections or such conditions as asthma or hay-fever. If there is evidence of these the practitioner would probably be content to rely on the action of penicillin alone.

Nephritis is a relatively rare complication at present, and when it occurs is much less formidable than formerly. It usually yields to dietetic restriction and alkalis. The occasional patient who develops rheumatic carditis should be carefully watched, and strict rest in bed controlled by blood sedimentation estimations and pulse recordings should be insisted on until these are satisfactory.

ASSEMBLY OF POLISH DOCTORS

The Assembly of Polish Doctors in Great Britain held a meeting in London, at the British Medical Association House, on January 19 and 20. The large attendance included Polish medical men and women from other Western countries. It was the first such meeting since the end of the war, and much fraternal feeling was manifested. The inaugural meeting was presided over by Dr. GEORGE DE SWIET, honorary president of the Polish Medical Association in Great Britain, who welcomed the company in their own language. The Poles, he said, had desired to combine a patriotic rally with some profitable scientific instruction, and had been grateful for this opportunity to come together and exchange views. Many Poles present on that occasion, like their British friends, had been combatants, and, having formed some very warm friendships on the battlefields of Europe and beyond, had felt that they would like to meet again.

Dr. de Swiet went on to thank the British Medical Association and its staff, both medical and non-medical, for the great sympathy and good will it had shown to Polish doctors who had come over to this country at a critical time in their lives. The Poles had felt the need for forming some kind of organization, and in this they had been greatly assisted by the B.M.A. He added that it was 40 years since he himself came to this hospitable country of Great Britain, and it had been his great endeavour to bring the British and Polish peoples together.

Polish Doctors in Britain

General W. ANDERS, who was commander-in-chief of the Polish Forces in the Middle East, paid a high tribute to the work done by the Polish Medical Corps during the war, in particular their treatment of malaria in Russia and their surgical work in Italy. It was stated that there were in Great Britain about 860 doctors of Polish origin and nationality, of whom some 700 had succeeded in finding a settled livelihood. The total number of Poles in this country was reckoned at some 140,000. Some further figures were given by Dr. E. L. STURDEE, who brought the greetings of the Ministry of Health. He said that about 600 Polish doctors now in this country had obtained their medical degrees in Poland before the war, and a small group had qualified in other European universities. There remained nearly 300 who had been trained in this country. He mentioned in particular the difficulties of those who were well on in middle age at the time of their exile and of those who had been specialists in Poland and could not easily find an opening for their specialty in the country in which they had taken refuge.

Dr. G. GREY TURNER, Assistant Secretary of the British Medical Association, spoke a welcome on behalf of that body. He was introduced as one who had fought side by side with the Poles at Monte Cassino, and he recalled the thrilling day when the pink and white flag of Poland was seen flying on the monastery which had for so many weeks menaced their lives. He conceived a great admiration and affection for the gallant Poles. He was also full of admiration for the way in which Polish doctors, some of them

relatively advanced in age, had learned a difficult language and adapted themselves to new conditions of life and practice. He had had the pleasure of meeting a number of Polish doctors working in the Colonies, and there, too, most of them had settled down quite well and found satisfying employment. Dr. L. Kocsis, a Hungarian doctor, offered a welcome on behalf of friendly Continental doctors. He hoped that in due time they would be able to pass on to their respective peoples the knowledge they had accumulated in exile.

Sir Alexander Fleming on Antibiotics

Sir ALEXANDER FLEMING then gave a brief address on "The Search for Antibiotics." Antibiotics, he said, were not a passing phase, they were here to stay. It could be said that their history started with penicillin; all the other antibiotics had been worked out in much the same way. The first discovery was made by the biologist; then came the chemist, who concentrated the material, purified it as far as he could, and synthesized it if possible; then the pharmacologist, who investigated its properties, in particular its toxicity, and how it was absorbed; and after him the physician came into the picture and assessed its therapeutic effect. The pharmaceutical manufacturer had to make the material in quantity, which might be the most difficult thing of all, though it was now not quite such a commercial hazard with the newer antibiotics as it had been with penicillin.

One recent development in the antibiotic field was not purely medical. It had been found that if certain of these substances were administered to chicks or young pigs there was a great enhancement of growth.

On the following day the Assembly met for a clinical session under the chairmanship of Dr. W. TOMASZEWSKI, when addresses were given by Professor Alexander Haddow, director of the Chester Beatty Research Institute, on "Latest Results of Cancer Research," and by Dr. F. A. H. Simmonds, physician and medical director, Clare Hall Hospital, South Mimms, Herts, on "Modern Aspects of the Treatment of Pulmonary Tuberculosis." These were followed by papers in Polish, one by Dr. M. ZALESKI on radiotherapy in cancer, and another by Dr. I. M. SPITZER on the treatment of tuberculous meningitis in the adult.

Cancer Research

Professor A. HADDOW, whose address was to some extent a repetition of one which he gave recently at the Royal Institution (reported in the *British Medical Journal*, December 2, 1950, p. 1271), spoke of the remarkable progress achieved in cancer research during the last 10 years. One of the main difficulties facing the investigator was that the change from the normal to the malignant type of cell was probably due to very subtle transformations, unaccompanied by any gross alterations in the protein structure of the cell or in its immunological properties; hence there was probably no protective reaction on the part of the host, such as was found in bacterial and other infections. Yet another difficulty was that the malignant cell was probably very stable. In the chemotherapy of malignant disease they were expected to undo what amounted almost to a natural process.

He then passed in review the endocrine treatments proposed for various forms of cancer, stating that, apart from stilboestrol in prostatic cancer, these still remained of limited use. He spoke also of the action of urethane, which was characterized by a high degree of chemical specificity. A great deal of attention had been paid to nitrogen mustards, the damaging effects of which on normal tissues had long been recognized. It was not unreasonable to expect to increase the therapeutic effect of these substances by chemical modification, and during the last three years about 250 new types had been synthesized and tested biologically, and some of them clinically as well. Remissions had been obtained in cases of Hodgkin's disease and chronic lymphatic