- Long, P. H. (1950). *Med. Clin. N. Amer.*, **34**, 307.
 — Chandler, C. A., Bliss, E. A., Morton, S. B., and Schoenbach, E. B. (1949). *J. Amer. med. Ass.*, **141**, 315.
 Parets, A. D. (1950). *Ibid.*, **143**, 653.
 Reyniers, J. A., Trexler, P. C., and Ervin, R. F. (1946). *Lobund Rep. Univ. Notre Dame*, **1**, 87.
 — Wagner, M., Luckey, T. D., and Gordon, H. A. (1949a). *Ibid.*, **2**, 1.
 — (1949b). *Nature, Lond.*, **163**, 67.
 Rich, M. L., Ritterhoff, R. J., and Hoffmann, R. J. (1950). *Ann. intern. Med.*, **33**, 1459.
 Smadel, J. E. (1950). *J. Amer. med. Ass.*, **142**, 315.
 Smith, G. N., and Worrel, C. (1949). *Fed. Proc.*, **8**, 253.
 Spink, W. W., Braude, A. I., Castaneda, M. R., and Goytia, R. S. (1948). *J. Amer. med. Ass.*, **138**, 1145.
 — and Yow, E. M. (1949). *Ibid.*, **141**, 964.
 Stephens, P. R. (1950). *Lancet*, **1**, 731.
 Tomaszewski, W. (1951). *British Medical Journal*, **1**, 117.
 Volini, I. F., Greenspan, I., Ehrlich, L., Gonner, J. A., Felsenfeld, O., and Schwartz, S. O. (1950). *J. Amer. med. Ass.*, **142**, 1333.
 Waisbren, B. A., and Glick, D. (1950). *J. Lab. clin. Med.*, **36**, 1001.
 Weinstein, L. (1947). *Amer. J. med. Sci.*, **214**, 56.
 Williams, B., jun. (1950). *Amer. Practit., Phila.*, **1**, 897.
 Zimmerman, L. E. (1950). *Arch. Path.*, **50**, 591.

CARCINOMA OF THE CERVIX UTERI IN AN INFANT

BY

M. D. BABER, M.D., M.R.C.P.
Paediatrician

E. ap I. ROSSER, M.B., M.R.C.O.G.
Gynaecologist and Obstetrician

AND

J. D. O'D. LAVERTINE, M.B., B.S.
*Senior Registrar, Pathology Department
Edgware General Hospital, Edgware*

[WITH SPECIAL PLATE]

With few exceptions malignant tumours of the uterus in children are sarcomatous, but cases of carcinoma have been recorded from reliable sources. Carcinoma of the cervix uteri is one of the very rare forms of malignant disease which occur in childhood. Pollack and Taylor (1947), in their review of the literature on this subject, listed 30 cases of histologically proved carcinoma of the cervix which occurred during the first two decades, and added another case of their own, a patient aged 18 years. Bowing and McCullough (1941) reviewed the case histories of 3,000 patients suffering from malignant disease of the cervix uteri referred to the Mayo Clinic for examination and treatment, and found only one patient of under 20. This patient, whose history they reported in detail and who is included in Pollack and Taylor's list, was a girl aged 13. There is no doubt, therefore, that the disease is very uncommon in childhood.

The youngest authenticated case of carcinoma of the cervix on record appears to be a baby of 6 months, operated upon by Professor Franz in the Charité Frankenklinik of Berlin, and reported by Palmer Findley (1924). Findley stated that he examined the gross and histological specimens, and was satisfied that the lesion was an adenocarcinoma of the cervix. The patient died 14 days after total hysterectomy had been performed.

Bumm (1909) also reported a case of carcinoma of the cervix in an infant of 7 months, but the macroscopic description of the lesion suggests sarcoma botryoides, and no histological confirmation of the diagnosis of carcinoma was offered. Pollack and Taylor included the case in their list, but Bowing and McCullough, who also reviewed the literature since 1862, were not satis-

fied that the criteria for a precise diagnosis were met. Adams (1915) described an advanced case of cancer of the cervix and body of the uterus in a child of 2½ years, but the Pathological Committee of the Royal Society of Medicine, reporting on the specimen, decided that it was a teratoma. Bowing and McCullough regarded a patient of 7 years described by Glöckner (1908) as the youngest case of carcinoma of the cervix on record but appear to have overlooked Palmer Findley's report.

We have recently observed a case of undoubted cervical carcinoma in a child of 15 months. It is presented as another reminder of the importance of malignant disease as a cause of death in childhood, increasingly so since infectious diseases have become well controlled by antibiotics.

Case Report

A baby girl born on March 17, 1948, was brought to hospital on June 30, 1949, because she had passed blood-stained urine during the previous four days, and her mother had also noticed a clot in the vulval orifice on the day before admission. On closer questioning the mother said that the child had recently been straining when passing urine and faeces. She had been drinking well but not excessively; the quantity of urine seemed to be normal and there was no increased frequency of micturition. The stools had been loose and frequent two weeks previously, during a cold, but this symptom had not persisted more than a few days, and the child was thought to be well in other respects. She had not complained of pain.

She was the second child of healthy parents and was born normally at full term. Her birth weight was 7½ lb. (3.4 kg.). She was breast-fed for ten months and there were no feeding difficulties, but in spite of this she had never gained well. On admission her weight was only 17 lb. (7.7 kg.), but her parents had noticed no evidence of weight loss, and, although on closer questioning they agreed that she had been fretful and "not quite herself" for about six weeks, they had always regarded the child as healthy except for a cold two weeks before admission.

Examination showed the patient to be irritable, small for her age, and underweight for her size, though there was no real evidence of wasting. She was pale, but her haemoglobin was 76% and the white cell count was within normal limits. No abnormal physical signs were observed in the heart and lungs. The abdomen was slightly enlarged, and a firm swelling could be felt rising out of the pelvis to the level of the umbilicus. On rectal examination this mass appeared to be cystic and was thought to be distended bladder. The liver was palpable one fingerbreadth below the costal margin and seemed to be somewhat harder than normal. There was no oedema, and the veins of the abdominal wall were not prominent.

On examination under anaesthesia on June 30 catheterization was carried out with some difficulty, as the urethral meatus appeared to be drawn upwards into the vagina: 240 ml. of urine was withdrawn. The abdominal enlargement was reduced but the main mass persisted. The cervix, viewed through a "nasal" speculum, appeared normal apart from a tiny erosion around the external os. No bleeding or discharge was present. Rectal examination revealed a smooth solid pelvic tumour which was continuous with that felt in the abdomen. The whole swelling was mobile to some extent and was roughly pear-shaped, the narrow end being in the pelvis. The uterus was not identified bimanually.

Examination of the urine revealed: specific gravity, 1012; albumin, faint trace; occasional R.B.C.; sterile on culture. The blood urea was 102 mg. per 100 ml. A provisional diagnosis of pelvic kidney was made.

Catheterization was again necessary on July 1, as the bladder was distended, and clear urine was again obtained. A little sanguineous vaginal discharge was observed on this occasion. Cystoscopy on July 2 revealed diffuse inflammatory

changes in the mucosa and suggested that the bladder was lying on the anterior aspect of a pelvic mass. Neither ureteric orifice could be identified owing to oedema of the bladder mucosa. An indwelling catheter was left in the bladder, and a continuous intravenous drip infusion of saline was set up; 250 ml. of blood was also transfused.

On July 4 her general condition improved, and blood urea was now 42 mg. per 100 ml. An intravenous pyelogram demonstrated both kidneys to be normally situated, with a slight degree of pelvic hydronephrosis bilaterally.

Abdominal Section.—On July 4 a midline subumbilical incision exposed the thick-walled bladder, which extended half-way to the umbilicus. The uterus was uniformly enlarged to the size of an adult organ; both tubes and ovaries were infantile and normal in appearance. The peritoneal covering showed no abnormality, but below the utero-vesical fold the bladder was adherent to the lower two-thirds of the enlarged uterus. A small finger in the vagina established that the vaginal cervix felt hard, the induration extending up into the uterus. There was gross supravaginal enlargement of the cervix and a diagnosis of endocervical neoplasm was considered. Hysterectomy was contemplated, but was not performed because of fixation of the uterus to the bladder.

The child's condition remained unchanged for the next week except that she started to pass urine without the help of the catheter. On July 13, after further vaginal bleeding, another examination under general anaesthesia was made. The vaginal vault was now the site of a friable ulcer. After dilatation of the lower vagina some necrotic tissue was removed from the cervical area, and free bleeding followed this procedure. A blood transfusion was given and the patient was returned to the ward, but in the evening of the same day the upper end of the abdominal wound "burst" and a few coils of small intestine prolapsed. The abdominal wall was resutured, and then healed satisfactorily. Report on the biopsy specimen suggested a mixed epithelial and mesodermal malignant tumour.

The patient's condition deteriorated gradually and the abdomen became very distended and hard. Epigastric veins became prominent, the liver enlarged progressively, and ascites developed. She died on August 10.

FIG. A.—Sagittal section of pelvic organs showing hypertrophied bladder, uterus, and vagina, with tumour mass and portion of rectum.

FIG. B.—Inferior surface of liver studded with secondary deposits.

Post-mortem Examination

This was performed on the day of death. The body was that of a rather emaciated female infant. The cervix uteri was greatly enlarged by a mass of hard white growth ($2\frac{1}{2}$ by 2 by $1\frac{1}{2}$ in.—6.25 by 5 by 4.1 cm.) which was friable and extensively ulcerated at the lower end. The body of the uterus was perched on top of the growth (Fig. A), and the tubes and ovaries seemed normal. The bladder was adherent to the front of the mass, drawn up, elongated, and somewhat hypertrophied. There was dilatation of the renal pelvis on both sides, but the kidneys themselves and the ureters appeared normal. Most striking of all was the enormous enlargement of the liver (weight 1,550 g.) due to the presence of large numbers of secondary deposits (Fig. B), many of which were cystic and contained mucoid material. No other metastatic deposits were found, and there was nothing abnormal in the cardiovascular, respiratory, alimentary, and endocrine systems. The primary growth and liver metastases were examined histologically, the former showing a poorly differentiated adenocarcinoma with more solid cellular areas (Plate, Fig. 1). The growth and invaded tissues were very closely intermingled, which was responsible for the original impression that the biopsy material represented a mixed tumour. In the liver, however, the growth was a well-differentiated papillary adenocarcinoma, in some places again showing intimate mixing with the invaded tissues (Plate, Fig. 2).

The sections were referred to the Consultant Panel in Morbid Histology of the Pathological Society of Great Britain and Ireland and to the British Empire Cancer Campaign. They agreed that the tumour was a papillary adenocarcinoma, probably of endocervical origin.

We are indebted to the Consultant Panel in Morbid Histology of the Pathological Society of Great Britain and Ireland and to the British Empire Cancer Campaign for their opinion on the sections, to Mr. R. Trevor Jones for performing the cystoscopic examination, and to Miss M. H. Shaw for the photographs.

REFERENCES

- Adams, J. E. (1915-16). *Proc. R. Soc. Med.*, 9 (Obst. Sect.), 45.
 Bowing, H. H., and McCullough, J. A. L. (1941). *Amer. J. Roentgenol.*, 45, 819.
 Bumm, E. (1909). *Disch. med. Wschr.*, 35, 1331.
 Findley, Palmer (1924). In *Abt's Paediatrics*, 4, 1234. Saunders, Philadelphia.
 Glöckner (1908). *Z. Geburtsh. Gynäk.*, 63, 182.
 Pollack, R. S., and Taylor, H. C. (1947). *Amer. J. Obstet. Gynec.*, 53, 135.

ADDENDUM.—Since submitting this report for publication another account of two cases of carcinoma of the cervix in the first year of life has come to our notice. This latest report, by G. P. Heckel in *Paediatrics*, 1950, 5, 924, describes two children of 7 and 11 months of age, both suffering from

adenocarcinoma of the cervix uteri, demonstrated by histological sections. The younger died of the disease on the day of admission, having had vaginal bleeding for about one month. The second was treated by total hysterectomy eight days after the onset of vaginal bleeding, and was reported well and developing normally 21 months after the operation.

DISSECTING ANEURYSM OF THE AORTA WITH RENAL COMPLICATIONS

BY

H. BREBNER, M.B., Ch.B., M.R.C.P.

Senior Registrar, Kingston Hospital

[WITH SPECIAL PLATE]

Chronic dissecting aneurysm has ceased to be a medical curiosity, and the absence of pain at the time of dissection has often been noted (Gouley and Anderson, 1940; Flaxman, 1942). In the following case there was no history of pain until one month before death, and at necropsy a long-standing dissecting aneurysm was found.

Case Report

The patient, a land-surveyor aged 52, had always enjoyed good health. His parents were over 80 years of age. He consulted his doctor in September, 1947, because of breathlessness on exertion. There was no history of chest pain. His blood pressure was 140/80 and there was no albuminuria. He continued to lead a normal active life. On November 18, 1949, he developed acute pain in the epigastrium radiating to the left side and left shoulder and finally settling in the left loin. He was admitted to a surgical ward, where tenderness was noted in the left loin. His blood pressure was 170/110 and the urine contained albumin but no red cells. Renal colic was diagnosed; the pain ceased in a few days, and he went home. On December 6 he was admitted suffering from increasingly severe dyspnoea. There

FIG. A.—Aorta at site of dissection. The normal ascending part is seen, then the slit of entry. The smooth surface of the false sac is seen just to the left of the probe at the top.

FIG. B.—Kidneys. Dissection of intrarenal branches of left renal artery can be seen. Multiple infarcts in left kidney are visible. (The left kidney is the lower in the Figure.)

had been no recurrence of abdominal pain and he had no pain during his stay in hospital. He gave no history of chest pain at any time.

On examination he was distressed, with acidotic breathing and bleeding from the lips and gums. His complexion was sallow and the mucous membranes were pale. Cardiac enlargement and gallop rhythm were present; there were no murmurs. Blood pressure was 220/150. Numerous moist rales were audible over both lower lobes, but there was no increase in jugular venous pressure nor any oedema. Abdominal palpation was negative. The fundi showed numerous exudates and a few haemorrhages. The urine contained a trace of albumin and very occasional white and red blood cells but no casts. The blood urea was 120 to 169 mg. per 100 ml.; haemoglobin, 74%; red blood cells, 4,020,000. The urinary output was maintained above 40 oz. (1.14 litres) daily, but uraemia persisted and the patient died on December 21.

Necropsy

The heart was enlarged, the right and left ventricular walls being thickened (0.6 cm. and 1.5 cm. respectively). The aorta (Fig. A) was normal up to a slit-shaped opening in the wall 1.2 by 0.3 cm. across, which lay between the left subclavian and the left common carotid arteries, and was the apex of a dissecting aneurysm. This did not involve either of these arteries. The false sac had a smooth lining and extended almost completely round the true descending aorta in its upper 3 in. (7.5 cm.). The dissection then veered more to the left side, where it continued, narrowing, to the iliac bifurcation and then along the left iliac artery to the inguinal ligament, but only a little way along the right iliac artery. No re-entrant was found. There was dissection of the left renal artery, the true lumen being flattened and narrowed. The dissection continued into the main intrarenal branches (Fig. B). The right renal artery was normal. The left kidney (9 by 5 by 4.5 cm., weight 106 g.) had an irregular shape and on section showed a number of pale yellow infarcts. The right kidney (10 by 7.5 by 5.5 cm., weight 210 g.) appeared normal. The lungs showed congestion and

J. B. RENNIE, J. A. MILNE, AND J. SOMMERVILLE: TRIAL OF A CINCHONINIC ACID
DERIVATIVE IN SOME COLLAGEN DISEASES

FIG. 1.—Case 6. Biopsy of deltoid muscle. A small artery is seen, the lumen of which is completely obliterated by fibroblastic proliferation. In the centre are some eosinophil leucocytes and particles of nuclear dust. The adventitia is infiltrated with inflammatory cells in which occasional eosinophil leucocytes are present. (H. & E. $\times 200$.)

FIG. 2.—Case 7. Skin of abdomen before treatment. Note the stretched atrophic-looking epidermis. The corium is made up of dense parallel bundles of collagen with no differentiation into papillary and reticular layers. (H. & E. $\times 70$.)

FIG. 3.—Case 7. Skin of abdomen before treatment. This section is stained for elastic fibres and shows these lying parallel to the epidermis. (Orcein. $\times 200$.)

M. D. BABER, E. ap I. ROSSER, AND J. D. O'D. LAVERTINE:
CARCINOMA OF THE CERVIX UTERI IN AN INFANT

FIG. 1.—Primary growth ($\times 380$): poorly differentiated area.

FIG. 2.—Metastasis in liver ($\times 380$): well-differentiated papillary adenocarcinoma.

FIG. 4.—Case 7. Skin of abdomen after treatment. The epidermis is thicker and there is evidence of rete pegs. The papillary layer of the corium has re-formed. The reticular layer is still composed of dense collagen, but near the papillary layer there is evidence of breaking up and the fibres are becoming wavy. (H. & E. $\times 70$.)

FIG. 5.—Case 7. Skin of abdomen after treatment. The elastic fibres in the upper part of the corium are seen to have assumed their normal relationship, being at right angles to the epidermis. (Orcein. $\times 200$.)