recovered it will be important to investigate its possible relation to the MNS system and to find out whether, for instance, the gene controlling the antigen is an allele of S and s, or possibly occupies a third locus closely linked to those for MN and Ss. The rabbit which yielded our serum had been immunized only with the red cells of group M persons. If the new antibody is an immune one, therefore, the antigen must in this case have been present in the blood of an M person. If this is so the slight possibility is eliminated that the agglutinations observed were the property of a rare allelic variety of N.

A Suggestion

Our object in describing this somewhat abortive investigation is to draw attention to the existence of one or possibly two antigens which appear to be commoner in negroes than in Europeans, antigens which are possibly related genetically to the MNS blood group system, and to which rabbits fairly readily make antibodies. It is also possible that such an antigen might, on very rare occasions, if present in blood used for transfusion, provoke the development of the antibody in the serum of the recipient. We should therefore like to suggest that workers who, unlike ourselves, have direct access to negro populations should consider making an attempt to repeat the preparation of such an antibody and to investigate the genetics and anthropology of the antigen detected by it. It is suggested that several batches each of about 10 rabbits should be immunized with the pooled red cells of about 20 group N negroes, each batch of rabbits being injected with a different pool of negro cells. As some group N negroes certainly carry the antigen and group M negroes may not, it seems best to choose only those of group N, whatever may be the true genetic relationship of the "new" antigen. The serum of each rabbit should be divided into two or three portions and each portion absorbed with the red cells of one group A₁, MN European, and then tested against the cells of each of the negro donors.

We shall be glad to exchange sera (including anti-M and anti-N) with any worker who is contemplating such an investigation, or with any worker who has found anti-M or anti-N sera giving anomalous results, or who has discovered an unidentifiable antibody in the serum of a patient transfused with the blood of one or more negroes.

At the time of going to press we have just obtained a further sample of the blood of the original positive subject, through the kindness of Dr. D. R. Cannon and Mr. N. A. Barnicot. We are continuing our investigations with the collaboration of Dr. J. N. Marshall Chalmers.

We are indebted to Dr. D. R. Cannon, of the Yaba Medical Research Institute, Lagos, for sending the specimens and to Dr. J. A. Fraser Roberts for advice on statistics.

REFERENCES

- Landsteiner, K., Strutton, W. R., and Chase, M. W. (1934). *J. Immunol.*, 27, 469.
Wiener, A. S. (1943). *Blood Groups and Transfusion*, 3rd ed., p. 264. Springfield, Illinois.

Adrenocorticotrophic hormone being in very short supply, every possible source is being explored. In *Nature*, February 17, page 280, is a brief note on its preparation from whale pituitaries by Norwegian workers. One kilo of pituitaries, from about 50 fin whales and blue whales, was extracted by methods similar to those used on pig and sheep pituitaries and yielded 0.5 g. A.C.T.H. with an activity equal to 1.4 times the Armour standard.

Medical Memorandum

Islet-cell Tumour of the Pancreas

Islet-cell tumour of the pancreas is a well-recognized, though uncommon, cause of spontaneous hypoglycaemia. It may occur at any age, but is rare in infancy and after the age of 50. Most cases are found between 20 and 50, although a case in a 24-hour-old infant has been reported (Sherman, 1947). Our patient was 69, older than any other upon whom a successful operation has been performed. Howard, Moss, and Rhoads (1950) summarized all the 398 cases described up to 1950. Of these, some 200 have been operated upon for hyperinsulinism. In 150 patients the tumour or tumours were enucleated, while in about one-third of this number, no tumour being found, a partial resection of the pancreas was carried out. The operative mortality was about the same in both types of operation (9%), and the end-results were good in both sets of survivors, some 85% being cured.

With its diverse symptomatology the disorder may readily be mistaken for epilepsy, cerebral tumour, neurosis, psychosis, narcolepsy, and many other conditions. In elucidating such symptoms, islet-cell tumour may profitably be remembered. The case here reported is typical except for the age of the patient.

Case Report

A married woman aged 69 was admitted to hospital on January 23, 1950. In childhood she had had attacks of measles, whooping-cough, and bronchitis, but during forty years of married life had had no illnesses or operations, apart from mumps some 20 years ago. She was an only child; her parents died in old age and she had a healthy son and daughter.

Earlier Symptoms

Symptoms due to the islet-cell tumour had first appeared in 1945, when one morning she failed to waken, remaining unconscious until 11 p.m. that day. She then recovered spontaneously, ate a meal, and slept throughout the night. She remained well until November, 1946, when, after an attack of tonsillitis, she again failed to waken at the usual time. Four or five similar attacks followed, consciousness being regained in the afternoon or evening. On November 12 she was first admitted to the Royal South Hants and Southampton Hospital for investigation. She was seen to be a healthy-looking woman, and appeared rather younger than her age. Mentally she was a little vague and was inclined to repeat phrases. Her speech was slurred. Examination of her central nervous system revealed no abnormalities. The arcus senilis was present. The blood pressure was 170/110 mm. Hg, but the heart was otherwise normal. No abnormalities were noticed in the chest or abdomen. Lumbar puncture showed a normal fluid.

On November 15 she refused her lunch and lapsed into semicoma. The blood sugar was 38 mg. per 100 ml. On November 17 two further attacks occurred, which were relieved 20 minutes after the oral administration of 10 g. of glucose. Next day another hypoglycaemic episode responded to intravenous administration of 50% glucose solution. On November 19 a glucose-tolerance test was performed, with these results: 8 a.m. (fasting), 44 mg. blood sugar per 100 ml. (capillary blood); 50 g. glucose was then given orally; 8.30 a.m., 99 mg.; 9.45 a.m., 117 mg.; 10.15 a.m., 153 mg.; 10.45 a.m., 124 mg.

On November 27 the patient had no breakfast, and two-hourly samples of blood were taken, the glucose content of which was: 8 a.m., 47 mg. per 100 ml. (capillary blood); 10 a.m., 38 mg.; noon, 35 mg.; 2 p.m., 31 mg.; 4 p.m., 89 mg. She thus fulfilled the diagnostic criteria of Whipple for the diagnosis of hyperinsulinism, which are that (1) the

fasting blood-sugar level should be less than 50 mg. per 100 ml.; (2) there are symptomatic attacks of hypoglycaemia, which occur only when the patient is fasting; and (3) relief of such attacks is produced by administration of glucose, without the patient knowing that glucose is being given.

The attacks of hypoglycaemia were therefore thought to be due to an islet-cell tumour of the pancreas, the low fasting blood sugar excluding the possibility of reactive hyperinsulinism. The patient refused to undergo operation and was discharged home on a high carbohydrate diet. She remained free from attacks until shortly before Christmas, 1949. She had been gaining weight considerably and decided to reduce it by restricting her intake of carbohydrate. Thereupon the attacks of hypoglycaemia returned, her failure to waken in the morning again drawing attention to them.

Present Attack

The patient was readmitted to hospital on January 23, 1950. During the first few days she had several attacks of hypoglycaemia shortly before breakfast. She usually responded to subcutaneous injections of adrenaline, 8 minims (0.5 ml.), followed by oral glucose. It was then found that, in order to prevent the attacks, it was necessary to give her a carbohydrate meal at 2 a.m., a procedure which was followed until her operation.

On January 25 her blood sugar at 11.15 a.m. was 64 mg. per 100 ml. after a normal breakfast at 7.30 a.m.; on January 28, at 4 p.m., after her usual lunch at midday, it was 32 mg. per 100 ml. On neither occasion was the patient clinically hypoglycaemic, a point which is referred to later.

On January 27 food was not given; her blood-sugar levels were: at 6 a.m., 65 mg. per 100 ml. (venous blood); 8 a.m., 31 mg.; 10 a.m., 14 mg.; 12.15 p.m., 12 mg.; 1 p.m., 23 mg.; 2 p.m., 28 mg.

These figures illustrate two features known to be associated with islet-cell tumours of the pancreas: (1) the tendency to spontaneous recovery from the hypoglycaemia; and (2) the lack of correlation between the clinical attacks and the blood-sugar level. At 12.15 p.m., when the blood sugar was 12 mg. per 100 ml., she merely felt faint, but was still able to co-operate, yet at 2 p.m., when her blood sugar had risen to 28 mg. per 100 ml., she was incoherent, restless, and so difficult to manage that the tests could not be continued.

On January 30 an intravenous glucose-tolerance test was performed, the patient fasting; the blood-sugar levels were: fasting, 33 mg. per 100 ml. (capillary blood); 50 ml. of 50% glucose solution was given intravenously; after five minutes the blood-sugar level was 158 mg. per 100 ml. (capillary blood); 15 minutes, 171 mg.; 35 minutes, 335 mg.; 45 minutes, 292 mg.; 60 minutes, 111 mg.; 90 minutes, 84 mg.; 120 minutes, 71 mg.; 180 minutes, 64 mg.

On February 2 a series of fasting blood sugars was obtained, followed immediately by intravenous injection of 40 ml. of 50% glucose solution, with the following results:

Fasting	Capillary Blood Sugar	Venous Blood Sugar
At 10 a.m.	39 mg. per 100 ml. . .	28 mg.
" 11 "	34 " " " " . . .	
" noon "	31 " " " " . . .	
" 1 p.m.	30 " " " " . . .	
" 2 "	31 " " " " . . .	22 mg.
5 min. after glucose . . .	225 " " " " . . .	
10 " " " "	284 " " " " . . .	
30 " " " "	128 " " " " . . .	
45 " " " "	77 " " " " . . .	
60 " " " "	69 " " " " . . .	
90 " " " "	38 " " " " . . .	
120 " " " "	36 " " " " . . .	
150 " " " "	28 " " " " . . .	

Gastric juice was withdrawn during the first part of the test to ascertain if there was any alteration in the secretion of free hydrochloric acid; unfortunately achlorhydria was found.

The patient now consented to operation. The tumour was enucleated on February 15 by Mr. Raymond Purvis, under general anaesthesia administered by Dr. Oakley White.

The condition of the patient was good, but by 10 p.m. she was not yet conscious. A catheter specimen of urine was found to contain large quantities of sugar and acetone. At 1 a.m. on February 16, 40 units of soluble insulin was given and 40 units more was injected at 5 a.m., the urine being still loaded with sugar and acetone. At 9 a.m. she was

conscious; her urine now contained only a trace of sugar and no acetone. Her blood sugar was 104 mg. per 100 ml. On February 17 the urine again contained large quantities of sugar and acetone; the blood sugar was 304 mg. per 100 ml. Thirty units of soluble insulin was given. Thereafter no further glycosuria occurred and no more insulin was needed.

The histological report of the tumour was as follows: "A tumour 1.2 cm. in diameter, of grey tissue, clearly defined but not encapsulated, with a little normal pancreas around. Histologically this is an islet-cell tumour. The cells resemble those of the islets, but are larger. No mitoses are seen in the cells. The cells are in masses divided by strands of dense fibrous tissue. There is no definite capsule. The pancreas around is compressed, and a little fat necrosis with leucocyte reaction is seen. The tumour appears simple."

On March 23, with the patient fasting, blood-sugar estimations were carried out at hourly intervals from 10 a.m. to

Islet-cell tumour. Van Gieson stain. ($\times 350$.)

2 p.m. At this point 30 ml. of 50% glucose solution was given by intravenous injection, and during the next 150 minutes eight further blood-sugar estimations were carried out. In this way a series of figures comparable to those found on February 23 was produced.

Fasting	Capillary Blood Sugar	Venous Blood Sugar
At 10 a.m.	118 mg. per 100 ml. . .	
" 11 "	97 " " " " . . .	
" noon "	102 " " " " . . .	
" 1 p.m.	94 " " " " . . .	
" 2 "	88 " " " " . . .	79 mg.
5 min. after glucose . . .	194 " " " " . . .	
10 " " " "	176 " " " " . . .	
30 " " " "	154 " " " " . . .	
45 " " " "	118 " " " " . . .	
60 " " " "	102 " " " " . . .	
90 " " " "	88 " " " " . . .	
120 " " " "	88 " " " " . . .	
150 " " " "	86 " " " " . . .	

The difference between the results of these tests before and after operation is well illustrated in the tables.

The patient was discharged from hospital on April 10, cured of her spontaneous hypoglycaemia and with her wound almost healed.

I am indebted to Dr. K. M. Robertson for permission to publish the case, and for his help and encouragement; to Mr. Raymond Purvis, who performed the operation; to Dr. Oakley White, the anaesthetist; to Dr. H. H. Gleave for the histological report; to Dr. R. H. Slater for the biochemical analyses; and to Dr. C. E. Brierley, the patient's general practitioner.

F. J. Y. WOOD, M.B., B.Chir., M.R.C.P.,
Late House Physician,
Royal South Hants and Southampton Hospital.

REFERENCES

- Howard, J. M., Moss, N. H., and Rhoads, J. E. (1950). *Int. Abstr. Surg.*, 90, 417.
Sherman, H. (1947). *Amer. J. Dis. Child.*, 74, 58.