REFRESHER COURSE FOR GENERAL PRACTITIONERS

EMPHYSEMA

BY

K. ROBSON, M.D., F.R.C.P.*Physician, St. George's Hospital, London*

Pulmonary emphysema is a term that has come to be applied in a rather loose way to a number of conditions which, aetiologically speaking, are quite dissimilar. We speak of large-lunged and small-lunged emphysema, the former conjuring up visions of breathless bandmen (who may have been drummers) and the latter of leaner, more scholarly individuals. Then there are interstitial emphysema, compensatory emphysema, and that peculiarly localized variety spoken of as bullous or cystic emphysema. Despite their apparent dissimilarity, there are features which unify all these varieties. These are loss of elasticity and the destruction by rupture of interalveolar septa.

Pulmonary Distension

Among the essential properties of lung tissue an important place must be given to elasticity and the power to inflate and deflate. It is a prerequisite of our existence that this inflation and deflation should take place. The capacity of the lungs to contain air varies enormously: a fit man has little difficulty in blowing three litres at a "vital capacity" determination, and all this air must have been accommodated in full inspiration by the stretching of pulmonary tissue and bronchi, and alteration of their mutual relationships. This expansion and recoil repeats itself about a thousand times in each hour in health. With the passage of years, even in the absence of disease, changes take place. In particular, the recoil diminishes and elasticity gives place to rigidity, fixity, and shrinkage—to small-lunged emphysema of the elderly.

A similar loss of elasticity is produced after some years of pulmonary distension: shrinkage of part of the lung owing to collapse or disease will necessitate compensatory changes, and if these cannot be wholly met by shift of the heart and mediastinum there will be overdistension of the parts of the lung which are still aerating. Compensatory or hypertrophic emphysema may for a time be a true description of this process, but later the continued overstretching will produce distension beyond the limit of reversal.

Rupture of the interalveolar septa is a feature, in greater or lesser degree, of all varieties of emphysema. It is seen in its grossest form in the bullous or cystic variety. Here, the very property of inflation is turned to a disadvantage. Some minor alteration in the relationship between bronchial tube and pulmonary tissue permits distension of an area of lung without the possibility of its deflation. Septal rupture proceeds apace and a localized pulmonary balloon within the lung results. This same process, essentially, operates in inflammatory disease; in certain varieties of bronchopneumonic affection and in tuberculosis with lesions of important extent in bronchial tubes it makes possible the development, from a small area of destructive pulmonary breakdown, of a tension (or, as it might equally

well be termed, a distension) cavity. It is probable that up to a certain limit of distension, and perhaps depending to some extent on the duration of the distension, the process may be a reversible one. Once this limit has been passed the process is irreversible.

Generalized bronchial inflammation, particularly if associated with bronchial spasm, is likely to lead to emphysema. In the production of cystic or bullous emphysema the state of affairs is postulated where the obstruction to the outlet of air is complete or nearly so. In bronchitis with spasm, well exemplified in chronic recurrent asthma, the obstruction is not complete, but over weeks and months it exerts a telling effect. Intimately connected with the whole subject is the fact that while inspiration is a powerful movement, involving active work for various groups of muscles, expiration is in the main passive, at least so far as costal respiration is concerned. This means that it is easier to inflate (or distend) the lungs than to deflate them. In fact, it is only by cultivation of diaphragm breathing that really forcible expiration is possible. The bearing which this functional distinction between inspiratory and expiratory movement has on pulmonary distension will be obvious.

Pulmonary Reserve: Effect on Function

Increasing experience of patients who have had a pneumonectomy or lobectomy, or of individuals in whom some disease process has reduced the available pulmonary tissue, is bringing a fuller realization of the immense reserves of respiratory function of which human beings are possessed. Equally striking is the dire plight of patients who have had a pneumonectomy after a faulty assessment of their reserves. Considerable damage must have accrued over the years in an individual before symptoms are produced by emphysema. On the whole, the younger the subject of lung ablation, whether surgical or the result of disease, the better is his adjustment. Emphysema produces some degree of anoxia. The total surface area over which air and haemoglobin can make contact is reduced. The distension obliterates altogether some capillaries and reduces the lumen of others. The air tends to be drawn into the relatively distended peripheral parts of the lung, and the impaired expiratory force makes for the stagnation of air there and hinders its replacement with fresh atmospheric air containing the normal complement of oxygen. All these factors make emphysema a serious disability.

Further aggravation may come through the operation of the Hering-Breuer reflex. Normally the impulse to terminate the process of inspiration arises in pulmonary stretch receptors. When a certain degree of lung distension has been reached these receptors are stimulated, inspiration ceases, and expiration begins. If there is gross localized emphysema, as for instance in bullous or

cystic emphysema, stimulation of the receptors may occur in the distended portion of lung, while the remainder is still capable of some expansion which, from the functional point of view, would be profitable.

Diagnosis

Emphysema is often diagnosed, and in many cases probably correctly so, but it is by no means easy to lay down reliable criteria for its recognition. It can occur as an acute process, though it is more frequently encountered as a chronic and static phenomenon. Sometimes there is a familial incidence. Patients are often in early middle life and their complaint is breathlessness. They are particularly troubled on exertion, when they may show a tinge of cyanosis.

The conformation of the thorax and the posture of the ribs often indicate some overdistension of the lungs. Measurement shows poor thoracic expansion, often less than 2 in. (5 cm.) being evident between full expiration and full inspiration. There may be hyperresonance and obliteration of liver and heart dullness, though these findings are frequently equivocal. With the stethoscope one is often struck by the poor air-entry and the very quiet respiratory murmur. Prolongation of expiration and added sounds, particularly rhonchi, are features of associated bronchitis and bronchial spasm rather than of emphysema itself. The radiographs are often striking. The ribs lie in an exaggerated inspiratory position, the outline of the diaphragm is ragged with a scalloped outline, and the heart shadow is narrow. One gets the impression of overdistended lung from the unduly black appearance of the parenchyma. If there are grossly localized areas of emphysema, they are often readily apparent, and sometimes the walls of cystic or bullous spaces can be traced as hair-like lines.

Chronic bronchitis may be the sequel of an acute attack or it may develop insidiously in association with emphysema. The two conditions are intimately connected and emphysema is undoubtedly aggravated by chronic bronchitis, more especially by exacerbations of acute bronchitis. The four months from November to February are the critical ones for bronchitic patients. They mostly lose time at work each year after the age of 50. As the years go by acute exacerbation tends to occur earlier in the winter, and it may be repeated once or twice before the spring.

Management

With degenerative changes at work and so much structural alteration in being it may well be thought that little enough can be done to help emphysematous people. The treatment of emphysema falls under three heads: (1) the prevention and cure of attacks of bronchitis with spasm and of asthma; (2) the institution of measures to produce the maximum use of the available pulmonary tissue and particularly to develop expiration; and (3) the consideration of any procedures which may be indicated to meet special varieties of emphysema.

It should be an axiom that bronchial spasm should be relieved as quickly and effectively as possible whenever it occurs. Once that pattern of functional behaviour which results in spasm of the bronchial wall has been set, the pathway to its production is facilitated and the spastic response to various stimuli occurs all too readily. Whatever remedy has been found to be most efficacious in an individual case, whether it be an antispasmodic drug, a member of the antihistamine series,

or one of the varieties of adrenaline or ephedrine, it must be used early and in adequate doses. In the bronchitis of middle-aged and elderly people it often pays to saturate the patient with an antispasmodic, and that good old-fashioned remedy "iodide and stramonium" has few equals.

Potassium iodide	3 gr. (0.2 g.)
Tincture of stramonium	15 min. (0.9 ml.)
Liquid extract of liquorice	30 min. (1.8 ml.)
Spirits of chloroform	5 min. (0.3 ml.)
Water	to 1 fl. oz. (28.4 ml.)

It should be taken regularly, not intermittently, until a full week has elapsed after the relief of symptoms. Penicillin and the sulphonamides have a distinct place in these cases, though their value is often discovered only casually when they are being given for some other condition, as, for instance, a urinary infection. Antibiotics are best given by injection; sulphonamides should be used in orthodox doses, to a total of 35 g. in a week. Inhalation therapy can sometimes be most valuable, but, whatever proprietary preparation is used, the instructions supplied with the form of apparatus concerned should be followed in exact detail.

To experience more than one exacerbation of acute bronchitis during a winter is very discouraging and the condition is often prolonged well into the spring. For the few who can manage it, three weeks spent at one of the French spas may help them to a better level of health. Prevention of bronchitis in those who are subject to it may involve a change of habits or even of work. Patients with this sort of chest trouble are often advised that they should have an outdoor job. Generally speaking, indoor work is to be preferred in Britain. As Sir Clifford Allbutt put it many years ago, "fresh-air treatment does not mean treatment by hurricane." Far too often the climatic conditions encountered in this country between November and March are such as to aggravate very seriously the bronchitic tendency. Restrictions must be accepted, and going out on cold, raw winter nights, especially when fog may be encountered, should be forbidden. Any abrupt change from warm to cold is a potential aggravating factor. The acceptance of this way of life will certainly have a preventive value, and it is essential if the bronchitis and emphysema are merely additional burdens in a respiratory system already impaired by scarred and fibrotic tubercle or other established disease.

Somehow or other an attempt must be made to improve the resilience of the thorax. The barrack-square ramrod posture, a lunge forward with the arms flung wide to greet the wintry dawn at an open window, is an approach as wrong as it can well be. Rather should the emphasis be on relaxation, with the patient lying down comfortably on a couch endeavouring to dispel all tension. Then when he is calm he makes a deliberate effort to move and control the diaphragm. The facts about diaphragm movement, and what it can achieve, and its particular value in expiration are explained to him. He must make a conscious effort to cultivate diaphragm breathing and lower-rib expansion. He must watch himself in a looking-glass part of the time to break the habit of upper respiratory breathing with tensing of the sternomastoids and trapezii and elevation of the whole shoulder girdle. He must practise his diaphragm diligently until the movement is second nature to him.

Just occasionally some special measures may be applicable. The induction of a pneumoperitoneum can

certainly give a ragged diaphragm a smooth contour, but on the whole the long-term results have been disappointing. Some unfortunate results also have attended well-intentioned efforts to remove emphysematous bullae or cysts. There is some evidence, however, that, paradoxically enough, a phrenic interruption may be beneficial in cases of gross localized emphysema at a base where the reflex control of respiration is dominated by the distension in this situation. When some relaxation is produced inspiration and expiration of the less distended parts of the lungs assume more normal proportions and rhythm.

DR. HELEN KELLER IN ENGLAND

On the eve of her departure from this country to South Africa Dr. Helen Keller was the guest of honour at a dinner-party given by Sir Ian and Lady Fraser at the House of Commons on Wednesday, February 28. Dr. Keller was accompanied by her lifelong friend and guide, Miss Polly Thompson, through whom she has contact with the world around her, and by Mr. Alfred Allen, the assistant director of the American Foundation for the Blind. A distinguished and representative company was gathered to pay tribute to

Sport and General Press Agency, Ltd., London

Dr. Helen Keller in conversation with Sir Ian Fraser.

Dr. Keller's courage and example. At the age of 2, following an attack of scarlet fever, she was deprived of sight, hearing, and sense of smell; yet in spite of these great handicaps she has not only learnt to talk and to communicate her ideas but has achieved high academic distinction in letters, languages, and mathematics, as well as world-wide fame for her work among the deaf-blind. There were at the dinner representatives from St. Dunstan's and the National Institute for the Blind, Members of Parliament from both sides of the House, M. Georges Raverat, European director of the American Foundation for Overseas Blind, and among the other guests Maestro J. Rodrigo, a noted blind Spanish composer.

Sir Ian Fraser, welcoming Dr. Keller and Miss Thompson, said that as one born in South Africa he was glad that they were to visit that country to see the organizations there for

the care of the blind and the deaf. He was sure that Dr. Keller would bring help and encouragement to all handicapped people in the Union, whether European, Indian, or African. Sir Ian Fraser's speech was interpreted to Dr. Keller by the movements of Miss Thompson's right hand in her own, and it was clear from the lively expression of pleasure on Dr. Keller's face how much she appreciated what was said. It was easy to see how completely she was able to participate in all that went on, and this became even more evident when she rose to reply to Sir Ian's toast. Her words required Miss Thompson's sympathetic interpretation, but it was remarkable how distinct were some of them, especially those which were charged with feeling. In a most gracious speech she thanked her hosts and all those who had come to welcome her.

Miss Florence Horsbrugh, M.P., then proposed the health of Maestro Rodrigo, who lost his eyesight after diphtheria when he was 3 years old. She told the company how, in spite of this handicap, he had become a professor of music at Madrid University; how he had recently composed a concerto for guitar and orchestra in braille, which was later transcribed by a copyist.

REMUNERATION OF GENERAL PRACTITIONERS

LETTER FROM MINISTRY

The B.M.A. has received the following letter from the Ministry of Health:

"I am directed by the Minister of Health to refer to the meeting between representatives of the General Medical Services Committee and the Secretary of State for Scotland and himself on February 28, and to say that the Minister himself is already convinced, as a result of his first discussion with the representatives—and the Secretary of State for Scotland agrees—that there is a good prima facie case for their reviewing, as a matter of urgency, both the adequacy of the total remuneration of general practitioners and its distribution. They will look without delay at the best way of speeding up that review.

"A further meeting will be arranged before the end of April."

A new amino-acid, hitherto unknown in Nature, has just been prepared from human urine by C. E. Dent and his collaborators (*Nature*, February 24, p. 307). It is a normal constituent of urine, but in 4.8% of 459 healthy people who were tested it was present in an unusually large amount. Since in these people the blood level of the new amino-acid was not raised, the authors argue that the unusual urinary excretion must be due to a low renal threshold (defective reabsorption by the tubule from the glomerular filtrate), as is the explanation of the amino-aciduria of cystinuria and Fanconi syndrome. The new amino-acid is α -methyl- β -alanine (or β -amino isobutyric acid); β -alanine has already been frequently met in biochemical work, since it is one of the two components of the B-complex vitamin called pantothenic acid, and also occurs linked to another amino-acid, histidine, in the substances carnosine and anserine, which occur in fair quantity in muscle and which are at present of unknown function or significance. Whether α -methyl- β -alanine has any metabolic connexion with them remains to be seen. The new amino-acid was discovered through paper partition chromatography of urine, using phenol and collidine-lutidine as solvents. When the paper was sprayed with ninhydrin to detect the dispersed spots of different amino-acids an unknown "T-spot" was noted: the cause has proved to be α -methyl- β -alanine.