A working hypothesis on the basis of these mitosis counts is that in shock there are at least two inhibitory mechanisms affecting carbohydrate catabolism—one operating hormonally in the stage of glycolysis, and the other operating (possibly through the direct action of adenylyl compounds) in the oxidative cycle. Stoner, Green, and Threlfall (1950) have already provided some chemical evidence of an interference with carbohydrate utilization in shock. Here we are mainly concerned to show how these studies may help to explain one aspect of the complicated action of cortisone and related steroids.

If the antimitotic effect is, as seems very likely, exerted on other tissues, such as the reticulo-endothelial, and in other species than the mouse, this powerful action must play a part (see Green, 1950) in the anti-allergic, anti-inflammatory, and anti-repair effects which in the mouse and rabbit, and possibly man, are so prominent. That such effects are incidental and not in any way linked up with the known therapeutic effects of cortisone and related substances is most unlikely. That they explain all the therapeutic effects is still less likely, for in that case adrenaline or A.T.P. should, in appropriate dosage, have the therapeutic virtues of cortisone, which they manifestly have not. The antimitotic effect of adrenaline is, however, interesting in view of its rather similar effects to A.C.T.H. and cortisone in reducing or abolishing some hypersensitive tissue reactions. This particular property is possibly due to the discharge, in excess, of a cortical steroid through adrenaline stimulation via the pituitary-adrenal route.

Summary

Cortisone, injected or applied locally, is a powerful inhibitor of mitosis in the pre-prophase stage in the epidermis of the adult male mouse.

Two other substances possibly concerned in the shock reaction—adrenaline and adenosine—when injected in adequate dosage have a similar action.

It is shown that these findings give indirect support to the hypothesis that in the shock state there is a profound interference with carbohydrate metabolism.

We are indebted to the Medical Research Council Committee on A.C.T.H., Cortisone, and Related Substances for supplying the cortisone, and to Mr. R. F. Wrighton, of the Department of Social Medicine, Sheffield University, for his valuable statistical help.

REFERENCES

- Bullough, W. S. (1948). *Proc. roy. Soc. B*, **135**, 212.
 — (1949a). *J. exp. Biol.*, **26**, 83.
 — (1949b). *J. Endocrinol.*, **6**, 350.
 Green, H. N. (1950). *British Medical Journal*, **1**, 1165.
 — and Bullough, W. S. (1950). *Brit. J. exp. Path.*, **31**, 175.
 — and Stoner, H. B. (1950). *Biological Actions of the Adenine Nucleotides*. Lewis, London.
 Long, C. N. H. (1947). *Rec. Progr. Hormone Res.*, **1**, 99.
 — (1949). In A. K. Parpart's (Editor) *Chemistry and Physiology of Growth*, p. 266. Princeton University Press.
 Hughes, A. F. W. (1950). *Quart. J. micro. Sci.*, **91**, 251.
 Peters, R. A. (1948). *Proc. roy. Soc. Med.*, **41**, 781.
 Stoner, H. B., and Green, H. N. (1950). *Brit. J. exp. Path.*, **31**, 603.
 — and Threlfall, C. J. (1950). *Proc. 18th Int. Physiol. Congr.*, p. 473.

Fresh water aboard ship has always been something of a problem, and its magnitude can be gathered from the fact that a modern liner uses about 220 tons a day in the Tropics, and must provide storage for 3,000 tons, representing a great loss in cargo capacity. The P. & O. are now for the first time to make all their fresh water by continuous distillation of sea water. In their new *Himalaya* special evaporators have been designed for heating by steam taken from the turbines after it has done its useful work there.

SHOCK AND NEOPLASIA

BY

H. N. GREEN, M.A., M.D., M.Sc.

AND

M. SAVIGEAR, M.Sc.

(From the Department of Pathology, University of Sheffield)

It was shown (Green, 1949) that in the rat the growth of neoplasms from the time of transplantation was unaffected, or relatively unaffected, by repeated induction of the shock state, a procedure which powerfully inhibits mitosis in normal epidermis and probably many other tissues (Green and Bullough, 1950). This was unexpected, since the pronounced fall in oxygen consumption in the tissues of the shocked animal (Green and Stoner, 1950) and some retardation of general body growth might have been expected to retard neoplastic proliferation somewhat. Preliminary evidence was presented at the International Cancer Congress in Paris (Green, 1950) that, as might now be expected, the mitotic rate of the cells of the transplanted tumour was relatively unaffected by such a strong shock-inducing measure as hind-limb ischaemia or by the injection of A.C.T.H. Some further details are now presented together with an interesting extension of this inquiry. The general methods used have already been described (Bullough, 1948; Green and Bullough, 1950).

Tumour Mitosis and Natural Antimitotic Agents

Since then further investigations make it clear that such strong antimitotic agents as the shock state or injected cortisone (Green and Ghadially, 1951) do not affect mitosis in various transplanted tumours of mice and rats to anything like the same degree as is observed in the epithelium of normal skin.

In Table I some figures are abstracted from two typical experiments to show the effect and time-relationships of ischaemic shock and cortisone injection on the mitotic

TABLE I.—Effect of Ischaemic Shock and Cortisone on Epidermal Mitotic Rates. Average Number of Mitoses per 10 cm. Length of Ear Epidermis in Sections Cut at 7 μ . The Figures Given are for 5 Strong's CBA Male Mice, and in the Case of Cortisone for 5 Kreyberg's "White Label" Male Mice

Time of Day	Untreated Normal Mice	Ischaemic Shock. Removal of Limb Tourniquets at 11.30 a.m.	Cortisone, 0.2 mg. I.P. Injected at 12 noon
10.30 a.m. ..	31 $\pm$ 2.2	27 $\pm$ 2.9	—
12 noon ..	52 $\pm$ 4.1	14 $\pm$ 1.1	58.8 $\pm$ 14.2
1.30 p.m. ..	73 $\pm$ 1.2	0	9.2 $\pm$ 7.0
3 p.m. ..	44 $\pm$ 1.2	0	6.5 $\pm$ 3.0

rate in the skin of normal mice. In Table II the effect of similar treatments in tumour-bearing mice and rats on the mitotic rates of the tumours and of the epidermis in the same animal is illustrated. It has been found difficult in some instances to determine the effect of antimitotic agents on the skin of tumour-bearing mice, and the presence of a rapidly growing neoplasm may conceivably depress the rate of mitosis in normal epidermis. This point, however, is being investigated further, for, if true, it could be important. It will be readily seen, particularly if Tables I and II are compared, that the tumour mitotic rate in both the mouse and the rat is relatively unaffected as compared with the epidermal rate, either by ischaemic shock or by cortisone injection.

TABLE II.—*Effect of Ischaemic Shock and Cortisone on Epidermis and Tumour Mitotic Rates in the Same Tumour-bearing Animal. Average Number of Mitoses per 5 mm.² Tumour Section (5 μ) and 10 cm. of Ear Epidermis (7 μ) Taken at 3 p.m.*

Type	Tissue	Untreated Tumour Mice	Ischaemic Shock	Cortisone, 0.2 mg.
C3H male mice bearing breast carcinoma	Tumour Ear	(10 Mice) 46.0 $\pm$ 3.2 15.5 $\pm$ 4.8	(8 Mice) 36.6 $\pm$ 5.0 0	(1 Mouse) 51 0
Wistar rats bearing RD3 sarcoma	Tumour Ear	(8 Rats) 53.6 $\pm$ 1.8 70.0 $\pm$ 21.5	(8 Rats) 25.6 $\pm$ 2.6 0.1	—

In some experiments ischaemic shock in the mouse did not reduce the tumour mitotic rate at all, and the most powerful reduction was only by about half as compared with complete suppression in the epidermis.

It must be recognized, of course, that we are not comparing the malignant cell with its normal prototype—that is, skin carcinoma with skin epithelium. There is, however, collateral evidence from human studies with A.C.T.H. and cortisone which seem to us to indicate that neoplasms of the lymphoid tissues are not so sensitive as normal lymphoid tissue to the “involuting” effect of these substances. Though striking shrinkage of enlarged lymph nodes was observed in chronic lymphatic leukaemia, lymphosarcoma, etc. (Pearson and Eliel, 1950), this was usually transitory and may have been due partly to “involution” of normal hyperplastic lymphoid tissue. It is perhaps significant that a variety of neoplasms, both carcinomata and sarcomata, were completely unaffected by this treatment. There is probably a relationship between this lack of response of many human malignant neoplasms to cortisone and A.C.T.H. and the relatively weak or negligible anti-mitotic effect observed in our animal neoplasms. If it could be generally established, a negative might prove of greater interest than a positive effect by revealing an essential difference, amenable to experimental approach, between the normal and the malignant cell. One approach we are trying is to treat chemically induced carcinomata of the skin in mice with cortisone, as cortisone has a definite local antimitotic effect on normal mouse skin (Green and Ghadially, 1951).

Another and perhaps more practical approach was to find the response of the skin pretreated with a chemical carcinogen to these powerful antimitotic agents. Does such a local treatment make the epithelium refractory to such agents in whole or in part? We have now good evidence that it does. After a short period (not more than four paintings over eight days) of treatment with the powerful carcinogen 9:10-dimethyl-1:2-benzanthracene (D.M.B.), the mitotic rate of the ear epidermis usually rises quite substantially above the normal rate and above that of the opposite ear treated with the solvent only. The high rate is maintained for some days after cessation of the painting, but appears to decline with the gradual disappearance of the increased vascularity produced locally by the treatment. This increase in the mitotic rate is rarely, though it is occasionally, as great as after croton-oil painting (see Table III).*

From the present standpoint this stimulant effect of the carcinogen is by no means so important as the response of the treated cells to shock-inducing measures

TABLE III.—*Response to Antimitotic Agents of Ears Treated with 9:10-Dimethyl-1:2-Benzanthracene (0.5% in Acetone) and Croton Oil (1% in Acetone). Five Kreyberg's "White Label" Male Mice were Used in Each Case*

No. of Paintings	Average No. of Mitoses per 10 cm. of Ear Epidermis (7 μ)				
	Day of Count	Antimitotic Agent	Time of Day	L. Ear (Control)	R. Ear (Treated)
D.M.B.; 4 applications over 9 days	9th	Ischaemic shock tourniquets released at 12.30 p.m.	10.30 a.m. 4.30 p.m.	13.0 $\pm$ 4.0 0	31.2 $\pm$ 6.4 42.3 $\pm$ 6.8
D.M.B.; 4 applications over 8 days	8th	„ „	10.30 a.m. 4.30 p.m.	3.8 $\pm$ 1.2 0	29.6 $\pm$ 11.8 36.6 $\pm$ 13.1
D.M.B.; 4 applications over 8 days	8th	Cortisone (0.2 mg.) I.P. at 12 noon	12 noon 3 p.m.	58.8 $\pm$ 14.2 6.5 $\pm$ 3.0	54.6 $\pm$ 12.0 44.8 $\pm$ 10.0
Croton oil; 5 applications over 10 days	10th	Ischaemic shock tourniquets released at 11.30 a.m.	10.30 a.m. 3 p.m.	26 0	293 3

and physiologically equivalent procedures. In Table III examples are given of typical experiments. The carcinogen-treated ear, in contrast to the untreated opposite ear, showed only a slight response to the decisive antimitotic stimulus following the release of limb tourniquets (Green and Bullough, 1950) or following the injection of cortisone (Green and Ghadially, 1951). In striking contrast is the effect of shock on the hyperplastic epithelium induced by croton oil (Table III). Here the excessively high mitotic rate was almost completely suppressed, though not quite so quickly as was normal mitosis in the untreated opposite ear.

Discussion

Two most interesting conclusions can be drawn from this work at its present stage.

Even a relatively short exposure (in terms of tumour development) to a chemical carcinogen makes the skin epithelial cells refractory in some degree to the antimitotic action of certain natural agents. The skin is now relatively insensitive to the profound antimitotic effect of induced shock. Of the known factors concerned in this effect only cortisone and A.C.T.H. have been tested, and the carcinogen-treated skin was refractory to both. We are, however, at the moment somewhat uncertain of the biological potency of the sample of A.C.T.H. used. Adenosine and its compounds have not yet been tested, but, as their effects are in the present respect exactly like those of shock (Green and Bullough, 1950; Green and Ghadially, 1951), the same lack of response in the carcinogen-treated skin may be anticipated. It is also highly significant that the abnormally high rate of mitosis induced by croton oil is nearly as responsive to shock inhibition as is normal skin. The contrast here between the behaviour of the cell involved in the inflammatory reaction and the cell involved in the “precancerous” reaction is remarkable.

If the skin paintings are stopped, at any rate at an early stage, the refractory state to these agents inhibiting cell division gradually disappears. We do not know as yet if, and at what stage of treatment, the refractory state becomes permanent. That it does become permanent is highly probable, for in a considerable number of tests on three neoplasms (a breast carcinoma and two types of chemically induced sarcomata) the maximum decrease in the mitotic rate following a strong shock

*The experiments with croton oil were done in collaboration with Dr. W. S. Bullough.

stimulus was only a half. In many instances the malignant cell appeared to be completely refractory to such stimuli. Of course the importance of this finding depends on how far it reflects an essential difference between the malignant cell and its normal prototype. The effects of local carcinogen treatment do, however, strongly suggest that the resistance of the fully developed cancer cell is acquired very early in neoplasia and gradually becomes a permanent property of the cell.

Although such a resistance to normal growth-regulating agents has often been postulated as a property of the cancer cell, and in fact is vaguely included in one definition—that is, “autonomous new growth”—the property has never been precisely demonstrated experimentally. It may be that this work not only reveals the existence of such a property but offers an approach to an elucidation of its nature. If, as we think (Green and Bullough, 1950; Green and Ghadially, 1951), these antimitotic agents act on the normal cell by interfering at one or more stages in carbohydrate utilization, the work of the Sheffield Traumatic Shock Team on carbohydrate metabolism in shock may prove to have some bearing not only on the mechanism of this general bodily reaction but also, somewhat surprisingly, on the intimate changes in cell metabolism, which in themselves reflect, or constitute, the essential nature of neoplastic transformation.

Summary

A short period of painting of the ear with the carcinogen 9:10-dimethyl-1:2-benzanthracene renders the epidermis relatively refractory to the antimitotic action of shock or of cortisone.

Various transplanted malignant neoplasms are also relatively insensitive to the same stimuli.

Whether these findings have a bearing on the essential nature of the neoplastic change is briefly discussed.

We are indebted to the Medical Research Council Committee on A.C.T.H., Cortisone, and Related Substances for supplying the cortisone; to Mr. R. F. Wrighton, of the Department of Social Medicine, Sheffield University, for his valuable statistical help; and to Dr. W. S. Bullough for his expert collaboration in one experiment.

REFERENCES

- Bullough, W. S. (1948). *Proc. roy. Soc. B*, **135**, 212.
 Green, H. N. (1949). *Rep., Brit. Emp. Cancer Campgn.*, **27**, 172.
 — (1950). *Proc. Fifth Congrès International du Cancer*, Paris. *Acta Un. int. Cancer*, **7**. In press.
 — and Bullough, W. S. (1950). *Brit. J. exp. Path.*, **31**, 175.
 — and Ghadially, F. N. (1951). *British Medical Journal*, **1**, 496.
 — and Stoner, H. B. (1950). *Biological Actions of the Adenine Nucleotides*. Lewis, London.
 Pearson, O. H., and Eliel, L. P. (1950). *J. Amer. med. Ass.*, **144**, 1349.

How the honey-bee discovers the first, newly opened flowers of a crop was discussed in a paper read to the Royal Society on March 8 by Dr. C. G. Butler, of Rothamsted Experimental Station, Harpenden. It has been known for a long time that bees possess good colour vision, though they can see into the ultra-violet where the human eye is blind and are insensitive to red—whence, it is suggested, arises the scarcity of pure red flowers. More recently Professor Karl von Frisch has described the amazing dance a scout bee performs on returning to the hive to let its worker colleagues know where it has found a new group of open flowers with pollen and nectar ready. But how does the scout discover the new flowers in the first instance—by sight or smell? Dr. Butler reported that colour is the chief attractive factor, though a really strong perfume acts in the same way. The scent of an individual flower is, however, often weak, and it serves only to induce the honey-bee to enter the corolla once the colour has done its work. If the scent is masked, the bee hesitates to enter on the first or subsequent visits.

CHEMOTHERAPY IN CHOLERA

BY

S. C. LAHIRI, M.D.

Associate Professor of Medicine, National Medical Institute,
and Visiting Physician, Chittaranjan Hospital, Calcutta

Really effective chemotherapy in cholera has hitherto proved to be a baffling problem. However, good results have been claimed for a few of the newer compounds. Among these, sulphaguanidine has been extensively used for the past few years, but opinions regarding its efficacy in this disease have not been uniform (Chopra *et al.*, 1941; Carruthers, 1942; Lahiri, 1945, 1948). In another series of observations (Lahiri, 1949; Lahiri and Dutta, 1949) thiodiamine was tried. This is derived from the decomposition product of the bark of *Crataeva roxburghii* and also of *Moringa pterygosperma*. *In vitro* it prevented the growth of *Vibrio cholerae* (both Ogawa and Inaba types) even in a concentration of 0.5 mg. in 10 ml. of media. On clinical trial in 30 bacteriologically positive cases the stool became free of vibrio in three to five days in all the 23 cases systematically observed, but the death rate was not materially lowered.

Bhatnagar *et al.* (1948) had remarkably good results with another compound—a condensation product of sulphathiazole and formaldehyde. Out of 85 patients treated with this drug in a part of the Madras Presidency during an epidemic in 1947 only three died (3.53%); the case mortality rate had been about 60% in epidemics during the previous seven years. Saline transfusion was not given to any of these patients, and there was no nursing and general medical care or adjuvant treatment, though 57 patients (67.06%) were in a critical condition with anuria, marked dehydration, “semi-comatose condition, algidity, imperceptible pulse, and body cramps.” Stools became free from vibrio in all the cases within six days. The prophylactic value of this drug has also been emphasized. Bose (1949), in experiments *in vitro* with different condensation products of formaldehyde and sulphonamides, found that formosulphacetamide and formosulphabenzamide exerted better bactericidal effect on *V. cholerae* than formosulphathiazole and other similar compounds.

A clinical trial of some of the above compounds—for example, sulphaguanidine, formosulphathiazole, and formosulphacetamide—was undertaken in cholera cases admitted into the cholera ward of the Chittaranjan Hospital, Calcutta, during the last epidemic (1950). The findings are discussed below.

Clinical Investigation

The period of observation of the cases discussed here was from May to July, 1950. The total number of patients admitted during this period was 377, and there were 122 deaths (32.35%). Many of these patients were refugees in a bad state of nutrition and with lowered vitality. Consequently the death rate was high. A total of 268 patients were under clinical observation for the present study. All these patients presented typical symptoms of cholera, with profuse rice-water stools, vomiting, and dehydration with anuria, prostration, and high specific gravity of blood. At the time of admission 133 patients (49.62%) were in a very bad state of collapse, with imperceptible radial pulse and the specific gravity of blood varying between 1068 and 1072.