In acute female schizophrenia, thyroid function was usually increased, but was normal or subnormal in the chronic form.

There were no psychiatric reactions characteristic of thyroid abnormality.

The significance of the findings and the possibility of thyroid therapy have been discussed.

REFERENCES

- Asher, R. (1949). *British Medical Journal*, 2, 555.
Bowman, K. M., and Fry, C. G. (1925). *Arch. Neurol. Psychiat., Chicago*, 14, 819.
— Miller, E. R., Dailey, M. E., Simon, A., and Mayer, B. F. (1950). *J. nerv. ment. Dis.*, 112, 404.
Haigh, C. P., and Reiss, M. (1950). *Brit. J. Radiol.*, 23, 534.
Hoskins, R. G. (1946). *The Biology of Schizophrenia*. New York.
— and Sleeper, F. H. (1929). *Endocrinology*, 13, 245, 459.
— (1932). *Amer. J. med. Sci.*, 184, 158.
Means, J. H. (1948). *The Thyroid and Its Diseases*. Philadelphia.
Reiss, M., Haigh, C. P., Hemphill, R. E., Maggs, R., Reiss, J. M., and Smith, S. (1951). In press.
Simpson, S. L. (1949). *British Medical Journal*, 2, 759.

ACUTE UNILATERAL DACRYOADENTITIS

A REPORT OF FOUR PATIENTS

BY

PETER WILSON, F.R.C.S.Ed., D.O.M.S.

Ophthalmic Registrar, General Infirmary, Leeds

During the six months October, 1949, to March, 1950, four cases of acute dacryoadentitis were observed among the 3,622 new ophthalmic patients at the General Infirmary at Leeds. This incidence contrasts with the views of many writers who regard it as a rare condition, and several articles describe a single case which is reported by virtue of its rarity. Arlt (quoted by Schirmer, 1877) stated that he had never met a case. Pooley (1884) stated that the condition was rare and reported one instance which was considered to be metastatic in origin, though at the time there was a diphtheritic infection of the conjunctiva of the other eye. Galezowski (1886) also stressed its rarity, but stated that he had met five examples. Weeks (1911) reported the incidence to be 1 in 10,000 ophthalmic cases. Richardson (1942) was able to compile from the literature 37 cases of gonococcal origin and mentioned the causes of acute dacryoadentitis which had been reported.

The clinical features of the four patients with acute dacryoadentitis are described below, and points of particular interest are discussed.

Case 1

Five weeks before admission this patient, a girl aged 16, had had what was thought to be influenza, which lasted two weeks. She made an apparently complete recovery and was able to resume work. The day before admission a painful swelling appeared in the upper lid of the left eye and was accompanied by malaise and lack of appetite.

On examination the patient was rather lethargic. Her temperature was normal, pulse 80, respirations 18. There was no cough, and the chest condition seemed normal. Radiologically the paranasal sinuses appeared normal.

Ocular Condition.—A swelling was seen in the upper lid of the left eye, situated mainly at the outer aspect but extending to involve the whole lid and adjacent frontal region. The swelling was tense and tender; the eye was considerably proptosed, but there was only slight chemosis of the bulbar conjunctiva. Eye movements were almost

absent. A radiograph of the sinuses showed no evidence of infection.

Systemic administration of penicillin and sulphanilamide was accompanied by the instillation of penicillin drops and application of heat to the eye. After a few days the swelling became less tense and subsided, allowing palpation of a firm lump in the upper lid which was felt to extend forwards from under the superior orbital margin. The general condition improved as the swelling diminished. When the patient was seen after four weeks no lump could be palpated and the chemosis had disappeared, but there was still slight ptosis and some weakness of the left superior rectus.

Case 2

Three weeks before attending hospital this patient, a boy aged 10, had had what was thought to be tonsillitis, for which he was confined to bed for one week. He made an apparently complete recovery and was able to resume school. Four days before admission a painful swelling appeared in the right upper lid, accompanied by considerable malaise and loss of appetite. One intramuscular injection of penicillin had been given by his own doctor.

On examination his general state was one of languor and apathy. His temperature was normal and pulse 80. The patient had a loose cough, and rales and rhonchi were heard in the chest. Radiological examination revealed some haziness of the paranasal sinuses.

Ocular Condition.—The right upper lid was swollen. This swelling was mainly in the outer part, but involved the whole lid and extended into the frontal region. It was tense and tender and there was complete ptosis. The eyeball was considerably proptosed, the conjunctiva was chemotic and bulging between the lids. Movements of the eye were almost absent. Acute tenderness was felt on palpation over the right malar bone. Both fundi appeared normal. Radiologically there was thickening of the mucosa of the frontal sinuses; this was not considered an adequate cause for the swelling of the lid.

Systemic administration of penicillin and "sulphatriad" was accompanied by instillation of penicillin drops and the application of heat to the eye. Next day the pulse rate had increased to 90 though the temperature remained normal, so an incision was made over the swelling and forceps were inserted. No pus appeared until the orbital septum had been pierced, when about 0.5 ml. of creamy pus was evacuated. During the next seven days the swelling and proptosis subsided and a lump could be palpated in the upper lid. This lump was felt to extend forwards from under the superior orbital margin. When the patient was seen five weeks later there was a complete recovery of both the general and the local condition.

Case 3

A girl aged 16 had recently had an attack of what was considered to be influenza. She was in bed for one week and then returned to school for two weeks. Four days before she attended hospital a painful swelling appeared in the upper lid of the left eye, mainly at the lateral aspect. The swelling increased during the next few days and was accompanied by ptosis and chemosis.

On examination her general condition was one of languor and malaise. Her temperature and pulse were normal.

Ocular Condition.—The left eye showed complete ptosis, with a tender tense swelling occupying the upper lid, mainly in its outer part and extending into the forehead. There was considerable chemosis, mainly of the lateral bulbar conjunctiva. Slight proptosis was present and movement of the globe was limited in all directions. Radiological examination revealed no evidence of infection of the paranasal sinuses.

Systemic administration of penicillin and sulphatriad was accompanied by the instillation of penicillin drops and the application of heat to the eye. The swelling subsided gradually, and after four to five days a lump became

palpable in the lateral part of the upper lid, emerging from under the superior orbital margin. Apart from some nausea from sulphatriad, the general condition improved as the swelling diminished. Two weeks later the proptosis had disappeared, but there was some diplopia on looking to the left. The diplopia was the last to clear up and was still noticeable after five months.

Case 4

A boy aged 14 began to complain of headache and feverishness six days before attending hospital. Three days later he had a painful swelling of the upper lid of the right eye and, at the same time, a sharp pain in the left side of the chest. He was seen by his doctor, who was doubtful whether to diagnose right frontal sinusitis or left lower lobe pneumonia. His temperature exceeded 100° F. (37.8° C.) during the next few days.

On examination signs of consolidation with a small effusion were found at the left base of the chest. A blood count showed 8,600 white cells with 81% polymorphs. There was

a swelling of the right upper lid, maximal at the lateral aspect but involving the whole lid and extending into the frontal region (see illustration). The swelling was tense and tender and was associated with complete ptosis, some proptosis, and chemosis. Movements of the right eye were considerably diminished in all directions but mainly upwards and outwards. The right malar bone was tender on palpation.

Systemic administration of penicillin and sulphadimidine was accompanied by the instillation of penicillin drops and by the application of heat to the eye. Aspiration of the pleural effusion was unsuccessful. During the following week complete resolution occurred at the left base, but no fluid was found, though transient inflammatory changes appeared at the right base, accompanied by fever. With treatment the swelling in the right lid gradually subsided during the course of seven days, and a lump became palpable in the lateral part of the upper lid. Three weeks

later there was only slight ptosis and slight thickening of the outer part of the lid.

Discussion

The diagnosis of infection of the lacrimal gland depends upon the palpation of a lump in the lateral part of the upper lid, extending forwards into the lid from under the outer aspect of the supraorbital margin; therefore the posterior limit of the lump cannot be determined, though the anterior edge is easily palpated through the thickness of the lid. During the initial stages of the infection the tenseness and tenderness of the region prevent any useful palpation, but the condition may be suggested by a swelling which is maximal at the lateral aspect of the lid. Such a swelling must be distinguished from other inflammatory swellings of the upper lid.

The most important conditions which may simulate acute dacryoadenitis are orbital cellulitis, osteitis, and acute inflammation of the Meibomian gland. Schirmer (1877) stated that many instances described as infection of the gland were really infections of the cellular tissues in the vicinity.

Orbital cellulitis arising from metastatic infection, from spreading thrombophlebitis, or by direct spread from a paranasal sinus usually produces a swelling which is diffuse throughout the length of the lid or maximal at the nasal aspect. Ethmoidal mucocoele typically produces a swelling which is at the medial aspect of the lid and displaces the eye laterally and downwards (Thomson and Negus, 1948).

Osteitis of the frontal bone may produce a swelling of the upper lid, but radiological changes in the bone would be expected, and such a subsiding osteitis would be unlikely to leave a lump at the site of the lacrimal gland. Benford and Brunner (1947), however, give details of a patient with a polypoid mass of sinus mucosa which had herniated through a fistula in the frontal bone.

Acute infection of a Meibomian gland may cause a swelling in the lid and on subsiding may leave a firm lump; the lump is likely to be confined to the lid and not to extend under the orbital margin; moreover, eye movements are not likely to be affected.

Some interesting features are presented by these four patients. In each there was an infection of the respiratory tract. The first three gave a history of such an infection two to three weeks before attending hospital and apparently had made complete recovery. Two of these febrile attacks had been ascribed to influenza and one to tonsillitis. The fourth patient apparently acquired infection of the lung and of the lacrimal gland simultaneously. Why infection of the lacrimal gland should have this association with infection of the respiratory tract is difficult to explain. It may be that the gland infection follows transport by the blood stream of material from a site of infection elsewhere in the body, and a common site for such an infection is the respiratory tract.

General symptoms were severe in the early stages. By the time the patients were seen in the eye department, all had had penicillin systemically and their temperature was normal; nevertheless there was considerable malaise and loss of appetite. Recovery of the general condition seemed to run concurrently with subsidence of the swelling, in contrast with the type of infection described by Toulant, Benhamou, and Foissin

(1947) in which the local condition cleared up in a matter of days but the general malaise was considerably prolonged.

Acute tenderness caused by pressure over the zygoma was noticed in Case 2, and thereafter was elicited in Cases 3 and 4. It was not noticed in Case 1, perhaps because that area was not palpated, but in Case 2 the provisional diagnosis of osteitis of the zygoma was made, owing to the situation and degree of the tenderness. This point was about 1 in. (2.5 cm.) behind and below the external canthus—a point quite distinct from that found in tenderness due to enlarged preauricular gland—and presents an interesting feature for speculation. It may be that the sensory nerve supply for this area of skin is derived from the zygomatic branch of the maxillary nerve, and as a result of the communication between the lacrimal and zygomatic nerves tenderness in this region reached a high degree. No reference to this phenomenon could be found in the literature, yet it appears to be characteristic of acute dacryoadenitis at an early stage, when swelling of the lid is too tense to permit palpation of a lump.

Ptosis was present on each occasion, and owing to the situation of the swelling, which was maximal at the lateral aspect of the lid, a sinuous outline was given to the lid margin. This appearance has been noted by Richardson (1942) and by Rosen (1944).

Restriction of eye movement was found in all patients in the early stages of the gland infection; in two it amounted to almost complete immobility of the eye, and a notable feature was the comparative slowness in recovery of full eye movements.

Proptosis was a pronounced feature in one instance. This patient had had the most severe general reaction, and one might visualize an abscess in the gland which had extended backwards to cause an orbital cellulitis.

Chemosis was seen in all patients, chiefly affecting the temporal bulbar conjunctiva. In two it caused bulging of the conjunctiva between the lid margins.

In conclusion, it may be stated that acute dacryoadenitis is not uncommon in some districts and at certain times of the year. Schmidt (quoted by Schirmer, 1877), stated that the condition was not uncommon; Wright and Nayar (1937) described three cases which seemed to be caused by *Diplobacillus morax-axenfeld*. Epidemics have been reported from time to time (Galezowski, 1875; Groenouw, 1889 (quoted by Nicolau, 1933); Nicolau, 1933; Toulant, Benhamou, and Foissin, 1947). The seasonal incidence is a matter of some interest; Nicolau described 10 cases which occurred during the winter months, and, on investigating 81 instances previously reported, she found an increase in numbers during the months of October to March, those months in which respiratory infection is most common. Respiratory infection may be a frequent cause of unilateral acute dacryoadenitis.

It appears that there may be a latent interval of two to three weeks during which the patient feels quite well before orbital symptoms become manifest; only careful history-taking will reveal that there has been a period of headache, malaise, sore throat, and feverishness. Fralick (1935) makes the following comment: "Seldom is the practising ophthalmologist called upon to treat conditions affecting the lacrimal gland. Consequently, little is written in the standard textbooks on the subject of acute inflammation of this structure. It is therefore conceivable that a few cases may leave our offices

improperly diagnosed because of our unfamiliarity with this particular group."

Summary

Four patients with acute dacryoadenitis have been seen during a six-months period. The clinical features are described and discussed. Attention is drawn to the following points: (1) the rarity of acute dacryoadenitis, (2) its association with respiratory infection, (3) its seasonal incidence, and (4) an early diagnostic sign of acute dacryoadenitis consisting of tenderness over the malar bone.

I would like to thank Dr. Maxwell Telling for much valuable criticism and for access to Case 4, and Mr. John Foster and Mr. George Black for their help and for allowing me to describe their patients.

BIBLIOGRAPHY

- Benford, M. C., and Brunner, H. (1947). *Amer. J. Ophthalm.*, **30**, 297.
 Boley, J. P., and Taylor, R. E. (1940). *J. Canad. med. Ass.*, **43**, 66.
 Byers, W. G. M. (1912). *Trans. Amer. ophthalm. Soc.*, **13**, 135.
 Foster, M. L. (1925). *Ibid.*, **23**, 50.
 Fralick, F. B. (1935). *Amer. J. Ophthalm.*, **18**, 19.
 Galezowski, X. (1875). *Recu. Ophthalm.*, **2**, 270.
 — (1886). *Ibid.*, **8**, 415.
 — (1892). *Ibid.*, **14**, 594.
 Nicolau (1933). *Ann. Oculist., Paris.*, **170**, 772.
 Pooley, T. R. (1884). *Trans. Amer. ophthalm. Soc.*, **3**, 729.
 Richardson, J. M. (1942). *Arch. Ophthalm.*, **Chicago**, **28**, 93.
 Rosen, E. (1944). *Amer. J. Ophthalm.*, **27**, 276.
 Schirmer, R. (1877). In A. Graefe and E. T. Saemisch's *Handbuch der Gesamten Augenheilkunde*, **7**, 1. Engelmann, Leipzig.
 Thomson, StC., and Negus, V. E. (1948). *Diseases of the Nose and Throat*, 5th ed., p. 297. Cassell, London.
 Toulant, Benhamou, and Foissin (1947). *Bull. Mém. Soc. franç. Ophthalm.*, **60**, 23.
 Weeks, J. E. (1911). *Treatise on Diseases of the Eye*, p. 190. Kimpton, London.
 Wright, R. E., and Nayar, K. K. (1937). *Brit. J. Ophthalm.*, **21**, 367.

BORNHOLM DISEASE

BY

T. GEFFEN, M.D., M.R.C.P.

Late Medical Registrar, Hampstead General and North-west London Hospital

Outbreaks of Bornholm disease have been described in this country at infrequent intervals since the work of Pickles (1933) in Yorkshire. The subject was recently reviewed by Hopkins (1950), who reported a further series of cases; and the possible connexion between this disease and the Coxsackie group of viruses was discussed by Findlay and Howard (1950). Very shortly after the publication of the latter two papers a small group of patients attended this hospital with clinical histories corresponding closely to those reported in previous descriptions of Bornholm disease. Dr. G. M. Findlay very kindly attempted to isolate Coxsackie virus from the nasal washings of some of the patients and to carry out the serological antibody reactions on them from the earliest stages of the outbreak. The clinical histories of these patients are briefly described, together with a summary of Dr. Findlay's findings.

Case Reports

Case 1.—A girl aged 18 first noticed mid-thoracic pain and pain behind the right shoulder on May 28, 1950. It was worse on inspiration but was not severe, and she did not mention her symptoms until three days later, when two