A Case of Large Laryngeal Diverticula

Large laryngeal diverticula, although a normal condition in certain primates, are rare in man. The following case is of interest.

Case Report

A driver in the R.A.S.C., aged 24, reported at the British Military Hospital, Kuala Lumpur, Malaya, on March 13, 1950, with a history of a swelling on the left side of the neck of 12 months' duration. This was symptomless until two weeks earlier, when increasing hoarseness developed.

On examination a soft fluctuant swelling was found completely filling the left submaxillary triangle, in which air crepitus could be elicited. A laryngeal diverticulum was suspected, and the diagnosis was confirmed by asking the patient to hold his nose and blow; this caused the swelling to distend, filling most of the anterior triangle of the neck. It could then be emptied by pressure, with an audible squeak from the back of the throat. Examination of the right side revealed a similar but much smaller swelling, of which the patient was unaware. X-ray examination after inflation

Radiograph showing the left air sac.

showed air sacs on each side of the thyroid cartilage—large on the left side, small on the right (see illustration).

Laryngoscopy revealed an intralaryngeal extension of the left pouch emerging from the ventricular fossa, which inflated with each expiration and completely filled the larynx. It collapsed completely during inspiration; this explained the hoarseness and stridor which were noticeable when the patient was lying down.

Operation.—On March 22 a large left laryngeal diverticulum was removed by dissection through a collar incision in the neck. It emerged through the thyrohyoid membrane at the upper border of the thyroid cartilage at the site where the superior laryngeal vessels entered. There was a small pouch-like extension into the larynx.

The neck of the diverticulum was dealt with as for a hernial sac. The sac on the right side, which was small and symptomless, was not touched. Microscopical examination of the sac showed it to be lined with ordinary respiratory epithelium. Convalescence was uneventful, the voice returning to normal almost immediately.

I am indebted to Lieutenant-Colonel D. T. Swift, Officer Commanding British Military Hospital, Kuala Lumpur, for

permission to report this case, and to Brigadier J. C. Collins, D.M.S., G.H.Q., FARELF, for permission to forward for publication.

A. J. CLYNE, M.B., B.S., F.R.C.S.,
Lieutenant-Colonel, R.A.M.C.

Actinomycosis in a Child Aged 6

Swellings of the cheek in children are an every-day problem to doctors, and as cases of actinomycosis are rarely found under the age of 10 (Cope, 1938) the following case is reported so that the possibility of this diagnosis may be brought to mind when a chronic inflammatory condition of unknown aetiology presents in a child.

Case Report

The patient, a boy aged 6, was sent to hospital in October, 1950, by the school medical officer because of a so-called "cyst" of his left cheek; this was diagnosed clinically as " ? eosinophil granuloma." The mother stated that the lesion began in June, 1950, as a small dark blister about the size of a pin-head, followed by an indurated swelling which did not respond to fomentos. He had had tonsillectomy in 1949, scarlet fever in 1946, and measles in 1947. He was a city child with no history of contact with animals, and apart from a holiday at the sea in 1950 he plays in his local playground and the street of his home.

On examination a hard firm swelling 2 cm. in diameter and 0.5 cm. in depth was seen on his left cheek (see illustration). The cervical and tonsillar lymph nodes were not enlarged. The first and second left molars showed dental caries. The swelling was aspirated and about 3 c.mm. of serous-like material sent to the laboratory for examination, " ? acid-fast infection."

Photograph showing hard firm swelling on left cheek.

Laboratory Findings.—A small amount of serous-like material was received in a glass bottle, and a few bright yellow granules were observed. Direct examination showed Gram-positive branching filaments with round and clubbed ends, non-acid-fast, on which a diagnosis was made. No other organisms were cultured from this material.

Swabs taken from the buccal mucous membrane, teeth, and tonsils produced no evidence of mycelial forms on direct examination or on culture.

The patient was given 1,000,000 units of penicillin daily, and at the end of three weeks the swelling had practically disappeared. He had dental caries in the region of the buccal side of the swelling, and although the organism was not found in this area it may have been the portal of entry of the original infection (Lord and Trevett, 1936; Slack, 1942; Cope, 1949).

Our thanks are due to Mr. J. E. Andrews, Southern Group Laboratory, for the photograph, and to Mr. A. Cooper for his technical assistance.

L. R. A. WELLS, M.R.C.S., L.R.C.P.
Surgeon.

JANE M. FULLERTON, M.B., D.P.H.
Pathologist.

St. Olave's Hospital, London, S.E.16.

REFERENCES

- Cope, Z. (1938). *Actinomycosis*. Oxford Univ. Press, London.
— (1949). *British Medical Journal*, 2, 1311.
Lord, F. T., and Trevett, L. D. (1936). *J. infect. Dis.*, 58, 115.
Slack, J. (1942). *J. Bact.*, 43, 193.