REFRESHER COURSE FOR GENERAL PRACTITIONERS

ALLERGY

BY

C. J. C. BRITTON, M.D., D.P.H.

Physician in Charge of Department of Allergy, Prince of Wales's General Hospital, London, N.15

The purpose of this article is to describe a practical approach to the investigation and treatment of allergic diseases, which can be followed by the general practitioner either by himself or in conjunction with a specialist. As in almost every case of allergy a whole host of factors may be of aetiological importance, investigation and treatment must cover them all if the best results are to be obtained. Too often in the past, cases have been treated by one method only, to the exclusion of all others, and this has led to pessimism and neglect. This is also true of the present time. In no disease is this defeatist outlook so marked as in asthma, and yet I know of few chronic disabling conditions in which such excellent results can be obtained from a properly planned regime.

Allergy may be defined as a state of exaggerated susceptibility to various foreign substances or physical agents that are harmless to the great majority of normal individuals. The reaction is of similar type in the same individual for each substance, generally appearing after minute dosage, and differing from any toxic action the substance might have in large dosage.

At present there is no completely satisfying theory for the cause of allergic manifestations, but the greatest bulk of evidence supports the so-called histamine theory, and it is the one of most practical help. This theory holds that the actual allergic reaction is due to liberation of histamine from the cells in the particular tissue. This liberation of histamine is due to the damage caused to the cells by the allergic antigen-antibody union taking place in or on the cells rather than harmlessly in the circulating blood stream as in a simple immunity reaction. There is considerable evidence both for and against this theory, but little would be gained by discussing this matter further.

The liberation of the histamine has two important physiological actions: first, it causes increased vascular permeability so that fluid is poured out into the tissues (for example, as in urticaria or angioneurotic oedema); and, secondly, it causes spasm of smooth muscle (for example, as in asthma). In asthma there is also swelling of the mucosa owing to the increased vascular permeability.

The type of allergic reaction, depending on the "shock tissue," is usually fairly constant in any person; thus one individual may react with bronchial spasm; another with urticaria, a third with eczema, and a fourth with hay-fever, but in all cases there will be evidence of increased capillary permeability and irritation. Certain types of allergen tend to be more likely to cause certain types of reaction. Thus sensitivity to foods tends to cause urticaria and eczema; to pollen, hay-fever; and to dust and animal danders, asthma and vasomotor rhinitis. Again, certain reactions are more likely in certain age groups; for instance, allergic eczema is more common below the age of 5: after this age the child tends to replace his eczema by asthma.

Causes of Allergy

The most important predisposing cause is heredity. Whereas in any group of supposedly normal individuals approximately 4-5% give a history of some type of allergic manifestation within the last two generations, 50-70% of allergic patients have an allergic family history. If both sides of the family are affected the children are much more likely to show allergic manifestations and to show them at an earlier age than if only one branch is affected. Another predisposing cause, at least in asthma, is a severe respiratory illness—for example, whooping-cough or pneumonia.

The precipitating causes of the actual attack are manifold and are of the greatest importance, as it is at this stage that we should make our most fundamental attack on the allergic disease. The causes may be specific (extrinsic allergy) or non-specific (intrinsic allergy), but almost always both these elements are involved. The extrinsic factors, or specific sensitizing substances, may be classified by their portal of entry into inhalants (such as dust, feathers, grass pollen), ingestants (such as egg, cheese, or fish), injectants (such as horse serum and other biological products), and contactants (such as primula leaves or rubber). The intrinsic, or non-specific, factors may be classified as psychic, climatic, dietary, endocrine, physical, chemical, and mechanical.

Practical Approach

To explain my method of approach to an allergic case I will deal in detail with asthma and later give consideration to any special factors in other types of allergy. However, the principles involved are the same in all cases. Nothing that I advise for the treatment of these cases is beyond the competence of the general practitioner, though he may find it very convenient to call in the aid of an allergist in the initial stages. The first essential is to take a very full history. Time given to this is well spent and will make it possible to assess the relative importance of the different factors. The first question I ask is whether there is a hereditary factor, mentioning specifically any history of asthma, hay-fever, stuffy noses, chronic catarrh, sneezing attacks, irritable skin conditions, or migraine. Next, I ask for the method of onset, the age at which the condition first appeared, and for any history of respiratory illness or of a severe shock. I ask the patient what he has found precipitates an attack—for example, exercise, excitement, dust, pollen, or contact with animals—and for the periodicity, whether winter or summer, night or early morning.

With the answers to these questions before me, I plan my skin tests. These are carried out by the intradermal method for most inhalants with the adult patient, the scratch or prick method for ingestants and in young children, and the patch method for contactants. As a routine, I include house dust, feathers, cat, dog, and

horse danders, orris root, grass, tree, shrub, and flower group pollens, and some 30 different foods, but especially chocolate, egg-white, cheese, fish, and wheat.

While allowing these tests time, usually 10–15 minutes, to show their reactions, if any, I discuss with the patient the whole plan of the disease and its causes. This discussion is of fundamental importance, as unless one can explain satisfactorily to the patient how the attack is caused his co-operation is unlikely, and without such co-operation satisfactory results cannot be expected. The explanation must be in simple terms so that each point is clearly grasped by the patient. In some cases this may mean much repetition. The general explanation I give is as follows—modified, of course, according to the history, the type of allergy, and the appearance of any reactions to the skin tests as the explanation is proceeding.

General Explanation

I explain that most cases of asthma are due to a greater or lesser extent to sensitivity to certain substances inhaled, ingested, or contacted, and that this tendency to sensitivity is inherited, although the ancestor need not necessarily have had the same allergic disease or sensitivity to the same substance as the patient. I then proceed as follows: "Each thing to which you are sensitive may be illustrated by comparing it to a brick in a wall. With every contact a 'brick' is formed and the 'bricks' pile up until they reach a certain level called the threshold. If this threshold is passed the attack will begin. The size of the 'brick' depends on the degree of contact. Thus supposing you are sensitive to household dust and feathers, and suppose you sleep on a feather pillow; then if you lie quietly the feather 'brick' will be a small one, if you are restless it will be bigger; similarly with the dust from the bedding. These two factors, textile dust and feather dust, are the commonest allergic factors in asthma and explain why most attacks come on at night or in the early morning."

The patient will probably ask, "Why is it that I have slept on feather pillows or a feather bed for years and yet I get my attacks only once in every two or three weeks or months?" The answer to this is often simple. "It means that your sensitivity to dust and feathers is not great enough to reach the threshold by itself, but other factors are needed before the 'bricks' will be brought up to this level. These other factors may be contact on the occasion of the attack with other things to which you are sensitive, as, for example, horses, pollens, foods, and so on. It is more likely, however, that these other factors will be non-specific or intrinsic ones. Probably the most important of these is the psychological factor—any upset, worry, shock, overtiredness, excitement, and so forth may act as a 'brick.' Another important factor is the climatic one: sudden changes of temperature, as when getting out of a warm bed into a cold room in the morning or going from the cold into a warm stuffy atmosphere, cold east winds, cold damp conditions rather than cold and dry conditions, sudden changes of barometric pressure—all these can act as non-specific physical irritants and thus act as a 'brick' in our illustration. Again, respiratory infections are potent irritants. The bacterial infection in the bronchial tree causes both oedema and irritation, thus aggravating any potential swelling or spasm due to a specific sensitivity."

"Other mechanical irritants are such things as fumes from the boiler, insecticides, soap flakes, etc.—these irritate everybody to a certain extent but may be sufficient to precipitate an attack in an allergic patient. Finally, foodstuffs can act as 'bricks.' You may be actually sensitive to a food as shown by a positive skin reaction, but very often the food reacts in a non-specific way. Thus any overloading of the stomach with rich or heavy food, especially at night, will

often cause a reflex from the stomach to the bronchioles and act as a further non-specific 'brick.' Any attack will thus be due to a combination of these different factors—one attack will be due to contact with dust and feathers alone; the next to dust, feathers, and some psychological upset; the next to dust, worry, and a respiratory infection, and so on."

Skin Testing

As the specific factors are the most easily controlled, the importance of adequate skin testing is obvious. These skin tests are now examined for reactions. A good reaction should show a swelling about 1 cm. in diameter, and pseudopodia are present in cases with marked sensitivity. The site of the reaction often irritates, and it is helpful to allow the patient himself to pick out the positive reactions and to explain to him that what he is seeing on his arm is what actually happens in the bronchioles with an attack of asthma—namely, swelling of the mucosa, tending to block the bronchioles, and spasm of muscle round the bronchioles due to the irritation. It is the attention to such little points as these that brings home to the patient the mechanism of his attacks, and with the knowledge of the how and why of an attack he often loses the fear of the attack. This loss of fear in its turn results favourably on the attack itself.

The next step is to discuss with the patient in a practical and common-sense manner the avoidance of the things to which he may be shown to be sensitive by the skin reactions. We will suppose that the patient is sensitive to dust and feathers, as these are the most common. First, an explanation about "dust." This dust is that formed by disintegration of textiles, bedding, and the like. The dust from the road is only a mechanical irritant. The textile or household dust is not, as has been stated by certain workers, a simple mixture of wool, cotton, and other textiles, but is a disintegration, probably oxidation, product of these substances. An experiment which I have carried out makes this clear. If a skin-testing extract be made of a new carpet or blanket and an intradermal skin test is carried out on a known dust-sensitive patient, no reaction will occur. If, however, this carpet or blanket be kept say in a glass case but exposed to sunlight for a year or more, an extract now made will give a significant reaction in the dust-sensitive patient. The recent work of Rimington and Maunsell on the preparation of dust extracts points in the same direction.

Avoidance of Dust and Feathers

Directions are now given to help the patient to avoid dust and feathers. This advice must be practical and attainable by the patient. For this purpose I am very strict about the conditions in the patient's bedroom, the place where, after all, he has his most intimate contact with textile dust and probably feathers; the place where he spends over one-third of his life; and also the place where, by and large, he gets most of his attacks. I know of one allergy clinic where those patients who are found sensitive to dust are given a foolscap sheet of instructions. Only two of these need be mentioned: first, all rugs and carpets must be removed from the whole house; and, secondly, all wallpaper should be taken down throughout the whole house and be replaced by paint or distemper. Such ideals, of course, are impossible

to attain by most patients and the common reaction to such advice is, as I have seen, to give up hope or else to regard the whole procedure as ridiculous.

It is my contention that if the bedroom is strictly dealt with, the rather scanty contacts in the rest of the house can be neglected, apart from such obvious precautions as to use a vacuum cleaner where possible and to dampen floors before sweeping. I insist, therefore, that for the rest of the patient's life the bedroom should be as bare as a hospital ward—no bookcases, toy boxes, bric-à-brac, or settees. There should be only a dressing-table, the bed or beds, side tables, a wardrobe, and possibly a tallboy. On the floor, linoleum or stained boards are best, with washable mats which should be washed three or four times a year. If a fitted carpet is present I do not, at first any rate, insist on its removal so long as it is vacuum-cleaned daily.

Next the bed or beds are discussed. Eiderdowns should be removed for all time and replaced by extra blankets or, if luxury is required, by an electric blanket. The blankets should be washed at least once a year and sheets turned back well over the blankets, and, in the case of a child, pinned down if he is restless. This helps to prevent the blanket dust reaching the nose. Sponge rubber pillows are preferable, but good wool or flock ones covered with jaconet or plastic under the pillowcase are the next best variety. The best mattress is a sponge rubber one, but most good mattresses produce little dust if well vacuumed. It must be remembered that all feathers must be removed from the bedroom in feather-sensitive cases. I have often found that the wife has retained her feather pillow while the affected husband has a rubber one.

The patient's frequent inability for logical thought stresses the need for a detailed discussion of the methods necessary for the removal of dangerous allergens. Once again I would point out that common sense is needed. Nothing could be worse than to convert a patient into a fussy, introspective person who flies if he sees a feather cushion or a dog in a friend's house. If the patient's own bedroom is free from allergens, contacts elsewhere can, within reason, be neglected. Instructions with regard to contact with other allergens should be given in the same detailed, practical, and common-sense manner.

Non-specific Factors

Now the non-specific factors should be discussed in more detail. With regard to the psychological factor, the explanation given above helps to combat the fear of the condition. If the patient is a child, and the parents can afford it, I insist that the child should go to a boarding school, as the discipline of the school and the example of other children are most helpful. Also the child gets relief from a mother or father who aggravates things by over-protection. If the parents cannot afford a boarding school, I assure them that it is not essential, but I explain the importance of not fussing over the child. The activities of the child should not be curtailed, although exposure to bitter east winds is better avoided.

This brings us to a consideration of climate. Let us face the fact that there is little difference in climate from this point of view in Great Britain, and of the cases I see almost as many come from the south coast as from the midlands and industrial towns. As a general principle, low-lying damp areas should be avoided, but the contacts with irritants in the home are usually of more

importance than the comparatively slight differences in climate met with in Great Britain. It is difficult to avoid respiratory infections, but I find it helpful if the patient has a cold to give, each night before going to bed during the period of the cold, a prophylactic dose of an anti-spasmodic medicine. As mentioned above, heavy meals and rich foods should be avoided, especially at night. With regard to symptomatic treatment of asthma, I would refer you to an earlier article on asthma in this series by Dr. Bruce Pearson (*British Medical Journal*, 1950, 1, 1311).

Where sensitivity to dust or pollens is present, or in some cases of other sensitivities in which it is impossible to avoid the allergen, desensitization should be carried out, and solutions for this purpose, with full instructions, can be obtained from several commercial firms.

Small skin-testing outfits for the general practitioner can also be obtained if the practitioner wishes to carry out his own testing, but the interpretation of such tests with an imperfect technique may not be easy. Specific desensitization when properly carried out as an adjunct to control of contacts and intrinsic factors as detailed above gives excellent results. The period of relief obtained by such desensitization varies from case to case, but is likely to be prolonged where the specific sensitivities are great and where the control of contact with the allergens is adequate. When the sensitivity is marked it is, of course, important to build up the desensitization gradually, by beginning with a weak solution and following this with one or more stronger vaccines as may be required.

Hay-fever

This condition is due to sensitivity to grass, tree, shrub, or flower pollen, but it is surprising how often sensitivity to dust and feathers or animal danders is also present. It is well known if hay-fever is untreated some 50–75% of cases become asthmatic after a varying number of years. It would seem that this happens as the sensitivity to dust increases. One of the disadvantages of the anti-histamine drugs now used for the symptomatic treatment of hay-fever is that with them the conversion rate from hay-fever to asthma appears to be much higher and to occur much more rapidly. This has been my own experience and has been reported in the United States of America. It would seem that the antihistamine decongests the nose and allows the pollen to penetrate deeply into the lung. For this reason I do not recommend treatment of hay-fever by antihistamines alone, though they are of great value in "topping off" cases in which desensitization therapy gives incomplete relief. They are also useful as palliatives while desensitization is proceeding or until it can be begun.

Better results are obtained with desensitization if the desensitizing solution is made up from the range of pollens to which the patient is sensitive rather than from a single pollen or group of pollens. With hay-fever it is usually necessary for courses of injections to be given for three or more consecutive years, with progressively stronger solutions. It is particularly important to stress the necessity for persevering with the treatment; if this is not done a patient who is so much improved after one course of injections may neglect to continue with a stronger solution in the following year, with the result that he relapses in the future and loses the ground already gained.

When antihistamine drugs are used for symptomatic relief certain precautions should be taken. All these drugs cause side-reactions in a certain number of cases, and these side-reactions may be unpleasant or even dangerous. With all these drugs, therefore, it is important to begin with a small dose. The first dose should not be given at a time when the patient will be called upon to perform skilled movements depending on acts of judgment—for example, driving a car. For an adult half a tablet or capsule should be the initial dose. If this is not followed by side-reactions or does not give relief, then a whole tablet should be taken on the next occasion, and the dose can be increased gradually up to the maximum dose of two tablets or capsules four times a day if necessary. The drugs should give relief of symptoms in 10–30 minutes and the relief should continue for 3–10 hours. If relief is not obtained after a trial with a dosage as outlined above then another variety of antihistamine should be given, using the same precautions and scheme of dosage, until one is found which is effective and suits the patient. To lessen the amount of the drug used, I recommend that it should be given only when symptoms are beginning, and not regularly three or four times a day in the absence of symptoms.

Urticaria and Angioneurotic Oedema

Injectants, such as penicillin, and foods are frequent factors in the causation of these conditions, but in many cases psychological factors are paramount. Where specific factors are found and cannot be reasonably avoided, desensitization is recommended, but a general discussion of the psychological and other non-specific factors should be given and will usually help considerably. The antihistamines are our standby in resistant cases. When the greatest reactions occur at night those antihistamines with a common soporific side-action, such as "benadryl" and "phenegan," are best used at night, and the less soporific types, such as "antistin" or "histantin," used in the day as necessary.

Vasomotor Rhinitis: Paroxysmal Rhinorrhoea

A full investigation and discussion, as for asthma, will usually lead to the proper regime for relief. It is a common finding to obtain from a case of asthma a history of years of rhinorrhoea. Antihistamines are valuable for symptomatic treatment in some cases while investigation and treatment are proceeding.

Allergic Eczema

This is extremely common in the child up to the age of 6 or 7, when it is often replaced by asthma. In the infant and young child food sensitivity is very common, chocolate, cheese, egg-white, fish, and wheat being the most common allergens. It would seem from Bray's work that the almost universal hypochlorhydria in children below the age of 7 allows absorption of unsplit proteins, which causes the eczema. After the age of 7 the digestive function usually improves and food reactions are less common. As these children tend to become sensitive to dust, etc., after a few years, it is wise to carry out all precautions in the bedroom as if these sensitivities were already present.

For the eczema, apart from local creams, the administration of antihistamines—for example, "elixir

benadryl" or "elixir anthisan"—one to four teaspoonfuls, according to age, especially at night but also in the day, if necessary, to relieve the irritation, gives great benefit. Dilute hydrochloric acid, 30 min. (1.8 ml.) in water three times a day with meals, despite its homoeopathic dose, seems to improve appetite and digestion. Any foods to which the child is sensitive should, of course, be banned. Milk is usually best after it has been boiled and allowed to cool and the thin albuminous skin has been removed. I rarely attempt to desensitize children before the age of 6–7, as I believe that in most cases the harm caused by the psychological trauma of the battle to give a course of injections outweighs the benefit obtained.

Conclusions

The prognosis in allergic conditions depends largely on the care with which the multiple factors concerned in nearly all cases are investigated and explained to the patient, and on the thoroughness of the steps taken to deal with these factors. In most cases sufficient specific factors are discovered to make desensitization an important part of the control, but, except perhaps in hay-fever, desensitization alone, without reference to the other factors discussed, will not be successful.

ORDER OF ST. JOHN OF JERUSALEM

The *London Gazette* has announced the following promotions in, and appointments to, the Venerable Order of the Hospital of St. John of Jerusalem:

As Knight: Dr. C. E. Salt. *As Commanders (Brothers):* Surgeon Rear-Admiral K. A. I. Mackenzie, C.B., M.R.C.S., L.R.C.P., K.H.P., Brigadier Sir A. E. Porritt, K.C.M.G., C.B.E., F.R.C.S., Wing Commander (Acting Group Captain) J. S. Carslaw, M.B., Sir H. K. Graham-Hodgson, K.C.V.O., F.R.C.P., Drs. W. J. Vickers, C.M.G., M.R.C.S., L.R.C.P., W. E. Hutchinson, and M. M. Scott. *As Officers (Brothers):* Surgeon Captains J. Hamilton, C.B.E., M.B., and W. P. E. McIntyre, M.D., R.N., Major-Generals W. E. Tyndall, C.B., C.B.E., M.C., M.B., K.H.S., and F. Harris, C.B., C.B.E., M.C., M.B., K.H.S., Brigadier H. C. Edwards, C.B.E., F.R.C.S., Major R. R. M. Porter, M.C., I.M.S., retired, Squadron Leader D. S. Grant, M.R.C.S., L.R.C.P., Drs. J. B. Mackie, W. Radcliffe, E. R. A. Merewether, C.B.E., C. J. P. Paglar, J. T. P. Handy, G. Swapp, D.C.M., M.M., H. M. Golding, D.F.C., M. V. Roberts, E. M. B. West, O.B.E., and T. H. Jenkins, and Mr. L. W. Hefferman, M.D., F.R.C.S. *As Associate Officers (Brothers):* Drs. Ip Kam Wah, P. C. Kwan, and S. I. M. Ibrahim. *As Serving Brothers:* Major S. Mackenzie, O.B.E., M.B., R.A.M.C., Professor R. E. Lane, M.D., Drs. W. C. Murray, G. S. Mackay, M.C., Y. C. Hooi, G. H. Oorloff, J. M. de Ville-es-Offrans, M.B.E., A. J. Amor, C.B.E., H. F. Chard, T. A. L. Davies, G. F. Keatinge, E. Holland, W. G. Arthur, J. P. Michael, R. L. Brown, A. Chalmers, T. E. Wood, and J. S. M. Wilson. *As Associate Serving Brothers:* Drs. O. H. Leong and T. C. Dey. *As Serving Sisters:* Drs. Margaret L. Dobbie-Bateman and (Mrs.) Margaret M. H. Gibson-Hill.

The Northumberland Association of Parish Councils thinks more doctors should be appointed as magistrates and is writing to the Lord Lieutenant asking him to arrange for the inclusion of at least one doctor on each bench in the area wherever this is possible, according to a report in the *Newcastle Journal*. It considers that as so many cases coming before the magistrates now have psychological or medical aspects the advice of doctors as magistrates would be very valuable. The honorary public relations secretary of the North of England Branch of the B.M.A. said that doctors would welcome the plan; at present only two doctors sit on Newcastle city bench, and in the county of Northumberland there are about four serving.