Effective adrenergic blocking agents able to prevent the hypertensive attacks have been described (Console *et al.*, 1950) and their severe side-reactions discussed (Swan, 1951). Some, such as benzodioxane, act quickly and are useful in the acute hypertensive attack whether spontaneous or provoked by histamine. Others, such as "dibenamine," are slower to act but more prolonged and complete in action, and preliminary reports (Bartels and Cattell, 1950; Crowther, 1951) suggest that these probably offer the best hope of success in preparing the patient for later surgery by preventing the state of shock.

We are grateful to Dr. Russell Fraser and Mr. M. R. Ewing for permission to publish the case, which was under their care.

REFERENCES

- Bartels, E. C., and Cattell, R. B. (1950). *Ann. Surg.*, **131**, 903.
 Console, A. D., Dunbar, H. S., and Ray, B. S. (1950). *Surgery*, **28**, 428.
 Crowther, K. V. (1951). *British Medical Journal*, **1**, 445.
 Duncan, L. E., Semans, J. H., and Howard, J. E. (1944). *Ann. intern. Med.*, **20**, 815.
 Goldner, M. G. (1947). *J. clin. Endocrinol.*, **7**, 716.
 Green, D. M. (1946). *J. Amer. med. Ass.*, **131**, 1260.
 Labbe, E. M., Tinel, J., and Doumer, E. (1922). *Bull. Soc. méd. Hôp. Paris*, **46**, 982.
 Pantridge, J. F., and Burrows, M. McC. (1951). *British Medical Journal*, **1**, 448.
 Rabin, C. B. (1929). *Arch. Path.*, **7**, 228.
 Smithwick, R. H., Greer, W. E. R., Robertson, C. W., and Wilkins, R. W. (1950). *New Engl. J. Med.*, **242**, 252.
 Swan, H. J. C. (1951). *British Medical Journal*, **1**, 440.
 Walton, J. N. (1950). *Lancet*, **1**, 438.
 Washington, E. L., Callahan, W. P., and Edwards, E. W. (1946). *J. clin. Endocrinol.*, **6**, 688.
 Wells, A. H., and Boman, P. G. (1937). *J. Amer. med. Ass.*, **109**, 1176.
 Wilkins, R. W., Greer, W. E. R., Culbertson, J. W., Halperin, M. H., Litter, J., Burnett, C. H., and Smithwick, R. H. (1950). *Arch. intern. Med.*, **86**, 51.

HEPATIC COMA WITH RECOVERY IN A CYSTINURIC

BY

E. D. H. COWEN, M.D., M.R.C.P.

Consulting Physician, Hitchin and St. Albans Hospitals

Recovery from hepatic coma is always an event of note. Latner (1950) reported the recovery of four out of five consecutive cases on his regime. Because no one physician is likely to see more than an occasional case of hepatic coma it is important to establish whether these results can be substantiated in other hands in a syndrome in which effective therapy has been conspicuously absent. Communication to Latner or to the medical press of all results of the use of his regime, whether favourable or not, would seem desirable.

It has been suggested that cystine deficiency may be a factor in human liver disease (Himsworth, 1945; Glynn, Himsworth, and Neuberger, 1945). The following case is therefore of especial interest in that one would think a cystinuric, losing as he does fair quantities of cystine continuously into his urine, would be most susceptible to liver disease. This case does not support that idea.

Case Report

The patient was a male physician aged 42 in good health. He had never had jaundice, but had recently attended a jaundiced pregnant woman. On October 10, 1951, he felt feverish and went to bed. Two days later jaundice

developed. His fever resolved in a further two to three days, the jaundice increased, and he began to vomit. The vomiting increased in severity until on the sixth day of his illness he could not retain water by mouth. Seen next day, he appeared very depressed and quiet, although mentally alert. He was markedly jaundiced. His liver was easily palpable but not very tender. He had a moist furred tongue and took water and glucose fluids well. He took 100 mg. of pethidine that night because he "could not get mental relaxation," and in the early hours of the following day (eighth day) he took 1½ gr. (0.1 g.) of "seconal" (usual sedative). He then slept deeply and all that day remained drowsy. Fluids were taken well, and he passed 3½ pints (2 litres) of urine. In the early hours of the ninth day, for the first time, he behaved rather oddly—pushing his wife away when getting out of bed.

When seen that morning (October 18) he lay quiet and comatose, unable to recognize relatives or to answer questions. His pupils were widely dilated. He was deeply icteric, with no petechiae. His liver was not palpable, but his respirations were shallow. There was no rise in jugular venous pressure. His blood pressure was 140/90. Ascites was not present. He was admitted to the Lister Hospital as an urgent case and placed on the Latner regime. This consists of two solutions: (a) 10% dextrose in normal saline, to each pint (570 ml.) of which are added 50 mg. of thiamin hydrochloride, 150 mg. of nicotinamide, 50 mg. of riboflavin, and 0.5 g. of potassium chloride; (b) plasma to which vitamins as in (a) but no potassium chloride have been added.

Four pints (2.3 litres) of the dextrose-saline solution were given in 24 hours, but only one pint (570 ml.) of plasma was given (October 19, morning) during the 50 hours in which the intravenous drip was running. Latner recommends three pints (1.7 litres) of dextrose-saline, followed by one pint (570 ml.) of plasma in every 24 hours. There were administered in addition to the intravenous drip: 300 mg. of tocopherol (100 mg. thrice daily) intramuscularly, 60 mg. of vitamin K daily, and 1,000,000 units of penicillin daily.

On the day of admission the patient remained very drowsy and became very restless, necessitating three 5-ml. doses of paraldehyde intramuscularly in the first 24 hours. The next day (October 19) his condition appeared very much the same (coma and restlessness). However, when examined at 5.45 p.m. he just opened his eyelids, and by 8.15 p.m. had become fairly orientated and alert. The intravenous drip was discontinued, after 50 hours, on October 20. During this time he had had 9 pints (5.1 litres) of fluid by drip (glucose 450 g., sodium chloride approximately 12 g.) and 55 oz. (1.6 litres) of fluid by mouth. He was now given tocopherol, B vitamins, and "hepovite" by mouth (Latner). From this time onwards recovery proceeded without significant episodes. He left hospital on November 7 and returned to work on January 8, 1951, feeling very well.

Laboratory Data.—On the day of admission serum bilirubin was 4.8 mg. per 100 ml., thymol turbidity 14 units, prothrombin time 27% (24 seconds), alkaline phosphatase 13.7 units, serum proteins 5 g. % (albumin 3.4 g., globulin 1.6 g.), Hb 106%. Urine examination showed bilirubinuria, urobilinogen and urobilin absent, with a trace of albumin and a few granular casts. The serum bilirubin rose to 8.2 mg. per 100 ml. two days later and to 12 mg. after one week. By December 1 it had fallen to 1 mg. per 100 ml. and by December 29 to 0.8 mg. Thymol turbidity, which lay between 14 and 10 units in the first week, fell steadily to 4.5 units on December 29. Urobilinogen and urobilin (Ehrlich and Schlesinger tests respectively) were present in quantity for the first time on November 1. The haemoglobin was 110% on November 3. On December 5 total proteins were 8 g. % (albumin 6 g., globin 2 g.).

Paper Chromatography

At the height of coma (October 18) a specimen of urine was sent to Dr. C. E. Dent through the kindness of Dr. S. P. V. Sherlock. It showed a slight generalized amino-aciduria with abnormal quantities of cystine, lysine,

and ethanolamine on paper chromatography. A further specimen of urine on December 29, when the patient was fully convalescent, showed abnormal quantities of cystine, lysine, and methyl histidine, but was otherwise normal.

Dr. Dent interprets these findings as follows. The first specimen (October 18) was compatible with severe liver damage, but not so severe as would be regarded as inevitably fatal. It also suggested that the patient was a cystinuric. The second specimen (December 29) confirmed the fact that he was a true cystinuric—that is, was one of those people who all their lives excrete in the urine exceptionally large quantities of cystine, lysine, and sometimes also arginine (Dent and Rose, 1951). The increased quantity of methyl histidine cannot yet be interpreted.

Summary

A record is given of a case of infective hepatitis developing coma which recovered after 36 hours. The Latner regime of therapy was used.

The patient showed a slight generalized amino-aciduria to paper chromatography at the height of coma. It was a surprise to find that he was a cystinuric, a fact which was confirmed later when he was convalescent. The significance of this is briefly discussed.

I am greatly indebted to Dr. C. E. Dent, of University College Hospital, for his collaboration and for permission to use his data and observations.

REFERENCES

- Dent, C. E., and Rose, A. (1951). *Quart. J. Med.* In press.
Glynn, L. E., Himsworth, H. P., and Neuberger, A. (1945). *Brit. J. exp. Path.*, **26**, 326.
Himsworth, H. P. (1945). *Proc. roy. Soc. Med.*, **38**, 101.
Latner, A. L. (1950). *British Medical Journal*, **2**, 748.

Medical Memorandum

A Successful Case of Intra-arterial Blood Transfusion

We understand that intra-arterial blood transfusion has been used in certain Continental countries, but have seen little reference to it in South African or British literature. It has several theoretical advantages over intravenous transfusion, particularly when blood is needed urgently. The blood supply to the vital cerebral centres and to the coronary arteries is directly dependent on the pressure in the aorta, and it seems that transfusion into an artery raises this far more quickly than intravenous transfusion. In the dog (Samson Wright, 1947) an increase in mean aortic pressure from 50 to 130 mm. Hg may increase the flow in the coronary arteries from 20 to 250 ml. a minute. In our case, soon after the transfusion started the aorta appeared bulging and distended, and it was obvious that the pressure within had been substantially raised. It would also seem that the risk of pulmonary oedema due to overloading the right side of the heart should be much diminished.

An essential requirement of the method is that the blood should be perfectly free of clots, otherwise there must be a considerable risk of emboli lodging in the arteries of the extremities or elsewhere. There is no longer the filtering action of the lungs as in intravenous transfusion.

When we applied the method to our case all concerned were convinced that she was moribund, and we believe that intravenous transfusion, even at the same speed, would not have saved her life. A tribute is due to the nursing staff for producing the necessary apparatus with such rapidity.

Case Report

The patient, a moderately obese woman aged 68 with carcinoma of the cervix infiltrating the bladder and rectum, was brought to the theatre to undergo the operation described by Brunschwig in which a complete pelvic clearance is carried out. The stages of the operation are as follows: panhysterectomy with excision of bladder, urethra, labia minora, rectum, followed by transplantation of ureters into the sigmoid colon and a colostomy. Her pulse was 76; blood pressure 150/90. A blood transfusion into the right leg was set up. Premedication with "omnophon," 1/3 gr. (22 mg.), atropine, 1/100 gr. (0.65 mg.), was given one hour before operation. Anaesthesia was induced with thio-pentone and maintained on nitrous oxide and oxygen with ether and the addition of fractional D-tubocurarine chloride. An oral endotracheal tube was passed.

For the first two hours of operating time the patient's condition was quite satisfactory. There was a slight rise in pulse rate and a slight fall in blood pressure. However, during the process of stripping the pelvic wall, difficulty was encountered in controlling bleeding. At this stage the haemorrhage became very profuse indeed. Within a few minutes the patient became very pale, the skin cold and moist, and the pulse at the wrists and temple unobtainable. Auscultation failed to reveal any heart sounds. Automatic respiration also ceased.

Artificial respiration with 100% oxygen was immediately instituted. Although the blood transfusion into the patient's leg was running at a fair speed she did not show any signs of recovery. Repeated injections of methylamphetamine hydrochloride ("methedrine") produced no response.

The surgeon was able to feel feeble rapid pulsation in the abdominal aorta. One of us at this moment set up another blood transfusion apparatus with all possible speed. The surgeon was provided with sterile rubber tubing and a large needle (aspirating needle, gauge 14) and requested to insert the latter into the common iliac artery with the end pointing towards the heart. The blood was now pumped in at high speed and within 10 minutes a litre had been given.

During this short period a remarkable recovery occurred. The patient started automatic respiration and a fairly good pulse was palpable at the wrist and temple.

The operation was completed. Oxygen was administered continuously until consciousness was regained. The patient left the theatre with a pulse of 90 and a systolic pressure of 65. During transport to the ward some two hours after the intra-arterial transfusion her condition deteriorated slightly. There was some bleeding per vaginam. She was put into bed in the Trendelenburg position and given oxygen and omnophon, 1/6 gr. (45 mg.) intravenously.

A fortnight after the operation the patient was making good progress and there was no sign of mental impairment, nor were there any respiratory complications. There was, however, some bowel atonia for about a week.

D. C. DEVITT, M.R.C.S., L.R.C.P., D.A.
C. WIGDEROW, M.B., B.Ch., B.Sc., D.A.

REFERENCE

- Wright, Samson (1947). *Applied Physiology*, p. 447. Oxford Univ. Press, London.

A note in *Nature* (July 14, page 86) discusses the possibility that the colour sensitivity of the human eye is influenced by the physical position of the observer. Usually, when the observer is in the erect position, both eyes show the same colour sensitivity, but if he lies on one side a difference between the colour responses of the two eyes develops; the lower eye becomes markedly red-sensitive compared with the other. When he turns on to his back the two eyes resume their original similar response, and by turning on to the other side the formerly red-sensitive eye, now uppermost, gradually develops blue-sensitivity.