The time taken for allergy to develop may be appreciably diminished by increasing the dose used.

Although complications are not generally harmful they are nevertheless undesirable, and the effort to secure a minimal conversion time should be restricted to selected infants for whose special circumstances the time factor is of great importance.

It is suggested that in such circumstances complications may be prevented or minimized by giving the injections in divided doses at different sites.

We are indebted to Professor F. R. G. Heaf for his help in arranging for our supplies of vaccine; to Dr. van Diense and Colonel Bensted for the freeze-dried vaccine; to Professor Arvid Wallgren and Dr. J. H. Magnusson for generous gifts of B.C.G. tuberculin; and to our obstetric colleagues at St. Mary's Hospital for their willing co-operation.

REFERENCES

- Debré, R., Brissaud, H. E., Mozziconacci, P., Cousin, Maud, and Kaplan, A. (1950). *Quart. Rev. Pediat.*, **5**, 143.
 Heaf, F. R. G. (1950). *British Medical Journal*, **2**, 1353.
 Magnusson, J. H., and Lithander, A. (1949). *Pediatrics*, **3**, 429.
 Rekling, E. (1945). *Nord. Med.*, **27**, 1819.
 Smith, H. V., and Vollum, R. L. (1950). *Lancet*, **2**, 275.
 Ustvedt, H. J. (1951). *Conference of European B.C.G. Programmes*, p. 253. Heinemann, London.
 Wallgren, A. (1950). *Tuberculosis and Other Problems of Pediatrics*, p. 33. Williams and Wilkins, Baltimore.

PETHIDINE AS AN ADJUVANT TO NITROUS OXIDE AND OXYGEN ANAESTHESIA

BY

P. DUNBAR JOHNSON, M.R.C.S., L.R.C.P., D.A.
Anaesthetist, Hastings Group of Hospitals

The scope of thiopentone in general anaesthesia, first popularized by Lundy in 1934, was enormously extended by the introduction of D-tubocurarine in 1942 by Griffith and Johnson. Administered by continuous intravenous drip or repeated fractional doses in combination with nitrous oxide and oxygen and, where indicated, a muscle relaxant, it has proved generally satisfactory in experienced hands.

This type of anaesthesia, however, is not without its disadvantages. Thiopentone is a powerful respiratory depressant with a weak analgesic action. Used with a gas-and-oxygen mixture, rich enough in oxygen to prevent anoxia, the amount required to obtund pain may cause undesirable respiratory depression, which, in long procedures requiring a relatively high total dose, may be prolonged into the post-operative period. Moreover, its powerful hypnotic but weak analgesic action is a cause of post-operative restlessness. The parasympathetic effect of thiopentone is, in theory at any rate, undesirable, especially when combined with a muscle relaxant, which, itself a potential liberator of histamine, may initiate bronchial constriction and spasm.

The search for a suitable adjuvant to nitrous oxide and oxygen anaesthesia led to the introduction into clinical anaesthesia of pethidine hydrochloride by Neff *et al.* (1947) in America, and by Mushin and Rendell-Baker (1949) in England.

First synthesized by Eisleb and Schaumann in 1939, pethidine is widely used in surgery and obstetrics for the relief of pain. Many other piperidine derivatives

related to it have since been synthesized, and the properties of two of these—bemidone “10446” and keto-bemidone “10720”—have been described by Thompson and Neff (1950). They found the keto-compound “10720” to be fifteen to twenty times more powerful than pethidine, with which it compared favourably as an adjuvant to nitrous oxide and oxygen anaesthesia in dogs and human beings.

Unlike morphine, pethidine produces a high degree of analgesia without undue effect on the respirations and the blood pressure. It exerts a weak narcotic effect, thus allowing, in contradistinction to thiopentone, a rapid return to consciousness with delayed post-operative pain. It possesses, in addition, other properties, which have perhaps not been stressed enough by anaesthetists.

Professor J. H. Burn (1948) has drawn attention to the common action of a wide group of substances used medicinally. This group includes procaine, quinidine, atropine, the antihistamines, and pethidine. All have in greater or lesser degree a local anaesthetic action, the power to relax smooth muscle and to lengthen the refractory period of the heart, an analgesic and an antihistamine effect. Thus pethidine, in addition to its powerful analgesic action, has 80% of the activity of quinidine on the auricle, is similar in potency to procaine as a local anaesthetic, and has an antihistamine effect which is considerable. Burn (1950) suggested that intravenous pethidine would be more effective than procaine for the release of bronchial spasm in man. That it has a selective action on the laryngeal reflex has been shown by Ruben and Andreassen (1951), who, using intravenous pethidine alone, have intubated a series of 25 cases without eliciting laryngeal spasm.

The action of pethidine on the heart has been noted by Mushin and Rendell-Baker (1950), who, in over 100 thoracic operations, were particularly impressed by the absence of pulse irregularities when using gas and oxygen, gallamine triethiodide, and pethidine, and who suggest that this combination of drugs is especially suitable for operations on the heart.

Morphine is known to cause spasm of the lower end of the common bile duct near the sphincter of Oddi, yet it still continues to be used by anaesthetists and surgeons before and after operations on the biliary passages. Pethidine, because of its spasmolytic effect, is the drug of choice. Similarly, pethidine exerts a spasmolytic effect on the distal colon and rectum, and Griffiths (1950) has noted its value in combination with gas and oxygen in such operations as haemorrhoidectomy and posterior colporrhaphy.

The theoretical advantages of pethidine as a substitute for thiopentone in the maintenance of nitrous oxide and oxygen anaesthesia led me to try it in a large number of cases. In an attempt to arrive at a satisfactory standard technique use was made of wide variations in premedication, induction, and the dose and concentration of pethidine employed for maintenance. Premedication was with morphine and atropine, “omnupon” and scopolamine, pethidine and scopolamine, or pethidine and atropine in appropriate doses. In most cases thiopentone was used for induction, but some were induced with nitrous oxide and oxygen and others with cyclopropane. Intubation was performed under thiopentone and a muscle relaxant, thiopentone and cyclopropane, or a combination of all three. Maintenance was with nitrous oxide and oxygen, using a circle absorber or the semi-closed

method. The concentration of oxygen was kept at 30 to 50%. Pethidine was given in fractional doses varying between 20 and 50 mg. and in concentrations of 5 to 50 mg. per ml. The muscle relaxants used were gallamine triethiodide, D-tubocurarine, dimethyltubocurarine iodide, and decamethonium iodide.

Present Series

As a result of this preliminary trial a standard technique was evolved which was adopted in a series of more than 200 unselected cases. This series covers a variety of planned and emergency operations. Many were poor-risk cases of a high average age, the oldest patient being 98 and the youngest 12.

The results were uniformly good compared with other methods of anaesthesia. Notable features were the absence of respiratory depression, the relaxation of the bronchial tree, which greatly facilitated controlled or aided respiration, the rapid return to consciousness, and the prolonged post-operative analgesia. Fluctuations in the pulse rate and blood pressure were minimal. The high degree of analgesia was shown in patients allowed to come round too rapidly. When almost fully conscious they tolerated the endotracheal tube and seemed insensitive to skin suture.

This technique was found especially suitable when using hexamethonium bromide for controlled hypotension.

(i) Premedication

All patients in this series were given an intramuscular injection of pethidine and scopolamine one hour before operation, the dose for a healthy adult being 100 mg. of pethidine and 0.43 mg. of scopolamine. The sedation obtained appeared adequate and did not give rise to the respiratory depression which often occurs on the subsequent injection of pethidine in patients too heavily premedicated with morphine. This preliminary injection acted as a test dose and gave some indication of the subsequent pethidine requirements. It appeared to facilitate intubation and counteract the parasympathetic effect of thiopentone.

(ii) Induction

A 5% solution of thiopentone was used as a routine, but a 2.5% solution was found safer in old and shocked patients. The dose varied from 0.25 g. to 0.75 g. Intubation was performed under thiopentone and gallamine triethiodide.

(iii) Maintenance

This was with nitrous oxide through a circle absorber, using a concentration of 40–50% oxygen. Immediately normal respiratory rhythm was established pethidine was injected slowly, using 50 mg. in a concentration of 10 mg. per ml. for the average patient and 25 mg. in a concentration of 5 mg. per ml. in the old or debilitated. Subsequent doses, which varied between 20 and 50 mg., were given when indicated. Timing and dosage are a matter of experience, but 20 mg. every 20 minutes proved to be the average. Relaxation was maintained by gallamine triethiodide in 20-mg. doses when indicated.

A Gordh needle was found useful for the repeated injections, but it must be remembered that when pethidine and thiopentone come in contact a milky precipitate is

thrown down, due presumably to the marked difference in pH. The indwelling needle must therefore be cleared of all traces of thiopentone before pethidine is injected.

The total quantity of pethidine needed varied between 50 and 200 mg., but one patient, an extremely powerful young man, required 225 mg. during the course of an operation for reconstitution of the shoulder-joint. He was fully conscious within 20 minutes of the termination of the operation.

A more satisfactory procedure, especially in relatively long operations, is the administration of pethidine by continuous intravenous drip. By this method a more uniform level of anaesthesia is maintained and the total quantity of pethidine required has been found to be less than when administered in repeated fractional doses. A suitable dilution is 100 mg. of pethidine in 500 ml. of saline delivered from a graduated flask. The rate of flow is varied to suit the circumstances, but an average of 1 mg. of pethidine a minute has been found adequate to maintain satisfactory anaesthesia. The drip is progressively slowed down as the operation proceeds, being discontinued altogether before the end of the operation if a rapid return to consciousness is desired. Robust patients, however, may require an additional initial dose of 25 mg. (10 mg. per ml.). If a flask graduated in 100, 50, and 25 ml. is used the total dose of pethidine administered may be calculated at any time.

(iv) Post-operative Treatment

Pethidine has a weak narcotic effect. In the less painful procedures sleep can be procured on the night following the operation by a sedative such as one of the short-acting barbiturates. The more painful conditions require an analgesic post-operatively, but injection can usually be delayed until six or eight hours after recovery of consciousness. It is well to remind the nursing staff that pethidine is an analgesic and not a sedative.

Discussion

In the experimental series of cases pethidine was first given in a concentration of 50 mg. per ml. In a certain number of patients a single dose of 25 mg. caused pronounced respiratory depression, amounting in a few instances to complete apnoea for varying periods. In some cases a transient fall in blood pressure was noted. Since adopting a dilution of 10–5 mg. per ml. for intermittent injections and 0.2 mg. per ml. for continuous administration, no significant effect on the respirations has been observed.

D. Zuck (1951) reported prolonged apnoea and a profound fall in blood pressure in a patient given 25 mg. of pethidine following induction with thiopentone and gallamine triethiodide. It is worth noting that thiopentone, especially if administered rapidly, will cause an appreciable fall in blood pressure, and that pethidine potentiates the effect of the thiopentone. It is doubtful whether 25 mg. of pethidine, even if injected rapidly, in a concentration of 50 mg. per ml. would have such a profound effect on the circulatory and respiratory systems when given alone. It has been shown by Thompson and Neff (1950) that pethidine, given rapidly in doses of 5–10 mg. per kg. to anaesthetized dogs in a concentration of 50 mg. per ml., invariably produced a prompt and significant fall in blood pressure. When the injection was made slowly and the concentration reduced to 5–10 mg. per ml. no significant variation

of the blood pressure occurred. They found that slow injections of dilute solutions minimized the undesirable effects on respiration and circulation of piperidine derivatives resembling pethidine when administered to patients under nitrous oxide and oxygen anaesthesia.

Summary

The employment of pethidine hydrochloride as an adjuvant to nitrous oxide and oxygen anaesthesia has been discussed, with special reference to its anticholinergic and antihistaminic properties.

A method of maintenance by a continuous intravenous pethidine drip is outlined.

The importance of slow injections of dilute solutions is emphasized.

REFERENCES

- Burn, J. H. (1948). *The Background of Therapeutics*. Oxford Medical Publ., London.
 — (1950). *British Medical Journal*, 2, 691.
 Griffiths, H. F. (1950). *Ibid.*, 2, 763.
 Mushin, W. W., and Rendell-Baker, L. (1949). *Ibid.*, 2, 472.
 — (1950). *Brit. J. Anaesth.*, 22, 235.
 Neff, W. B., et al. (1947). *Calif. Med.*, 66, 67.
 Ruben, H., and Andreassen A. K. (1951). *Brit. J. Anaesth.*, 23, 33.
 Thompson, R. C., and Neff, W. B. (1950). *Curr. Res. Anesth.*, 29, 77.
 Zuck, D. (1951). *British Medical Journal*, 1, 125.

POST-MORTEM EXAMINATIONS ON AIR-CRASH VICTIMS

BY

DONALD TEARE, M.D., M.R.C.P.

*Assistant Pathologist and Lecturer in Forensic Medicine,
St. George's Hospital Medical School; Lecturer in Forensic
Medicine, St. Bartholomew's Hospital Medical College*

On October 31, 1950, a Viking aircraft of British European Airways, flying from Paris to London, crashed in thick fog at London Airport. The wreckage burst into flames and 28 out of the 30 occupants of the plane were killed. Post-mortem examinations on all the victims were ordered by the coroner of the district, for the purposes of identification and to determine the cause of death. Since the religious tenets of many of the victims required a rapid burial the examination of the remains became something of an urgent problem.

Organization

The facilities of the local mortuary were not ideal for examination of such a large number of bodies. It had a refrigeration chamber for six, one fixed post-mortem table, and room for three trolleys. A marquee was set up close to the mortuary, in which the bodies were first placed and numbered. They were there stripped of such clothing as remained, and this was put into bags and similarly numbered. To assist in the identification, there were present a coroner's officer and a senior member of an undertaking firm, who had had experience in similar types of disaster. They had a complete passenger list of the occupants of the plane when it left Paris. There were two mortuary assistants and several bearers. The post-mortem reports were typed out directly in the mortuary by my secretary. The general principle employed was to examine those readily identifiable first, and so narrow down the number which might prove difficult of identification towards the end.

As it was thought necessary to estimate the carbon monoxide haemoglobin of each victim, 4 ml. screw-cap bottles were numbered in advance and 28 post-mortem reports were similarly numbered.

The specimens of blood were taken into these screw-capped bottles, filled to the brim to obviate any loss of carbon monoxide from the blood—a loss which may vitiate the value of a carboxyhaemoglobin estimation, particularly if the bottle is shaken in transport. Urea estimations were performed on the specimens of blood at varying intervals, up to seven days, in the hope of finding some variation which might assist in the estimation of the time of death in cases in which such time was unknown. They proved valueless.

Identification

Though most of the bodies were very severely burnt, particularly about the face and head, the problem of identification did not prove as difficult as was at first thought. Facial recognition was, except in a few cases, quite impossible, but in most cases the clothing of the chest and abdomen was relatively spared and passports were recovered from the inside coat pockets of the majority of the men. The women, keeping their passports in handbags, proved the more difficult to identify. The question of sex never presented any difficulty, but stature was difficult to estimate owing to the destruction and contortion of the limbs as the result of heat. The age could be roughly estimated by the state of the costal cartilages, skull sutures, and other skeletal changes.

As soon as a body was definitely identified, either by papers or by clothing, the name was added to the number on the body, on the clothing, and on the post-mortem report and specimen of blood. These steps led to the rapid identification of 25 of the victims. There remained two women and one man, and these three were all identified positively by means of their teeth. It must be remembered that, while a person's name may appear on the passenger list of a crashed aircraft, the relatives are not unnaturally loath to accept this as the only evidence of the death of one close to them. The man was an American, and his dental chart was transmitted from New York for identification.

The Cause of Death

The post-mortem examinations revealed several important points. All the victims had received injuries which would lead to immediate loss of consciousness or death. Twenty-three of the victims had no measurable concentration of carbon monoxide in their blood; three had less than 15%; and two between 15% and 20%. None of the latter showed any burning of the internal air passages such as is seen in the victims of a bursting boiler or the "flash and flame" discharge of a high-voltage electric current. It was reasonable to suppose, therefore, that most of the victims were dead before the outbreak of fire, and that the remainder, whose injuries were very severe, were unconscious and lived only for a matter of seconds in the fire. Furthermore, it would appear that had no fire occurred the fate of the occupants would have been the same. The actual causes of death may be summarized as follows:

Fractured skull	..	..	..	..	..	10
Fractured cervical spine	..	..	..	..	..	1
Traumatic subarachnoid haemorrhage	..	..	..	..	..	1
Ruptured aorta	..	..	..	..	..	8
Ruptured lungs	..	..	..	..	..	3
Ruptured heart	..	..	..	..	..	3
Ruptured liver and spleen	..	..	..	..	..	2