TROMEXAN AND DICOUMAROL

BY

D. G. ILLINGWORTH, M.B., M.R.C.P.Ed.*Formerly Medical Registrar, Northern Hospitals Group,
Edinburgh*

There is now wide acceptance of the value of anti-coagulant therapy. The necessity for prolonged courses of heparin was obviated by the introduction of dicoumarol, which brought the advantages of oral administration and relative cheapness. The disadvantages of dicoumarol, such as its slowness of action, its cumulative toxicity with the attendant risk of haemorrhage, and the necessity for daily prothrombin estimations often caused a hesitancy in employing it. Rosicky (1944) produced a new coumarin drug—bis-3:3'-(4-oxycoumarinyl) ethyl acetate—which is now available as "tromexan" (Geigy). Evidence is accumulating that this drug has advantages over dicoumarol.

Pulver and von Kaulla (1948) observed that in man tromexan disappeared from the blood in 36 hours; with dicoumarol the time lag was five days. They also recorded a method for assessing the tromexan level in plasma. Reinis and Kubik (1948) claimed for tromexan a shorter latent period than that of dicoumarol, easier control, and a more rapid return to normal of the prothrombin level after cessation of treatment. They observed a certain individual variation with regard to tromexan activity and only slight toxicity. Von Kaulla and Pulver (1948) adduced, by animal experiments, evidence supporting these contentions.

Deutsch (1949) observed the production of initial hyperprothrombinaemia in man, as with dicoumarol, while Bickel and Della Santa (1949) put forward a plea for divided dosage and recorded only mild and transient haemorrhagic incidents.

The low toxicity of tromexan was illustrated by the proposal of Della Santa and von Kaulla (1949) to use the effect of the drug on the prothrombin level as a test of liver function, while Lian *et al.* (1950) employed the drug for as long as 11 weeks without tolerance or cumulative toxicity occurring.

With dicoumarol the maximum effect on the prothrombin level may be delayed as long as 72 hours. Burt, Wright, and Kubik (1949) pointed out that the tromexan lag period is usually 8 to 24 hours. According to Burke and Wright (1950) the maximum effect was obtained in 28 to 36 hours, although an appreciable therapeutic effect was reached in 18 to 24 hours, while the prothrombin level returned to normal in 60 to 72 hours. Reinis and Kubik (1948) found that this "hang-over" effect might, however, be prolonged up to seven days. It is estimated by Burke and Wright that tromexan has one-sixth the potency and one-twentieth the toxicity of dicoumarol. Experimental evidence has been adduced by Kubik and Reinis (1950) to demonstrate the effectiveness of tromexan in the reduction of intra-arterial clot. All publications claim for the drug relative safety and good clinical results.

Administration and Control of Tromexan Therapy

The drug is administered orally in tablets of 300 mg., which are grooved to facilitate smaller dosage. In the present series no patient complained of the taste of the drug, and there was no gastric upset.

Prothrombin estimation was achieved by Quick's method, using human brain as a source of thromboplastin. It was found to be an advantage, though usually impracticable, to have a reading before starting treatment. In the absence of complicating factors these initial readings varied between 60 and 100%. With congestive cardiac failure preliminary readings as low as 45% were recorded. Adjustment of the initial loading dose was therefore necessary.

A therapeutic level of between 30 and 10% of normal prothrombin was aimed at and a course of three weeks' duration envisaged, as with dicoumarol. A loading dose of 1,200 mg. proved safe and became standard, although 2,100 mg. was given twice without incident. All cases of pulmonary embolism and those of cardiac infarction with marked systemic upset were given heparin initially. At the outset of the investigation an intravenous drip of normal saline with 300 mg. of heparin was provided concomitantly for the first 24 hours, as this was thought to be the average tromexan lag period. In three out of five cases, however, a prothrombin concentration of less than 10% resulted within 24 hours. Latterly there was evolved the plan of giving one intravenous dose of 100 mg. of heparin along with the loading dose of tromexan. Multiple divided doses of heparin in the first 24 hours of treatment were found not only unnecessary and economically wasteful but they apparently invoked a complicating factor in prothrombin depression, for which allowance had to be made in estimating tromexan dosage.

Between 12 and 24 hours after starting treatment, and before further tromexan was administered, a prothrombin concentration test was performed: 70% of the cases were found to have less than 50% prothrombin, and 34% had entered the therapeutic range of 10 to 30% prothrombin. A few cases showed resistance to the drug and two demonstrated an initial increase in prothrombin.

At this point it should be made clear that before starting the course of tromexan it is impossible to forecast the dosage necessary for a given patient to enter or remain in the therapeutic zone. Age and body weight give no assistance, and even the response to the loading dose is occasionally deceptive in that a high degree of drug resistance may be soon overcome with intensive dosage, and minimal doses suffice thereafter.

Subsequent to the loading dose the daily ration was divided into two or three doses, but no tromexan was given until each morning's prothrombin value was to hand. A technique of trial and error was employed, although usually the necessary maintenance dose lay between 300 and 600 mg. Two patients were well controlled on a daily average of 150 mg.; two others required between 900 and 1,200 mg. each day. No case was maintained on a constant daily dosage. A rough guide to dosage is given in Table I.

TABLE I

Prothrombin Concentration	Daily Ration of Tromexan
100-80%	1,200 mg.
80-60%	900-750 mg.
60-40%	750-600 ..
40-20%	600-300 ..
20-10%	300-150 ..
Under 10%	0

Unfortunately, most patients exhibited a sudden escape from control after being well maintained for several days. This seldom took the form of a dangerous drop in prothrombin, but consisted of a sudden rise requiring increased dosage to recover equilibrium.

There is therefore no escape from frequent, preferably daily, prothrombin estimations. It is significant that in association with these episodes of apparent drug resistance were recurrences of ischaemic pain in the cardiac cases, and in one case a fresh cardiac infarction. The basic defect in such incidents is possibly in intestinal absorption of the drug, and it may be that inactivation takes place under certain intestinal conditions.

Depression of the prothrombin concentration below 10% was not infrequent, usually in the first days of treatment. It gave rise to little of the concern that one felt when dicoumarol was being used as the anticoagulant. Of the 72 cases treated with tromexan only one who reached the "danger-zone" bled, when haematemesis resulted from an active duodenal ulcer. Dicoumarol compares unfavourably in this respect, and it seems unwise to assess the patient's reaction to the coumarin drugs purely on prothrombin levels (Bickel and Della Santa, 1949).

Frequent search was made for microscopical haematuria. Bleeding incidents occurred in two of the 72 patients treated. Both suffered a haematemesis from a clinically active duodenal ulcer. One of these had a prothrombin concentration of less than 10%, which was regarded formerly as dangerously low. The other was in the centre of the therapeutic range, however. Neither case required any more active treatment than the cessation of tromexan therapy. It is considered that in the presence of peptic ulceration tromexan should be used with caution. A serious haemorrhage may be anticipated, and possibly averted, by the use of frequent benzidine tests of the faeces of such patients.

After the last dose of tromexan the prothrombin level was followed in 20 patients. The average time for recovery was 60 hours, the extremes being 14 hours and 14 days. Prolongation of action could not be correlated with delayed initial effect or drug resistance (Burt *et al.*, 1949).

Results of Tromexan Therapy

(a) *Cardiac Infarction*.—Sixty cases of cardiac infarction were treated (Table II). Confirmation of the clinical diagnosis was obtained by electrocardiogram or necropsy in all. Fifty-one cases left hospital, the immediate end-

TABLE II.—Results

	Anti-coagulant	No of Cases	Discharges	Hospital Deaths	Bleeding Manifestations
Cardiac infarction	Tromexan	60	51	9	2
	Dicoumarol	36	29	7	7
Extracardiac thrombosis	Tromexan	12	12	0	0
	Dicoumarol	22	19	3	6

result being satisfactory. Nine patients died in hospital, but two of these succumbed after the cessation of treatment—one from pneumonia and the other from renal failure secondary to gross atheroma and hypertension. Of the remaining seven who died, each of whom had suffered an anterior infarction, one died twelve hours after admission with haemopericardium and ventricular rupture. Two others, who died within a few days, had rupture of the left ventricle, while another two came to necropsy with previous multiple infarcts and a grossly scarred myocardium. Necropsy was refused in one aged woman who succumbed after attacks of acute left ventricular failure. A cerebro-vascular accident was coincident. The remaining fatal case never entered the thera-

peutic range of prothrombin. Sudden death occurred on the seventeenth day of treatment, and necropsy revealed a large infarct which had probably slowly extended.

(b) *Extracardiac Group*.—Twelve cases of extracardiac thrombo-embolic incidents received tromexan. Six suffered from phlebothrombosis, while six had also sustained a pulmonary embolus. The end-result was regarded in all as satisfactory.

Bleeding Incidents.—Among these 72 patients, as previously recorded, there were two cases of haematemesis in duodenal ulcer sufferers. The only necessary measure was the withdrawal of tromexan.

Results of Dicoumarol Therapy

(a) *Cardiac Infarction*.—Thirty-six cases of cardiac infarction were treated (Table II). The clinical diagnosis was confirmed in 35. In one case the electrocardiogram was normal but the diagnosis indisputable. Twenty-nine cases left hospital, the immediate end-result being satisfactory. Two patients died after the cessation of dicoumarol—one from a recurrence of infarction, the other from pneumonia. The remaining five who succumbed during treatment had sustained anterior infarcts. One died from a recurrence while in the therapeutic range of prothrombin. A further death occurred 36 hours after admission with extensive and recurrent infarcts and gross cardiac failure. One who died suddenly had a grossly scarred left ventricle, which was freshly infarcted while the patient was in the therapeutic range. Gross scarring was likewise noticed in a further patient who suffered profuse haematuria till the time of death in spite of transfusion. The last had suffered what was clinically ventricular rupture. No necropsy was permitted.

(b) *Extracardiac Group*.—Twenty-two cases of extracardiac thrombo-embolic incidents received dicoumarol. Eleven suffered from phlebothrombosis alone; seven had also sustained a pulmonary embolus. Four had developed peripheral arterial obstruction. Nineteen left hospital with an end-result regarded as satisfactory. Of the three deaths, two occurred after the cessation of therapy—one from renal carcinoma, the other from cerebral thrombosis. The third patient died during the dicoumarol treatment of a pulmonary embolus. The main cause of death was prolonged epistaxes and haematuria which failed to respond to transfusion.

Bleeding Incidents.—Of the 58 patients who received dicoumarol 13 bled. Five of these were not out of control as judged by prothrombin levels. Three lost blood from several orifices simultaneously. Haematuria occurred in nine, epistaxes in four, melaena in two, while there was one case of haematemesis and one of subcutaneous bleeding. All received intravenous vitamin K, and three required transfusion. Two of those who bled succumbed; the haemorrhage was an important factor in their death.

Conclusions

Tromexan in clinical practice is superior to dicoumarol. As it has a shorter "lag period" the use of heparin may be either obviated or restricted to a single intravenous dose, with great economic benefit.

The clinical results compare favourably with those of dicoumarol therapy; indeed, the fatal cases, with the exception of one patient who was inadequately treated, were poor risks. Haemorrhagic complications are not only less frequent but infinitely less severe. A striking

comparison is possible in the effects of a fall in prothrombin concentration below 10%, which was regarded in the tromexan group with relative equanimity.

None of the cardiac patients receiving tromexan suffered a recurrence of ischaemic pain while in the therapeutic range of 10 to 30% prothrombin. Two patients in the 40 to 50% range sustained further thrombotic episodes, and the return of ischaemic cardiac pain, not amounting to infarction, was always coincident with inadequate dosage. Two of the cardiac cases receiving dicoumarol developed fresh infarctions while ostensibly admirably controlled.

There were marked variations in susceptibility to tromexan. Indeed, a given patient's tolerance may alter widely from day to day, and there is a tendency to sudden escapes from control. It is therefore not desirable to give the drug without laboratory control. No correlation of dosage with weight or age is possible, but an initial prothrombin reading may assist in gauging the size of the loading dose.

Caution is necessary in treating peptic ulcer patients. Benzidine tests on the faeces may be helpful in the control of their treatment.

Summary

Clinical results are recorded in 130 patients suffering from vascular thrombosis and its sequelae.

Ninety-six of these had sustained a cardiac infarct. Sixty received tromexan and 36 dicoumarol.

A scheme for the use of tromexan is proposed and the literature briefly reviewed.

I wish to express my thanks to Dr. H. N. Robson, who suggested this investigation; to the physicians and surgeons of the Northern Hospitals Group, Edinburgh, who permitted access to their patients; and to the staff of the haematological laboratory for technical assistance. The tromexan was supplied by Pharmaceutical Laboratories, Geigy Ltd., Manchester.

REFERENCES

- Bickel, G., and Della Santa, R. (1949). *Rev. méd. Suisse rom.*, **49**, 545.
 Burke, G. E., and Wright, I. S. (1950). *Bull. N.Y. Acad. Med.*, **26**, 264.
 Burt, C. C., Wright, H. P., and Kubik, M. (1949). *British Medical Journal*, **2**, 1250.
 Della Santa, R., and von Kaulla, K. N. (1949). *Helv. med. Acta*, series A, **16**, 251.
 Deutsch, E. (1949). *Schweiz. med. Wschr.*, **79**, 1010.
 Von Kaulla, K. N., and Pulver, R. (1948). *Ibid.*, **78**, 806.
 Kubik, M., and Reinis, von Z. (1950). *International Congress of Haematology*, Cambridge.
 Lian, C., Siguier, F., Welti, J. J., Piette, M., Coblentz, B., and Trélat, J. (1950). *Bull. Soc. méd. Hôp. Paris*, **66**, 110.
 Pulver, R., and von Kaulla, K. N. (1948). *Schweiz. med. Wschr.*, **78**, 956.
 Reinis, von Z., and Kubik, M. (1948). *Ibid.*, **78**, 785.
 Rosicky, J. (1944). *Cas. Lék. ces.*, **83**, 1200.

In New York during the 32 years 1918-50 there were 104 fatal injuries from competitive sports, and 73 post-mortem examinations were made. Writing in the *Journal of the American Medical Association*, August 18, page 1506, Dr. T. A. Gonzales discusses the kinds of lethal injury sustained in different sports and their frequency, without, unfortunately, stating the numbers at risk in each activity. There surely cannot be many people in New York who play cricket, for instance, yet one death is recorded in this game—that of a player struck in the abdomen by a ball which ruptured the small intestine and led to acute peritonitis. This must be as rare a cause of death in England as that of the relay runner who ran into a railing alongside the track and ruptured his large bowel. Baseball has the highest total, with 43 deaths, and football (with 22 fatalities) was about level with boxing (21); but surely there are fewer boxers than football players in New York City?

DOSAGE OF PENICILLIN IN WEIL'S DISEASE

BY

C. W. LAWSON, M.B., M.R.C.P.

Senior Medical Registrar, Royal Infirmary, Leicester

Penicillin has been used for some years in the treatment of Weil's disease (leptospirosis icterohaemorrhagica), and its efficacy, especially in the pre-icteric stage of the disease, is generally accepted. There is, however, wide variation in the dosage recommended. Chang (1946) recommended a total daily dosage of 250,000-300,000 units of sodium penicillin. Patterson (1947) used three-hourly injections of 15,000 units. Suchett-Kaye (1951) used 100,000 units four-hourly and raised the dose to 500,000 units four-hourly if the patient's condition warranted it. The following cases illustrate some of the important clinical aspects of the disease and lend support to the use of a dosage of the order recommended by Suchett-Kaye.

Case 1

A butcher aged 32, who had bathed several times in a canal during the three weeks preceding his illness, was admitted on the sixth day of the disease, having had severe pains in the limbs for five days and frequent vomiting for three days.

He was a big, well-nourished man, obviously ill, with a temperature of 102.6° F. (39.2° C.). He had an erythematous rash on the face and trunk. There was marked suffusion of the conjunctivae, which were bright red. Jaundice was not present. The tongue was coated and dry. Slight neck rigidity was present, but Kernig's sign was negative. The muscles of the thighs and forearms were tender. The liver was not palpable, but he was tender in the right hypochondrium. The spleen was not palpable. His blood pressure was 110/80. The rest of the physical examination was negative. A provisional diagnosis of Weil's disease was made and various confirmatory tests were carried out. The urine contained a moderate amount of albumin, but was otherwise normal. The haemoglobin was 84%; white cells, 2,900 (polymorphs 74%, lymphocytes 22%, monocytes 4%); serum bilirubin, less than 0.5 mg. per 100 ml.; blood urea, 165 mg. per 100 ml. N.P.N., 97 mg. per 100 ml. A radiograph of the chest was normal. Blood taken at the time of admission for agglutination for *Leptospira icterohaemorrhagiae* was later reported as negative.

Soon after admission, before penicillin was started, his temperature rose to 104.8° F. (40.5° C.) and he became delirious and hallucinated, seeing people moving on the wall. Penicillin was therefore begun in doses of 30,000 units four-hourly by intramuscular injection. Thirty-six hours later his condition had not improved much, though his temperature was a little lower and he was no longer hallucinated. He was very drowsy. The dosage of penicillin was therefore increased to 100,000 units six-hourly. After this his temperature fell to normal over the next 48 hours and his clinical condition rapidly improved. This dosage was continued for eight days until a total of 3,500,000 units had been given. Convalescence was uneventful apart from a mild attack of gout in the left great toe on the eleventh day of the disease. He had never had gout before, but his mother suffered from it.

Blood taken on the eleventh day of the disease gave positive agglutinations for *L. icterohaemorrhagiae* at a dilution of 1/3,000.

Case 2

A hosiery packer aged 17, who had often bathed in a river during the month preceding his illness and whose workplace was infested with rats, was admitted on the third day