The reported claims of recent isolations of classical influenza virus A strains are considered. The evidence suggests that these claims should be viewed with great caution.

We should like to thank Dr. A. W. Gledhill for his help in preparing immune ferret sera and Mr. E. Owen for his most valuable technical assistance.

REFERENCES

- Andrewes, C. H. (1950). *New Engl. J. Med.*, **242**, 161.
 — (1951). *Proc. roy. Soc. Med.*, **44**, 803.
 — Glover, R. E., Himmelweit, F., and Smith, W. (1944). *Brit. J. exp. Path.*, **25**, 130.
 Bradley, W. H. (1951). *Proc. roy. Soc. Med.*, **44**, 789.
 Burnet, F. M., and Bull, D. R. (1943). *Aust. J. exp. Biol. med. Sci.*, **21**, 55.
 Chu, C. M., Andrewes, C. H., and Gledhill, A. W. (1950). *Bull. World Hlth Org.*, **3**, 187.
 Freyche, M. J., and Klimt, C. (1951). *World Hlth Org., Epidem. and Vital Statist. Rep.*, **4**, 141.
 Fulton, F., and Dumbell, K. R. (1949). *J. gen. Microbiol.*, **3**, 97.
 Hilleman, M. R., Mason, R. P., and Buescher, E. L. (1950). *Proc. Soc. exp. Biol., N.Y.*, **75**, 829.
 Isaacs, A., and Bozzo, A. (1951). *Brit. J. exp. Path.* In press.
 Magrassi, F. (1949). *Minerva med., Torino*, **40**, 565.
 Meenan, P. N., and Clarke, M. (1951). *J. Irish med. Ass.*, **29**, 3.
 Nagler, F. P., van Rooyen, C. E., and Sturdy, J. H. (1949). *Canad. J. publ. Hlth*, **40**, 457.
 Svedmyr, A., von Magnus, P., and Freundt, E. A. (1951). *Acta path. microbiol. scand.*, **29**, 96.
 van Rooyen, C. E., McClelland, L., and Campbell, E. K. (1949). *Canad. J. publ. Hlth*, **40**, 447.
 Veen, J. van der, and Mulder, J. (1950). "Studies on the Antigenic Composition of Human Influenza Virus A Strains." Stenfort Kroese, Leyden.
 Ward, T. G., and Eddy, B. E. (1950). *Science*, **112**, 501.

RELATION OF ADRENAL CORTICAL FUNCTION TO SCURVY IN GUINEA-PIGS

PRELIMINARY EXPERIMENTS

BY

B. E. CLAYTON, Ph.D., M.B., Ch.B.*

AND

F. T. G. PRUNTY, M.A., M.D., F.R.C.P.

(From the Department of Chemical Pathology, St. Thomas's Hospital Medical School)

It has been known for many years that the adrenal gland contains a very high concentration of ascorbic acid (Szent-Györgyi, 1928), and more recently the increasing amount of information on adrenal steroids has centred interest on the relationship between ascorbic acid and adrenal steroid metabolism.

Following the work of Sayers *et al.* (1944, 1946) it has become well known that pituitary adrenocorticotrophic hormone, which stimulates the release of adrenal steroids, also causes a fall in adrenal ascorbic acid. This is true for the rat, which can synthesize ascorbic acid, and for the guinea-pig, which is unable to do so (Sayers *et al.*, 1946; Long, 1947a, 1947b) and therefore is dependent upon an adequate dietary intake. In the absence of ascorbic acid the guinea-pig develops scurvy with all its well-known manifestations (Bessey *et al.*, 1934).

The present investigation was undertaken to determine the effect of scurvy on urinary 17-ketosteroid excretion.

These excretory products are generally thought to arise from the adrenal cortex and, to a less extent, from the testes in the male (Fraser *et al.*, 1941). Observations have also been made on the effect of A.C.T.H. and cortisone in scorbutic guinea-pigs, with particular reference to the effect on development of symptoms and the excretion of 17-ketosteroids.

17-Ketosteroid Excretion During Untreated Scurvy

Methods

Male and female guinea-pigs of varying weights (300 to 590 g.) obtained from two registered breeders were kept in groups of two or three in metabolism cages. They were maintained on the scorbutic diet described by Harris and Ray (1932), which was supplemented by cabbage *ad lib.* where healthy pigs were required. Tap-water and the scorbutic diet were supplied *ad lib.* but the amount eaten each day was usually measured. Body weights were determined daily.

Daily, 24-hour specimens of urine were collected over concentrated A.R. hydrochloric acid, except on Sundays. Urinary 17-ketosteroids were determined at frequent intervals. After acid hydrolysis and carbon tetrachloride extraction, the Zimmerman reaction as described by Callow *et al.* (1938) was carried out, and a colour correction was finally applied according to Talbot *et al.* (1942). Of each group of animals one was killed when moribund and the organs were sectioned. All animals were examined at death, and the diagnosis of scurvy was confirmed. In some healthy and scorbutic animals the ascorbic acid content of the adrenal glands was determined by the method of Roe and Kuether (1943).

Results

In normal male and female guinea-pigs the daily 17-ketosteroid excretion varied between 0.09 and 0.30 mg./24 hours.

In Table I a series of results is given for four separate groups of animals. This shows the daily average body weight in grammes, and the average 17-ketosteroid excretion in milligrams per 24 hours. In addition, for group I, details of daily food and daily urine volume are given. In Table II the day on which a fall in body weight began and the day of death are given.

1. All animals showed a fall in body weight beginning on the 9th to the 15th day after starting the scorbutic diet.

2. The daily food intake began to fall rapidly about the 12th to the 16th day after the scorbutic diet began.

3. Death from scurvy occurred in all animals on the 21st to the 29th day of the experiment.

4. In all groups of animals the urinary 17-ketosteroid excretion showed a gradual but definite increase of varying degree as the animals became scorbutic, and excretion reached a peak in the terminal phases.

5. When not required for histological examination the right adrenal gland was weighed to the nearest 0.1 mg. Scorbutic glands showed an increase in weight over glands from healthy pigs, and the mean ratio of right adrenal gland (mg.) to body weight (g.) was 0.40 ± 0.03 . Corresponding figures for healthy pigs are 0.30 ± 0.03 .

6. The ascorbic acid content of the adrenal glands of healthy pigs varied between 106 and 164 mg./100 g. adrenal tissue. Ascorbic acid was not detected chemically in the glands of grossly scorbutic animals, neither could it be demonstrated histologically.

*Holden Research Fellow.

TABLE I.—17-Ketosteroid Excretion During Untreated Scurvy

Diet	Day of Experiment	Group I: Experiment I (2 Female Pigs)				Group II: Experiment II (3 Male Pigs)		Group III: Experiment III (2 Female Pigs)		Gp. IV: Experiment IV (2 Male Pigs)	
		Average Body Weight in g.	Average Food Intake in g./24 hr.	Average Urine Vol. in ml./24 hr.	Average 17-Ketosteroid Excretion in mg./24 hr.	Average Body Weight in g.	Average 17-Ketosteroid Excretion in mg./24 hr.	Average Body Weight in g.	Average 17-Ketosteroid Excretion in mg./24 hr.	Average Body Weight in g.	Average 17-Ketosteroid Excretion in mg./24 hr.
Scorbutic diet + cabbage <i>ad lib.</i>	1 2	521.5				485.0 450.0	0.200 0.100	302.5	0.119	350.0	0.098
Scorbutic diet	1	521.5				451.0	0.300	281.5	0.076	336.5	
	2		24			442.0		337.0		326.0	
	3	455.0	24	12.5	0.176	437.0		347.5		364.0	
	4	439.5	23	10		435.5	0.300	357.0		382.0	0.198
	5	465.0	43	7		434.0		321.5			
	6	496.5	45	12	0.119	426.5					
	7	487.5	42	16				340.0		377.5	
	8	494.0	43	11	0.204	426.0	0.360	353.5	0.187	380.5	0.148
	9		27	12		421.5		358.0		396.5	
	10	484.5	27	12		419.0		349.5		402.0	
	11	502.5	43	13		418.0	0.300	341.5		390.5	0.238
	12	496.0	42	9	0.119	414.0		337.5			
	13	502.0	34	9		408.0				371.5	
	14	504.0	32	11			0.470			352.5	
	15	493.0	33	11		379.0	0.640	342.5		331.0	
	16		26	7		377.5	0.650	339.5	0.336	318.0	0.418
	17	464.5	26	7		375.0		320.5		316.5	0.413
	18	466.0	10	6	0.192	359.0	0.680	322.5	0.320	297.0	0.505
	19	452.5	11	4		340.5	0.880	307.0			
	20	434.5	7	17	0.350	321.5					
	21	422.5	6	6		285.5	0.650	254.0	0.265	265.0	0.485
						(2 dead)				(dead)	
	22	398.0	4	8	0.390	247.0	0.960	259.5	0.630		
						(3rd dead)					
	23		3	2				242.0			
	24	346.0	3	2	0.740			221.0	0.685		
	25	344.5	3	3				208.5			
								(dead)			
	26	327.5	2	1	0.700						
	27	309.0	3	1	0.945						
	28	288.5	1	3	0.785						
	29	275.5 (dead)	1	1	0.640						

TABLE II.—Influence of Cortisone and A.C.T.H. on Fall in Body Weight, Survival Time, and Adrenal Weight in Scurvy

Treatment	Body Weight in g. Beginning of Experiment	Day of Experiment on which Fall in Body Weight Begins	Day of Death or Day of Killing When Moribund	Right Adrenal Weight in mg.	Ratio of Right Adrenal Weight (mg.) to Body Weight (g.)*
None: healthy	587			214.8	0.37
	542			156.0	0.23
	608			230.8	0.38
	655			188.2	0.29
	324			90.7	0.28
	328			76.2	0.23
Scorbutic diet: no treatment	507	12	21	202.8	0.34
	425	11	21		
	415	12	22		
	540	15	29	208.4	0.39
	500	13	29		
	435	12	21	121.4	0.25
	320	11	21	127.7	0.39
	305	9	25		
	300	9	25	182.9	0.52
	590	15	27	282.4	0.47
	545	15	27	326.4	0.50
	395	15	28	146.5	0.32
	320	15	28	184.8	0.47
Scorbutic diet: 5 mg. cortisone daily	525	17	28		
	470	17	28		
	345	17	24	125.1	0.36
	445	17	24		
	415	14	22	118.5	0.29
Scorbutic diet: 20 mg. cortisone daily	380	14	21		
	620	12	22	172.2	0.28
Scorbutic diet: A.C.T.H. 20 mg. daily	485	12	21	139.7	0.29
Scorbutic diet: A.C.T.H. 10 mg. daily	385	10	22	259.0	0.67
	340	11	18	186.6	0.54
Scorbutic diet: A.C.T.H. 10 mg. daily	710	15	27	311.5	0.44
	585	17	27	285.5	0.47

* Body weight is the maximum recorded during the experiment prior to A.C.T.H. or cortisone treatment.

Effect of A.C.T.H. and Cortisone

Method

A.C.T.H. or cortisone administration was begun at the same time as the scorbutic diet. A.C.T.H. in physiological saline was administered daily to two groups of pigs (male). The first group received the equivalent of 10 mg. La-1-A (Armour's unit) as four subcutaneous injections daily at two-hour intervals. The second group received 20 mg. daily as eight subcutaneous injections at hourly intervals. Cortisone acetate (Merck's suspension) was given to male and female animals as one daily subcutaneous injection of 5 or 20 mg. Daily body weight, urine volume, and food intake were measured, and 17-ketosteroids were determined at frequent intervals.

Results

1. All animals showed a fall in body weight and subsequently died with manifestations of scurvy (Table II). Applying a "t" test for the day on which the fall in body weight began, the 95% confidence limits for the mean difference are -5.012 or +1.578—that is, it is unlikely that treatment increased the time of the first loss of weight by more than five days or decreased it by more than one and a half days. The 95% confidence limits for the mean difference in survival time are -3.019 and +5.019, so that, similarly, it was unlikely that treatment increased the survival time by more than three days, or decreased it by more than five days. No definite anti-scorbutic effect has therefore been demonstrated.

2. Food intake began to fall at the same time as in untreated scorbutic controls.

3. Preliminary experiments suggested that daily cortisone in a dosage of 5 mg. suppressed the endogenous urinary 17-ketosteroid excretion. The results are summarized in Table III, and, though not suitable for statis-

TABLE III.—Effect of 5 mg. Daily Cortisone on 17-Ketosteroid Excretion

Healthy Females		Scorbutic Females		Scorbutic Females		Scorbutic Males	
Day of Experiment	17-Ketosteroid Excretion	Day of Experiment	17-Ketosteroid Excretion	Day of Experiment	17-Ketosteroid Excretion	Day of Experiment	17-Ketosteroid Excretion
2	0.210	8	0.181	2	0.090	4	0.189
6	0.320	11	0.102	5	0.125	7	0.216
13	0.365	14	0.340	8	0.150	10	0.175
21	0.335	17	0.370	11	0.130	13	0.121
22	0.385	19	0.240	14	0.113	17	0.200
27	0.453	22	0.440	17	0.232	20	0.248
		24	0.390	19	0.219	22	0.092
		26	0.456	21	0.119	24	0.261
		28	0.655				

Results are given in mg. per animal per day, in groups of two.

tical analysis, they are very suggestive. A daily dosage of 20 mg. raised the excretion to about 1 mg. in control and scorbutic pigs. Sprague *et al.* (1950) suggested that 17-ketosteroids which appear in the urine during cortisone administration are largely derived from administered cortisone.

The findings with A.C.T.H. were inconclusive. During the first 15 days or so of the scorbutic diet a response to A.C.T.H. undoubtedly occurred, but without a large number of experiments it will not be possible to tell whether the grossly ascorbic-acid-depleted animals are responding to A.C.T.H., as the high urinary 17-ketosteroid excretion may merely be the normal response to acute advancing scurvy. Experiments are in progress to try to determine whether the grossly depleted adrenal gland can respond to stimulation by A.C.T.H. It is interesting to note, however, that the highest 17-ketosteroid excretion yet found—1.7 mg. in 24 hours—occurred in scorbutic animals receiving daily A.C.T.H. For comparison, it is interesting to note that healthy animals excreted 0.7 mg./24 hours after four days on 10 mg. A.C.T.H. daily in divided doses; with a daily dosage of 20 mg. A.C.T.H., excretion varied between 0.8 and 1.2 mg./24 hours.

4. The administration of cortisone prevented the increase in weight of the adrenal glands normally found in scurvy, and the mean ratio of right adrenal weight (mg.) to body weight (g.) was 0.30 ± 0.02 . A.C.T.H.-treated animals had enlarged glands with a mean ratio of 0.5 ± 0.05 .

5. Histological examination of liver, spleen, kidney, wrist-joint, and ankle-joint revealed no significant differences between A.C.T.H.- or cortisone-treated and untreated scorbutic animals. No ascorbic acid could be demonstrated histologically in the adrenal glands of treated scorbutic pigs.

Discussion

In a long series of papers, Giroud *et al.* (1939, 1940a, 1940b, 1940c) have concluded that ascorbic acid favours the production of adrenal cortical hormones and improves their utilization. They consider that a sufficiency of vitamin C is necessary for the secretion of cortical hormones, that a fall in vitamin C is accompanied by a fall in the suprarenal functional capacity, and that vitamin C and the cortical steroids are used simultaneously. The evidence that the concentrations of

steroids and vitamin C in adrenal glands paralleled each other was based, for the steroids, on biological assay. In this assay the partial contraction of chromatophores in certain fish was considered a specific response to cortical steroids. This is, of course, open to question, and no urinary steroid determinations were carried out. They lay much stress on their considered opinion that the symptoms of scurvy in guinea-pigs are similar to those of adrenal insufficiency, and that both conditions show disturbances of water, salt, carbohydrate, and protein metabolism. After the administration of cortical extract they claimed survival of scorbutic pigs for 40 instead of 28 days, though, apart from less apathy and muscle atrophy, morphological differences were not evident.

Later it was concluded by Schaffenburg *et al.* (1950) that cortisone inhibited many of the manifestations of scurvy in the guinea-pig. Their experiments were continued for only 21 days, but during that time body weight was maintained, joints remained mostly normal, and haemorrhage was less severe. Hyman *et al.* (1950) reported similar findings with 25 mg. A.C.T.H. daily and 12.5 mg. cortisone daily, but added a footnote to the effect that the protective effect might not be apparent owing to a "toxic" effect of these substances. They also claimed an increased survival time in A.C.T.H.-treated pigs. On the other hand, during a study of wound-healing in normal and scorbutic pigs, Upton and Coon (1951) investigated the effects of A.C.T.H. and cortisone in scurvy, but they could obtain no conclusive evidence of any effect on growth rate, nutrition, or degree of haemorrhage. Our results are in accord with the view of these authors, since they lend no support to the idea that A.C.T.H., or cortisone in dosage of 5 or 20 mg. daily, can alleviate the development of scorbutic symptoms in these animals, and therefore they do not form a substitute for vitamin C.

Dugal and Thérien (1949) showed that the typical enlargement of the adrenals which occurs in response to the stress of cold in rats and in guinea-pigs can be completely prevented by large doses of ascorbic acid. They thought that ascorbic acid apparently played a compensatory part rather similar to that of one of the adrenal hormones. Sayers and Sayers (1947) had already shown that if adrenal cortical extract was administered to rats immediately before a stress, then the enhanced adrenal activity such as normally occurs is suppressed. On the other hand, the development of scurvy itself may be considered a "stress situation," since adrenal enlargement occurs, as we have shown in confirmation of the earlier work of LaMer and Campbell (1920). That this is prevented by cortisone is established in our experiments and was also noted by Schaffenburg *et al.* (1950).

It has been suggested by Sayers *et al.* (1946) that changes in ascorbic acid and cholesterol produced by A.C.T.H. are associated with the formation and release of adrenal steroids. Long (1947b) has suggested that ascorbic acid may be directly concerned with the formation of adrenal steroids, or may merely be one of a series of metabolic changes associated with increased secretion by the gland. He described an experiment in which A.C.T.H. was administered to scorbutic guinea-pigs. Though it produced no further fall in adrenal ascorbic acid, there was a fall in cholesterol and the development of lymphopenia, so that a release of cortical hormones had apparently taken place. Jailer and Boas

(1950) found no change in the ascorbic acid content of chick adrenals even though corticosteroids were released.

While our experiments were in progress Nadel and Schneider (1951) reported that the "formaldehyde-genic substances" in guinea-pig urine showed a five-fold increase in the later stages of scurvy.

A.C.T.H. can produce a fall in the level of circulating eosinophils in scorbutic animals, as shown by Eisenstein and Shank (1951). From this, and the fact that scurvy is accompanied by adrenal hypertrophy, they concluded that ascorbic acid was not essential for adrenal function and not directly concerned in the formation and release of 11-oxysteroids.

Our experiments seem to confirm an increase of adrenal cortical activity in scurvy as judged by adrenal hypertrophy, and an increase of cortical hormone from the increase of 17-ketosteroid excretion. Both of these changes are reversible by cortisone, although this has no effect on the disease. A.C.T.H. is capable of causing additional hypertrophy. The 17-ketosteroid excretion is certainly increased during the earlier stages of the development of scurvy, but in the fully scorbutic animal, even in the absence of exogenous A.C.T.H., it appears to be nearly maximal. A much larger number of observations must be made before it is definitely established whether the output of these substances is significantly increased under these conditions by A.C.T.H. The balance of evidence favours the view that adrenal cortical activity is increased in scurvy, and can be further increased, in the absence of vitamin C, by A.C.T.H.

Summary

Guinea-pigs placed on a scorbutic diet showed a gradual well-marked increase of varying extent in urinary 17-ketosteroid excretion. This reached a peak in the terminal phases.

Daily administration of A.C.T.H. and cortisone failed to influence the fall in body weight and subsequent death of the animals on a scorbutic diet. Such animals showed similar manifestations of scurvy as untreated scorbutic controls.

We are grateful for the statistical assistance given us by Mr. P. Armitage, of the M.R.C. Statistical Unit; for histological assistance from Dr. H. E. M. Kay; and for technical assistance from Miss J. Hammant, Mr. A. E. Clark, and Mr. R. J. Hurst.

REFERENCES

- Bessey, O. A., Menten, M. L., and King, C. G. (1934). *Proc. Soc. exp. Biol., N.Y.*, **31**, 455.
- Callow, N. H., Callow, R. K., and Emmens, C. W. (1938). *Biochem. J.*, **32**, 1312.
- Dugal, L. P., and Thérien, M. (1949). *Endocrinology*, **44**, 420.
- Eisenstein, A. B., and Shank, R. E. (1951). *J. clin. Endocrinol.*, **11**, 791.
- Fraser, R. W., Forbes, A. P., Albright, F., Sulkowitch, H., and Reifenshtein, E. C. (1941). *Ibid.*, **1**, 234.
- Giroud, A. (1940a). *Pr. méd.*, **48**, 841.
- and Ratsimamanga, A. R. (1940b). *Ibid.*, **48**, 449.
- and Santa, N. (1939). *C.R. soc. Biol., Paris*, **131**, 1176.
- and Martinet, M. (1940c). *Ibid.*, **134**, 23.
- Harris, L. J., and Ray, S. N. (1932). *Biochem. J.*, **26**, 2067.
- Hyman, G. A., Ragan, C., and Turner, J. C. (1950). *Proc. Soc. exp. Biol., N.Y.*, **75**, 470.
- Jailer, J. W., and Boas, N. F. (1950). *Endocrinology*, **46**, 314.
- LaMer, V. K., and Campbell, H. L. (1920). *Proc. Soc. exp. Biol., N.Y.*, **18**, 32.
- Long, C. N. H. (1947a). *Recent Prog. hormone Res.*, **1**, 99.
- (1947b). *Fed. Proc.*, **6**, 461.
- Nadel, E. M., and Schneider, J. J. (1951). *J. clin. Endocrinol.*, **11**, 791.
- Roe, J. H., and Kuether, C. A. (1943). *J. biol. Chem.*, **147**, 399.
- Sayers, G., Sayers, M. A., Lewis, H. L., and Long, C. N. H. (1944). *Proc. Soc. exp. Biol., N.Y.*, **55**, 238.
- Liang, T. Y., and Long, C. N. H. (1946). *Endocrinology*, **38**, 1.
- Sayers, M. A., and Sayers, G. (1947). *Ibid.*, **40**, 265.
- Schaffenburg, C., Masson, G. M. C., and Corcoran, A. C. (1950). *Proc. soc. exp. Biol., N.Y.*, **74**, 358.
- Sprague, R. G., Power, H. M., Mason, H. L., Albert, A., Mathieson, D. R., Hench, P. S., Kendall, E. C., Slocumb, C. H., and Polley, H. F. (1950). *Arch. intern. Med.*, **85**, 199.
- Szent-Györgyi, A. (1928). *Biochem. J.*, **22**, 1387.
- Talbot, N. B., Berman, R. A., and MacLachlan, E. A. (1942). *J. biol. Chem.*, **143**, 211.
- Upton, A. C., and Coon, W. W. (1951). *Proc. Soc. exp. Biol., N.Y.*, **77**, 153.

PREPARATION OF CLEAN CHEMICAL SOLUTIONS

WITH SPECIAL REFERENCE TO RADIOACTIVE PHOSPHORUS SOLUTIONS FOR CLINICAL USE

BY

G. E. HARRISON, Ph.D., F.Inst.P.

AND

W. H. A. RAYMOND

(From the Medical Research Council Radiobiological Research Unit, Harwell)

In the application of radioactive isotopes to biological studies the preparation of clean solutions is a matter of considerable importance. Especially is this the case in the medical use of radioactive isotopes, for, in addition to the possible unsuspected localization of the particulate material after injection (see the following article by Lamerton and Harriss), there is the further complication of the uncertainty of the dosage delivered to a particular organ. Indeed, this very effect may account for some of the wide variations in the dose required to produce a given biological change reported in the literature—variations which are commonly attributed to differences in the biological sensitivity of a particular organ in different individuals. There is also the obvious physical complication that the realization of standard solutions of a radioactive isotope is greatly complicated if part of the activity is in a non-exchangeable, particulate form.

The present experiments were carried out on radioactive phosphate solutions as prepared at Harwell for clinical use. It should be noted, however, that the methods employed are of general application and are likely to be effective in all cases in which a high degree of cleanliness of a preparation is of importance.

In the dispensing of radioactive phosphate solutions for clinical use the inactive carrier is isotonic saline (0.9 g. NaCl %) buffered to pH 7 with sodium phosphate containing 1 mg. P/ml. A small aliquot of carrier-free radioactive phosphoric acid is then added to a measured volume of the buffered saline in a "pyrex" ampoule, to give an activity of about 1 mc./ml. As the phosphorus content of the mixture is practically the same as that of the carrier, the ratio of P^{32}/P^{31} in this solution is about 3×10^{-6} . The ampoule is sealed and then autoclaved at 20 lb./sq. in. (1.4 kg./cm.²) pressure (temperature 126°C.) for 20 minutes. Specimens of phosphate solution prepared by the methods in use to the end of 1950 were faintly hazy, and on centrifugation at 2,500 r.p.m. for 15 minutes up to 50% of the activity could be removed in the centrifugate. When