electroencephalogram the next day showed mild to moderate diffuse cerebral dysfunction, but no epileptiform activity. On follow-up examination weeks later, no spontaneous or stimulus-sensitive myoclonus was observed.

Discussion

Myoclonus is a known manifestation of neuroexcitotoxicity in patients receiving long-term opioids. In palliative care settings, wherein patients are given long-term high-dose opioids, the incidence of opioid-induced myoclonus ranges from 2.7% to 11%.¹ With the exception of a single case report,² opioid-induced myoclonus has been described only in the setting of long-term and high-dose administration of opioids.¹ It has been hypothesised that neuroexcitotoxicity is secondary to a gradual accumulation of several morphine and hydromorphone metabolites in plasma and the CSF and is not due to the opioid itself.^{1,3} Animal and in vitro studies of opioid-induced myoclonus have almost unanimously implicated the build-up of neuroexcitotoxic opioid metabolites, such as morphine-3-glucuronide or hydromorphone-3-glucuronide.^{1,3} For this reason, periodic rotation of different types of opioids, which reduces CSF levels of individual metabolites, is usually the treatment of choice.¹

In contrast with previously reported cases, our patient developed severe neuroexcitotoxicity within a brief interval after the acute administration of a single dose of hydromorphone. This strongly suggests that in at least a subset of cases, neuroexcitotoxicity is not dependent only on accumulation of opioid metabolites but can be induced by the opioids themselves.

Controversy exists as to whether neuroexcitotoxicity of opioids is reversible with naloxone. Early studies suggested that the neuroexcitotoxicity of a morphine metabolite was inversely proportional to its affinity to the opioid receptor. Whereas certain in vitro and animal studies showed diminution or abolishment of neuroexcitotoxicity with the administration of naloxone,⁴ others found opioid antagonists to be ineffective at reducing experimentally induced neuroexcitotoxicity.⁵ Perhaps in cases where accumulation of opioid metabolites is responsible for neuroexcitotoxicity, naloxone may be incapable of reversing the side effect. On the other hand, our patient had a complete abolition of myoclonus with a single dose of naloxone, suggesting that in acute-onset cases naloxone may be of use.

It is interesting to note that our patient had multiple system atrophy. Myoclonus is often seen in multiple system atrophy and is thought to be cortical in origin. This is, however, generally mild and usually consists of infrequent axial, stimulus-sensitive myoclonic jerks. Our patient's underlying condition, however, may have led to a predisposition to such a strikingly unusual presentation of opioid-induced myoclonus.

This case report shows that, in contrast to previous studies, opioid-induced myoclonus can occur acutely and with administration of low doses. The acuity of its development suggests that myoclonus is not reliant on accumulation of metabolites but can be caused by the drug itself. Naloxone may be of use in acute-onset cases.

A Hofmann, N Tangri

Department of Internal Medicine, McGill University, Montreal, Quebec, Canada

A-L Lafontaine, R B Postuma

Department of Neurology, McGill University

Correspondence to: Ronald B Postuma, Department of Neurology, L7-305 Montreal General Hospital, 1650 Cedar Avenue, Montreal, Quebec, Canada H3G 1A4; ron.postuma@mcgill.ca

doi: 10.1136/jnnp.2005.076588

Competing interests: None declared.

References

- 1 Mercadante S. Pathophysiology and treatment of opioid-related myoclonus in cancer patients. *Pain* 1998;**74**:5-9.
- 2 Stuerenburg HJ, Claassen J, Eggers C, et al. Acute adverse reaction to fentanyl in a 55 year old man. *J Neural Neurosurg Psychiatry* 2000;**69**:281-2.
- 3 Thwaites D, McCann S, Broderick P. Hydromorphone neuroexcitation. *J Palliat Med* 2004;**7**:545-50.
- 4 Tortella FC, Robles L, Mosberg HL. Evidence for mu opioid receptor mediation of enkephalin-induced electroencephalographic seizures. *J Pharmacol Exp Ther* 1987;**240**:571-7.
- 5 Frenk H, Watkins LR, Mayer DJ. Differential behavioral effects induced by intrathecal microinjection of opiates: comparison of convulsive and cataleptic effects produced by morphine, methadone, and D-Ala2-methionine-enkephalinamide. *Brain Res* 1984;**299**:31-42.

Transient global amnesia after gastroscopy

Transient global amnesia (TGA) is an amnesic syndrome characterised by sudden-onset, clear-cut, anterograde amnesia during the episode, but otherwise normal neurological functions.¹ Episodes end within 24 h.¹ TGA may follow various precipitating events, such as emotional stress, contact with cold water, physical exercise or sexual intercourse.^{2,3} Medical action precipitating onset of TGA is reported to be coronary angiography or painful procedures, but reports of TGA after endoscopic procedures are few. We present three patients who developed TGA after gastroscopy.

Case reports

One morning, a 58-year-old woman, a nurse's aid, underwent gastroscopy at the hospital where she worked. After the operation, she immediately returned to work on a ward. A co-worker noticed she had amnesia because she was repeatedly asking, "Where do I bring this patient?" She was taken to the neurology department for examination. Her blood pressure was 140/70 mm Hg, and heart rate 68 beats/min. She demonstrated anterograde amnesia, the inability to learn new facts and repetitious questioning. She was unable to recall any of three objects after 5 min. No cranial nerve dysfunction was presented. Tendon reflexes in the four extremities were normal, as were muscle strength and sensory examination results. Cranial CT scan was negative, and an electroencephalogram (EEG) detected no epileptic abnormalities. Brain MRI diffusion-weighted images obtained during the episode were normal. Seven and a half hours after the onset of symptoms she was reoriented, but did not recall anything that happened during the episode.

A 59-year-old woman was taken by a nurse to the emergency room because of amnesia after gastroscopy. Her blood pressure was 150/90 mm Hg and heart rate 90 beats/min.

She was alert, but had had retrograde amnesia for several years as well as anterograde amnesia. She repetitively inquired, "Why do I stay here?" Her pupils were 3 mm in diameter and equal. She had no neurological deficits, except recent memory disturbances. Brain CT scan was normal. She was admitted for observation. Her EEG detected no epileptic discharges. She regained her memory within 15 h. Brain MRI diffusion-weighted images 5 days after onset showed no abnormal lesions.

A 65-year-old woman who had undergone gastroscopy at a hospital afterwards sat on a chair in the waiting room completely conscious. Her daughter asked her for the results of her procedure, but she could not remember what the doctor had told her. She was alert, but had anterograde amnesia and repeatedly asked the same questions. She regained her memory 8 h after onset, and was brought to our hospital 10 h later. On arrival, no neurological deficits were found, except the recent ameliorated memory disturbances already described. Brain MRI diffusion-weighted images and an EEG 12 days after onset were normal.

Clinical profiles of these three patients are given in table 1. A veteran endoscopist with more than 10 years of experience handled the endoscope. The patients were placed at the left lateral position during gastroscopy. Patients 2 and 3 had been premedicated with an intramuscular injection of scopolamine butylbromide (20 mg). All three patients received 8% topical pharyngeal lidocaine spray before undergoing endoscopy. None had had endoscopic polypectomy, nor a history of migraine, epilepsy or stroke.

Discussion

These three patients showed sudden amnesia after gastroscopy. The common clinical feature, typical of TGA, was the repeated asking of questions with retrograde and anterograde memory disturbance. The patients had no other neurological abnormality, and fulfilled the diagnostic criteria for TGA.¹ All of them had a gastroscopy history, and the endoscope had been handled by a veteran endoscopist. Their cases suggest that gastroscopy may have adverse effects resulting in TGA. Although scopolamine easily penetrates the blood-brain barrier and scopolamine intoxication can cause anterograde amnesia,⁴ these patients had no anti-cholinergic symptoms such as mydriasis, tachycardia, urinary retention or dry mouth. In addition, patient 1 had not received any anti-cholinergic agent before gastroscopy. Even without such agents, therefore, TGA may occur after gastroscopy. Autonomic function of these patients during gastroscopy could not be clearly evaluated. Autonomic dysfunction or emotional stress are, however, possible reasons for their TGA. Changes in autonomic nervous activity during endoscopy are reported to occur reflexively when an endoscope passes through the upper digestive tract, because of excessive parasympathetic tone, anxiety, premedication, discomfort or rapid gastric dilatation,⁵ and the memory disorder of TGA is speculated to be triggered by these factors. As shown here, gastroscopy may precipitate TGA. These atypical features of TGA indicate that further study is required to confirm the pathophysiological mechanism. TGA as a result of gastroscopy is rare, but doctors need to be aware of the possibility of amnesic events after this procedure.

Table 1 Clinical features of patients with transient global amnesia after gastroscopy

Age (years), sex	Vascular risk	Episode duration (h)	Previous gastroscopy (times)	Premedication	Pharyngeal lidocaine spray	Experience of endoscopist (years)
58, Female	Hypertension	7.5	1	None	Yes	35
59, Female	None	Approximately 15	2	Scopolamine	Yes	14
65, Female	Hyperlipidaemia	8	12	Scopolamine	Yes	35

A Hiraga, T Matsunaga

Department of Neurology, Asahi General Hospital, Chiba, Japan

Correspondence to: Dr Akiyuki Hiraga, Department of Neurology, Asahi General Hospital, 1-1326 Asahi, Chiba 289-2511, Japan; hiraga@mail3.alpha-net.ne.jp

doi: 10.1136/jnnp.2006.087775

Competing interests: None declared.

References

- Hodges JR, Warlow CP. The aetiology of transient global amnesia: a case-control study of 114 cases with prospective follow-up. *Brain* 1990;113:639-57.
- Fisher CM. Transient global amnesia: precipitating activities and other observations. *Arch Neurol* 1982;39:605-8.
- Rösler A, Mrass GJ, Frese A, et al. Precipitating factors of transient global amnesia. *J Neurol* 1999;246:53-4.
- Ardila A, Moreno C. Scopolamine intoxication as a model of transient global amnesia. *Brain Cogn* 1991;15:236-45.
- Hayashi T, Nomura M, Honda H, et al. Evaluation of autonomic nervous function during upper gastrointestinal endoscopy using heart rate variability. *J Gastroenterol* 2000;35:815-23.

Hu-antibody-positive patients with or without cancer have similar clinical profiles

Hu-antibodies (Hu-abs) are markers of paraneoplastic neurological syndrome (PNS), almost always associated with small-cell lung carcinoma. A few patients with PNS who are Hu-ab positive, however, never develop cancer after long-term follow-up.¹ If there were clinical differences, they could help identify those Hu-ab-positive patients who would not need serial screening for cancer. In this study, we analysed the clinical profile of Hu-ab-positive patients without cancer after a follow-up of more than 3 years and compared the findings with those in Hu-ab-positive patients with a tumour.

Methods

Of a total of 608 patients from our databases, we retrospectively identified 68 patients with PNS who were positive for Hu-abs, and had a follow-up of at least 3 years from the onset of the neurological syndrome. Patients were divided into two groups, according to the presence or absence of cancer during the follow-up. All patients without cancer underwent at least a CT of the chest. All of them had serial clinical and radiological (CT or chest x ray) examinations during the follow-up.

We analysed the following variables: age, sex, smoking habit, type of PNS, time to diagnosis of Hu-ab syndrome and Rankin score at diagnosis and at last visit. Data were analysed with SPSS V.10.0. The ethical

committee of the hospital clinic approved the study.

Results

We identified 68 Hu-ab-positive patients (3% of the whole series) with a clinical follow-up of more than 3 years, 49 with and 19 without cancer. Table 1 summarises the clinical features of these patients. In the group without tumour, the youngest patient was a 28-year-old woman with polyradiculoneuropathy and systemic lupus erythematosus (SLE).² The remaining patients had no other autoimmune diseases. A detailed report on the presence of autoimmune disorders, however, was not routinely registered in the databases. A 61-year-old man, with cerebellar ataxia and opsoclonus, had a severe premalignant bronchial dysplasia at initial bronchoscopy, but the mucosa was normal at three subsequent bronchoscopies over the following months. Six patients without cancer died during the follow-up: four from PNS, one from heart disease and one of unknown cause. Postmortem examination was not performed.

In the group of 49 patients with cancer, the tumour was diagnosed before the third year in 46 (94%) patients. The other three patients

were diagnosed with metastatic small-cell lung carcinoma at 40, 41 and 72 months, respectively, after the onset of PNS.

We found no differences between the groups in any of the variables analysed, with the exception of the average duration of follow-up. This was significantly longer in the group without tumour (84 v 68 months, $p = 0.048$). In this group, the follow-up was longer than 72 months in 11 (58%) patients. Because one patient had cancer diagnosed 72 months after the onset of PNS, we compared the clinical variables of those 11 patients without cancer with those of patients with cancer after more than 72 months of follow-up. We found no differences, except for an increase in the time between the onset of PNS and detection of Hu-abs (table 1).

Discussion

In our study, 3% of all Hu-ab-positive patients with PNS did not develop a tumour during prolonged follow-up. We found no clinical features that distinguished the few Hu-ab-positive patients who did not develop cancer from those who did. Our use of the 3-year cut-off was based on previous studies on PNS in which almost all tumours were

Table 1 Clinical characteristics of patients with and without tumour with a follow-up of more than 36 and 72 months

	Tumour*	No tumour >36 months	p Value	No tumour >72 months	p Value
Number of patients	49	19		11	
Age of onset, years (range)†	58 (37-74)	54 (28-81)	0.24	52 (28-68)	0.121
Male	36 (73.5)	10 (52.6)	0.19	5 (45.5)	0.086
Smoker‡	32 (97.0)	15 (83.3)	0.12	9 (90.0)	0.415
Rankin score at diagnosis					
0-3	37 (75.5)	15 (78.9)	1.0	9 (81.8)	1.0
4-5	12 (24.5)	4 (21.1)		2 (18.2)	
Rankin score at follow-ups					
0-3	25 (64.1)	13 (72.2)	0.80	8 (80.0)	0.45
4-5	14 (35.9)	5 (27.8)		2 (20.0)	
Predominant syndrome					
Involvement of CNS¶	6 (12.2)	4 (21.0)	0.26	2 (18.2)	0.60
No involvement of CNS**	43 (87.8)	15 (79.0)		9 (81.8)	
Time between onset of PNS and detection Hu, months (range)	12.8 (0-60)	19.8 (3-65)	0.08	24 (5-65)	0.04

CNS, central nervous system; PNS, paraneoplastic neurological syndrome; SCLC, small-cell lung carcinoma.

Values in parentheses are percentages, unless otherwise stated.

*SCLC, 36; non-SCLC, 6; breast cancer, 3; bladder cancer, 1; ethmoidal epithelioma, 1; thymoma, 1; germ cell tumour, 1.

†Patients under 40 years (1 with tumour, 4 without tumour).

‡Status unknown in 17 patients (16 with tumour, 1 without tumour).

§Rankin score unknown in 11 patients (10 with tumour; 1 without tumour).

¶Tumour group: cerebellar degeneration, 3; limbic encephalitis, 2; encephalomyelitis, 1. Non-tumour group: cerebellar degeneration, 3; brain stem encephalitis, 1.

**Tumour group: sensory neuronopathy, 41; dysautonomia, 2. Non-tumour group: sensory neuronopathy, 11; sensorimotor neuropathy, 3; dysautonomia, 1.