neurological rehabilitation, cognitive performance improved considerably. Seven weeks after the operation, she was orientated in all qualities and initial deficits in attentional performance and executive functions recovered. However, neuropsychological testing at this time revealed an average performance on general and selective attention, whereas the performance on divided attention and on the Wisconsin Card Sorting Test was impaired. The spatial-constructional abilities tested by the Rey-Osterrieth Complex Figure test and the mosaic test of Hamburg Wechsler Intelligence Test for Adults were normal. In the Auditory-Verbal Learning Test, she could initially recall the words in five trials with borderline values but over time or with interference of other words she could not. The storage process for long term memory was more impaired than for short term memory. The activities of daily living were not disturbed.

The postoperative angiogram demonstrating narrowing of the median callosal artery is shown in fig 1D.

MRI performed 1 day after surgery showed a nearly symmetric signal increase of the body of the fornix and of the adjacent part of the genu and anterior body of the corpus callosum on fluid attenuated inversion recovery images (fig 1A, 1B). Diffusion weighted images also revealed hyperintense signal changes with reduced values on apparent diffusion coefficient (ADC) maps using a region of interest (ROI) analysis with 32 voxels (mean (SD): $0.49 (0.04) \times 10^{-3} \text{ mm}^2/\text{s}$; normal value from a separate control subject $0.76 (0.11) \times 10^{-3} \text{ mm}^2/\text{s}$). Diffusion tensor images (DTI: single shot echoplanar imaging sequence with TR 7200 ms; TE 80 ms; 12 non-collinear diffusion directions; 128×128 matrix, voxel dimensions $1.6 \times 1.6 \times 1.6 \text{ mm}^3$; $b = 700 \text{ s/mm}^2$) and fibre tracking were performed 13 days later. Fibre tracking was performed using DTI studio (Johns Hopkins University, Baltimore, Maryland, USA). The propagation algorithms were medium lengths of fibres, $\text{FA} > 0.3$ for each voxel, and line deviated by an angle $> 70^\circ$. DTI revealed an increase in ADC ($1.21 (0.19) \times 10^{-3} \text{ mm}^2/\text{s}$; normal value $0.71 (0.16) \times 10^{-3} \text{ mm}^2/\text{s}$) and a decrease in FA ($0.32 (0.06)$) in the region of interest of the lesion in the fornix whereas ADC ($0.72 (0.16) \times 10^{-3} \text{ mm}^2/\text{s}$) and FA were normal in the body of the fornix outside the lesion ($0.62 (0.13)$; FA value from a separate control subject $0.64 (0.12)$). Fibre tracking using a low threshold (FA 0.3) showed that some fibres could be followed even in the infarction of the anterior fornix (fig 1E, 1F).

Discussion

Amnesia is an uncommon, but increasingly recognised syndrome after damage of the anterior fornices. The fornix forms an integral part of the hippocampal-anterior thalamic pathway mediating memory retrieval, so that lesions of the fornix may selectively impair recall processes.¹ Infarction of the anterior fornices in conjunction with the rostrum and genu of the corpus callosum has been discussed previously by Moussouttas *et al.*² The vascular supply of these structures emanates from the subcallosal or the median callosal artery.³ In our patient, a dominant median callosal artery supplied the fornix. The damage of the basal forebrain, including the septohippocampal complex supplied by the anterior perforation arteries, is a well known complication of

rupture or repair of ACoA aneurysms. However, the presented type of focal infarction has not been reported previously. Amnesic syndromes often complicate the clinical outcome after rupture and surgery of ACoA aneurysms. They are commonly explained by damage of the nuclei of the medial septum and of the diagonal band of Broca, whereas detailed information about the fornices is lacking. The initial disturbances of the motivated behaviour and arousal in our case are explained by the concomitant involvement of the precallous region.⁴ Anterograde amnesia and compromised verbal recall in retrieval processes are described both in fornix infarction and after rupture and repair of ACoA aneurysms.^{2,4,5} The memory impairment of our patient is consistent with this memory profile. Therefore, it may be supposed that fornix infarction after treatment of ACoA aneurysms is a complication that has not been assessed to date.

Changes in ADC values reflect alterations at the cellular level whereas DTI mirrors the functional integrity of axonal structures. Our case is the first in which DTI and diffusion tensor tractography in infarction of the fornix were performed. Thus the significance of the FA changes in this fibre bundle has not been investigated previously. Despite half the reduction of FA in the lesion compared with normal, the clinical outcome as well as fibre tracking indicated that some fibres might be preserved. These findings imply that fibre tracking may predict the outcome after damage to this fibre system. Nonetheless, the tracing of designated fibre bundles has technical pitfalls because of the curved course and the partial volume effect of the adjacent CSF. However, increased signal resulting from the higher field strengths of 3 T in the presented case allows acquisition with smaller voxel sizes, improving substantially the tracking conditions.

Elke Hattingen

Institute of Neuroradiology, Johann Wolfgang Goethe University of Frankfurt/Main, Germany

Julian Rathert, Andreas Raabe

Department of Neurosurgery, Johann Wolfgang Goethe University of Frankfurt/Main, Germany

Angela Anjorin, Heinrich Lanfermann, Stefan Weidauer

Institute of Neuroradiology, Johann Wolfgang Goethe University of Frankfurt/Main, Germany

Correspondence to: Dr Elke Hattingen, Institute of Neuroradiology, University of Frankfurt, Schleusenweg 2-16, D - 60528 Frankfurt, Germany; elke.hattingen@kgu.de

doi: 10.1136/jnnp.2006.109801

Competing interests: None.

References

- 1 Gaffan EA, Gaffan D, Hodges JR. Amnesia following damage to the left fornix and to other sites. A comparative study. *Brain* 1991;114:1297-313.
- 2 Moussouttas M, Giacino J, Papamitsakis N. Amnesic syndrome of the subcallosal artery: a novel infarct syndrome. *Cerebrovasc Dis* 2005;19:410-14.
- 3 Ture U, Yasargil MG, Krisht AF. The arteries of the corpus callosum: a microsurgical anatomic study. *Neurosurgery* 1996;39:1075-84.
- 4 Böttger S, Prosiel M, Steiger HJ, *et al.* Neurobehavioural disturbances, rehabilitation outcome, and lesion site in patients after rupture and repair of anterior communicating artery

aneurysm. *J Neurol Neurosurg Psychiatry* 1998;65:93-102.

- 5 Moudgil SS, Azzouz M, Al-Azzaz A, *et al.* Amnesia due to fornix infarction. *Stroke* 2000;31:1418-19.

Myopathy with tubular aggregates and gyrate atrophy of the choroid and retina due to hyperornithinaemia

Gyrate atrophy (GA) of the choroid and retina with hyperornithinaemia is a rare inherited autosomal recessive metabolic disease. It is due to a mitochondrial enzymatic deficiency in ornithine-amino-transferase (OAT), leading to hyperornithinaemia. Most frequently, ocular findings are clinically isolated and spread to blindness. Associated muscular weakness is rare.

Case report

The present case is that of a patient, from a consanguineous Turkish marriage and the third of seven brothers and sisters, who developed, at age 15 years, loss of night vision with bilateral retinopathy, myopia and cataract. Ophthalmological symptoms progressively worsened. At the age of 18 years, mild proximal weakness appeared in the right arm and slowly spread to the four limbs. At the age of 33 years, he had proximal muscular atrophy of the lower limbs without pain. Clinical examination showed marked bilateral atrophy and mild weakness of the gluteus maximus, and detachment of the shoulder blades. There was no sensitive symptom, reflexes were normal, and there was no sign of central nervous system involvement.

Visual acuity was estimated to be 6/30 (right eye) and 6/15 (left eye). Fundoscopy revealed major chorioretinal atrophy with a typical pattern of gyrate atrophy: garland-shaped, sharply defined, zones of atrophy in the midperiphery of the fundus, and large atrophic areas with festooned edges (fig 1).

Standard biological investigations were normal; creatine phosphokinase serum level was 189 U/l (N: 40-250).

Electromyography showed a myopathic pattern in the proximal muscles.

Histological analysis (no 2349-95 IPS) of the deltoideus muscle revealed subsarcolemmal and intermyofibrillary slits or lakes affecting numerous muscle fibres. These areas were pink on H&E and red on Gomori's trichrome stain. They were strongly stained with nicotinamide dinucleotide tetrazolium reductase and were unstained with succinic acid dehydrogenase. They were exclusively found in type II fibres. On semi-thin toluidine blue-stained resin sections, these masses appeared pale blue, subdivided by thin septae. Electron microscopy showed that these areas consisted of densely packed tubules with a diameter of about 50 nm. Some of these tubules were dilated. This pattern corresponded to tubular aggregates.

Enzymatic activities of the mitochondrial chain and mitochondrial DNA were normal.

Serum and urinary levels of ornithine were very high, respectively, $1300 \mu\text{mol/l}$ (N: 111 ± 64) and $633 \mu\text{mol/mmol creatinine}$ (N < 5).

Cerebral MRI was normal.

The diagnosis of myopathy with tubular aggregates and GA of the choroid and retina due to hyperornithinaemia was made, and

Figure 1 Fundoscopy of gyrate atrophy.

treatment with daily oral pyridoxine was started and a low arginin diet ensured.

Discussion

Two genetic autosomal recessive disorders result from hyperornithinaemia: hyperornithinaemia-hyperammonaemia-homocitrullinuria syndrome, and GA of the choroid and retina with hyperornithinaemia. GA of the choroid and retina is a rare disease, first reported in the Finnish population, but now described all over the world.¹

The main clinical feature of hyperornithinaemia is its ocular abnormalities. First symptoms are non-specific and appear during the second or third decade, consisting of hyperopia, myopia and constricted visual field. Fundoscopy generally reveals a bilateral typical pattern of festooned chorioretinal atrophy, initially occurring in the midperiphery, and posterior subcapsular cataract.² If not treated, this condition can lead to blindness between the ages of 40–55 years.

Plasma and urinary ornithine levels, measured by chromatography, are constantly raised (5–20 N).

Ornithine is an amino acid that plays a role in the metabolism of urea, creatine and polyamines and that can be consumed during a reaction catalysed by the OAT in the mitochondria.

Hyperornithinaemia is caused by OAT deficiency secondary to OAT gene mutation. More than 50 pathogenic mutations of the OAT gene, mapped in 10q26, have already been detected.³ Eighty-five per cent of patients with gyrate atrophy have a normal amount of normal-sized OAT mRNA, but only 10% have normal amounts of normal-sized OAT protein. This finding suggests a normal transcription but abnormal transduction.

Systemic involvement, such as thin and rare hairs, mental retardation with diffuse brain atrophy, slow background electro-encephalographic abnormalities and muscular weakness, has been described previously in association with GA.⁴ Muscular histological examination, even in patients without weakness, can show muscular abnormalities such as type II fibre atrophy and sarcoplasmic tubular aggregates. The pathogenic role of tubular aggregates is unknown and uncertain. These distinctive inclusions are not specific and have been described in normal subjects, in some other muscular diseases, such as acromegaly, and in porphyria cutanea tarda.

An arginine-restricted diet can reduce plasma ornithine levels and completely prevent retinal degeneration, over a 12-month period, in a mouse model of OAT deficiency.⁵ Long-term reduction of ornithine levels by the

arginine-restricted diet has been proposed in patients with GA; progression of chorioretinal lesions can be slowed if the diet is started at an early age.⁶

Administration of pyridoxine stimulates residual OAT activity and is associated with a significant reduction in ornithine plasma levels in some genetically determined responsive patients. The daily dose of pyridoxine is still under discussion (between 20 and 600 mg). However, the pyridoxine therapy does not stop the progression of chorioretinal degeneration.

Conclusion

GA with hyperornithinaemia is a metabolically inherited disease and is one of the rare causes of both ocular and muscular involvement. Lack of specificity of initial ophthalmological symptoms can be responsible for a misdiagnosis. When muscular symptoms are associated with these ophthalmological signs, the diagnosis of GA of the choroid and retina due to hyperornithinaemia has to be considered and ornithine plasma levels have to be assayed.

M Fleury, R Barbier

Department of Neurology, Hôpital civil de Strasbourg, Strasbourg, France

F Ziegler

Department of Neurology, Hôpital de Belfort, Belfort, France

M Mohr

Department of Pathology, Hôpital civil de Strasbourg, Strasbourg, France

O Caron

Department of Ophthalmology, Hôpital civil de Strasbourg, Strasbourg, France

H Dollfus

Department of Medical Genetics, Université Louis Pasteur, Strasbourg, France

C Tranchant, J M Warter

Department of Neurology, Hôpital civil de Strasbourg, Strasbourg, France

Correspondence to: M Fleury, Department of Neurology, Hôpital civil de Strasbourg, 1 place de l'hôpital, 67091 Strasbourg, France; marie-celine.fleury@chru-strasbourg.fr

doi: 10.1136/jnnp.2006.101386

Published Online First 6 November 2006

Competing interests: None declared.

References

- 1 Cutler CW. Drei ungewöhnliche Fälle von retino-choroidal Degeneration. *Arch Augenheilk* 1895;30:117–22.
- 2 Wilson DJ, Weleber RG, Green WR. Ocular clinicopathologic study of gyrate atrophy. *Am J Ophthalmol* 1991;111:24–33.
- 3 Brody LC, Mitchell GA, Obie C, et al. Ornithine delta-amino-transferase mutations in gyrate atrophy. Allelic heterogeneity and functional consequences. *J Biol Chem* 1992;267:3302–7.
- 4 Kaiser-Kupfer MJ, Kuwabara T, Askonas V, et al. Systemic manifestations of gyrate atrophy of the choroid and retina. *Ophthalmology* 1981;88:302–6.
- 5 Wang T, Steel G, Milam AH, et al. Correction of ornithine accumulation prevents retinal degeneration in a mouse model of gyrate atrophy of the choroid and retina. *Proc Natl Acad Sci USA* 2000;97:1224–9.
- 6 Kaiser-Kupfer MJ, Caruso RC, Valle D. Gyrate atrophy of the choroid and retina: further experience with long-term reduction of ornithine levels in children. *Arch Ophthalmol* 2002;120:146–53.

Stiff person syndrome (SPS), a basal ganglia disease? Striatal MRI lesions in a patient with SPS

Stiff person syndrome (SPS) is characterised by muscle rigidity and episodic spasms involving the axial and limb musculature. The hallmark is co-contraction of agonist and antagonist muscles, with continuous motor unit firing at rest. An autoimmune pathogenesis is based on the presence of anti-glutamic acid decarboxylase antibodies (anti-GAD, rate-limiting enzyme to synthesise γ -aminobutyric acid (GABA), in presynaptic GABAergic inhibitory synapses), and association with other autoimmune disorders and autoantibodies, and on response to immunotherapy. There is only a limited number of cases with brain MRI abnormalities. Our patient is the first reported case of SPS with MRI striatal abnormalities.

Case report

A 69-year-old woman came in for consultation with an 18-month history of back pain, stiffness and progressive difficulty at walking. She had developed asymmetrical rigidity in her legs, with the left one being worse, from falls. In one of these falls she broke her left humerus. That limb remains useless, painful and rigid. She had progressive difficulty in turning over in bed and in getting up. In stressful situations and when trying to move, she had painful spasms in the left leg starting distally with toe and ankle extension, and progressing to knee and hip flexion while raising the leg from the bed. She became progressively bedridden. Her medical history was notable for diabetes mellitus (controlled with acarbose) and hypertension (controlled without drugs). A neurological examination showed stiffness and limitation in the range of active and passive movements in the left leg, left arm and, less pronouncedly, in the right leg. Also, there was slowing of finger, hand and arm movements in the left hand. She found it difficult to raise her left leg, which sometimes developed into painful spasms. She had decreased facial expression, cogwheel rigidity in both superior extremities and stony lumbar rigidity. Plantar reflexes were flexor and deep tendon reflexes were symmetrically brisk. She needed help to sit down and was afraid to walk even with aid. On the other hand, her mental status, speech and cranial nerves were normal.

The results of blood tests, including anti-nuclear antibody, endomysial, gliadin, transglutaminase, mitochondrial, smooth muscle, parietal, anti-LKM1 (autoimmune hepatitis), thyroid peroxidase and thyroglobulin, were negative. Levels of folic acid, vitamin B₁₂, T4, thyroid-stimulating hormone, rheumatoid factor, C-reactive protein, anti-streptolysin O, immunoglobulins and complement were normal. Serum anti-GAD level was 14 000 U/ml (radioimmunoassay). Also, the patient tested negative for anti-Yo, anti-Hu, anti-Ri and antiampiphysin. Serum anti-GAD increased to 60 400 U/ml (N<1 U/ml).

The patient rejected lumbar puncture. Nerve conduction was normal and electromyography showed normal motor unit potentials, an inability to relax any of the muscles tested, and continuous motor unit activity.

Cervical and lumbar MRI was normal. Cerebral MRI showed bright hyperintense changes on fluid-attenuated inversion recovery (FLAIR) and T2 in both striatal regions (more intense in the right striatum corresponding to