kg.) of body weight gave adequate resistance. Apart from its influence on the dosage, however, the time factor is of importance, since it appears that immunization, to be effective, should be given earlier than six days before the onset of symptoms.

Although gamma globulin has been used to control outbreaks of infective hepatitis in America, this appears to be the first occasion on which it has been used in this country. Indeed, as yet there has been no evidence to indicate that gamma globulin obtained from blood donors here contains sufficient antibodies to be of any immunizing value. The outbreak described, while in no way proving the efficacy of gamma globulin, does, however, suggest that it is a potent agent in controlling the spread of infective hepatitis. This is of considerable economic value in closed communities where infective hepatitis entails the restriction of admissions; and it may be that increased supplies will enable the use of gamma globulin as a protective agent to be more fully exploited in the future.

I should like to thank Dr. C. M. Smith, county medical officer, for his advice and help in preparing this report.

M. J. PLEYDELL, M.C., M.D., D.P.H.,
Deputy County Medical Officer of Health,
Northamptonshire.

REFERENCES

- Havens, W. P., jun., and Paul, J. R. (1945). *J. Amer. med. Ass.*, 129, 270.
Stokes, J., jun. (1953). *Amer. J. publ. Hlth*, 43, 1097.
— and Neefe, J. R. (1945). *J. Amer. med. Ass.*, 127, 144.

Scurvy

Frank scurvy is now almost a clinical curiosity in this country. The following case report is therefore of interest and worthy of publication.

CASE REPORT

A postman, aged 56, was admitted to the Royal Surrey County Hospital on July 22, 1952. He had lived in southern England all his life, except for one short period of service with the Army in France during the 1914-18 war, and up to the onset of his present illness had always enjoyed good health.

The present illness started in 1945, when he began to experience epigastric pain occurring approximately one hour after meals. A barium-meal x-ray examination confirmed the presence of a gastric ulcer and he was given medical treatment in a local cottage hospital. With treatment he became completely symptom-free, and after discharge he decided to restrict himself to a diet consisting mainly of steamed fish, eggs, bread-and-butter, and milk puddings. He avoided all fruit and vegetables except for an occasional small quantity of mashed potatoes. At no time was his diet supplemented with vitamins. He remained very fit on this diet until two weeks before admission, when he began to complain of lethargy and aching in the muscles of the legs. After approximately 48 hours the legs became swollen and bruises appeared; these slowly spread so as to involve both legs diffusely. At no time has he complained of epistaxis or soreness of the gums, but on close questioning he did admit to some dyspnoea on exertion.

The patient stated that he thought his illness had been precipitated by the recent death of his wife, which, he said, had caused him considerable worry and extra work.

On examination he was pale and of average stature. There was a low-grade pyrexia of 99.6° F. (37.6° C.), with a pulse rate of 90. He was edentulous, but the mouth, throat, and tongue appeared quite normal. In the chest, abdomen, and central nervous system there were no definite abnormal physical signs, and in the cardiovascular system the only abnormality discovered was a markedly positive Hess capillary fragility test. The skin on the upper parts of the arms and thighs showed evidence of early follicular hyperkeratosis, and the legs were grossly discoloured by

a haemorrhagic rash in which three types of lesion could be identified. These can best be described as small circum-follicular purpuric spots, and larger raised, red, tender plaques approximately 2 to 3 cm. in diameter; there were also large subcutaneous extravasations of blood involving the major part of the feet, legs, and thighs.

Apart from the individual lesions mentioned above, the legs were not tender at the time of examination, although the patient did state that the calves had been tender prior to admission.

On investigation the following results were obtained: haemoglobin, 41% (6.56 g.); R.B.C., 2,200,000; colour index, 0.93; M.C.D., 7.1 μ ; W.B.C., 4,200 (73% polymorphonuclears, 25% lymphocytes, 2% monocytes). The platelet count was 280,000 per c.mm. and the prothrombin index 83%. The bleeding-time was 2½ minutes, and the clotting-time 9½ minutes (venous blood). The blood Wassermann reaction and Kahn test were negative. The urine contained a trace of albumin and small numbers of red cells and leucocytes, but no casts were seen and culture was sterile. The stool was cultured for pathogens with a negative result. An x-ray film of the tibiae revealed no bony or periosteal abnormality.

An ascorbic acid saturation test was performed, the technique of which was to give 10 mg. of ascorbic acid per kg. body weight at 9 a.m., followed by a determination of the total urinary excretion of ascorbic acid between 1 and 3.15 p.m. It is stated that a fully saturated person should excrete 50 mg. or more during this 2½-hour period.

Synchronous with this test a daily reticulocyte count was done in order to determine the haematological response to ascorbic acid only, no iron being given. The results are shown in the Table.

Day	Ascorbic Acid Excreted in 2½-hour Period	Reticulocyte Count
1	1.3 mg.	1.8%
2	1.92 "	2.2%
3	2.32 "	3.4%
4	4.27 "	6.6%
5	3.97 "	4.2%
6	8.26 "	7.0%
7	8.77 "	8.4%
8	31.3 "	8.6%
9	64.0 "	8.6%
10	50.0 "	11.7%

After the completion of the saturation test, treatment was continued with ascorbic acid. By the time of his discharge, after three weeks in hospital, the patient was feeling extremely fit, the lesions on the legs had completely resolved, and his haemoglobin had risen to 80% (12.8 g.). He has since been seen again as an out-patient, and has remained quite well.

COMMENT

The diagnosis in the above case can be in no doubt, as the clinical findings, the ascorbic acid saturation test, and the response to treatment were all confirmatory.

Although subclinical deficiency of vitamin C is not uncommon in this country, scurvy is rare. If one considers the possibility that a number of people may be living on inadequate self-imposed diets, such as our patient was doing, this is perhaps surprising. One can only suppose, however, that very few people, except obsessionists, would in fact avoid all fruit and vegetables so completely as to cause frank scurvy. Nevertheless, the possibility of scurvy occurring in a patient on a diet is a very real one, and consideration should therefore always be given to the vitamin content in any prescribed diet.

I wish to thank Dr. H. Bell, under whose care the patient was admitted, for permission to publish this case.

H. R. C. RICHES, M.B., B.S., M.R.C.P.,
Medical Registrar, Royal Surrey County Hospital, Guildford.