BRITISH MEDICAL JOURNAL

LONDON SATURDAY SEPTEMBER 11 1954

THE DORMANT CANCER CELL*

BY

GEOFFREY HADFIELD, M.D., F.R.C.P.

*Sir William Collins Professor of Pathology and Dean of the Institute of Basic Medical Sciences,
Royal College of Surgeons of England*

[WITH SPECIAL PLATE]

The time elapsing between the excision of a human malignant tumour and the appearance of a clinically recognizable recurrence varies from a few months to thirty years. It is usually short, being less, sometimes appreciably less, than a year or so, and when of this order of duration it is probable that the secondary tumours are the product of continuous and unsuppressed proliferation of the cells of microscopic fragments of growth, some of which are portions of the primary growth which have escaped excision, whilst others had already reached a distant organ or tissue by lymphatic, venous, or arterial embolization before the tumour was removed.

Temporary Mitotic Arrest

When the interval between excision and the appearance of secondary tumours exceeds two and reaches four or more years, it becomes increasingly difficult to believe that cellular proliferation in the residual tumour fragments has been continuous and uninhibited. We cannot, for instance, assume that these longer intervals are characteristic of slowly growing tumours, because remarkably long intervals have often been described in tumours whose original and subsequent rates of growth were very rapid. When the interval is prolonged to six years or more it seems impossible to escape the conclusion that the cells of the dormant growth are in a state of *temporary mitotic arrest*, no matter how long the period may be. This conclusion becomes all the more probable when we witness the clinical resurrection of these minute foci of dormant cancer cells after a long period of latency and satisfy ourselves that they have lost none of their former capacity to grow rapidly and infiltrate widely.

In cancer literature there are many accounts of instances in which the clinically silent periods of mitotic arrest have been remarkably long, and the following table is a summary of some of these reports gathered together by Willis (1948).

Site	Latent Period
Breast (carcinoma)	6 to 20 years
Cutaneous and ocular melanoma ..	14 to 32 "
Kidney (carcinoma)	6 to 8 "
Tongue (carcinoma)	8 to 15 "
Stomach and colon (carcinoma) ..	5 to 6 "

Long latent periods of this order are certainly uncommon in the natural history of malignant disease. The usual course, although often punctuated by short intermissions, is one of continuous cellular proliferation and progressive clinical deterioration, its length being

primarily determined by the average growth rate of the tumour cells.

Instances in which the clinical course shows an uncommonly long period of mitotic arrest seem to be reported in cancer literature only when the latent period ending in recurrence is exceptionally long. In one sense this practice is misleading, for periods varying from three to six years are far from exceptional in the natural history of malignant tumours, and I find it difficult to escape the conclusion that during periods of this duration the tumour cells are in the same state of mitotic arrest as they almost certainly are when the period is exceptionally long.

In any case, the phenomenon of temporary growth suppression in tumour pathology is of obvious biological importance, and for the sake of brevity in choosing a title for this lecture I have used the word "dormant" to describe malignant cells which, although remaining alive in the tissues for relatively long periods, show no evidence of multiplication during this time, yet retain all their former and vigorous capacity to multiply.

Anatomical Site of Dormant Growths

Local Recurrence

There are three situations in which such dormant cancer cells may exist. Some lie in or near the operation site, where it is highly probable that they are involved in post-operative scar. When they eventually recur they produce secondary growths which not infrequently lie in close relationship with the skin incision or its deep extensions. Here they exist for long periods of time in poorly vascularized mature collagenous connective tissue, an environment which possibly subjects them to long periods of relative anoxia. Other foci of dormant growth may lie in regional lymph nodes, where in some growths, especially those of the breast, they may again survive but fail to multiply in a fibrous and possibly anoxic environment. It may not be unreasonable to assume, therefore, that in these situations mitotic arrest has been induced by anoxia or some similar and closely related metabolic deficiency, and, alternatively, that mitosis would be re-established if adequate oxygen became available by vascularization of the scar, an event which may be brought about by an inflammatory response to one or more of many varieties of local injury, possibly including irradiation.

Metastatic Growth

On the other hand, we have no information which might help us to understand the induction of the dormant state and the reactivation of microscopic foci of metastatic tumour tissue in deeply placed organs. Such foci must have been originally implanted before excision of the primary growth. In all probability they were small enough to have arisen

*Kettle Memorial Lecture delivered at the Royal College of Surgeons of England on December 14, 1953.

as the result of malignant embolism of small arteries and arterioles, and if these emboli were impacted in the systemic arterial circulation the tumour fragments must have either directly traversed the lung capillaries or have broken away from a far larger pulmonary venous metastasis which itself becomes dormant. The first of these alternatives would appear to be far the more probable.

The anatomical site of these minute quiescent growths must be significant, for in at least 50% of cases of malignant disease recurring after latent periods varying from five to thirty years reactivated metastatic growth of macroscopic size was found in the haemopoietic bone marrow. In 1912, E. H. Kettle, to whose memory this lecture is dedicated, drew attention to the presence of microscopic foci of metastatic carcinoma in the pulp of the normal-looking unenlarged spleen. He proposed that such growths lying in the territory of the portal vein may be the source of hepatic metastases when the primary growth lies outside the portal area.

More recently, Clifton and his co-workers (1952) have reported that in 33% of a series of 57 cases of carcinoma of the prostate carcinoma cells were demonstrated in the sternal bone marrow. The cells occurred with equal frequency in marrow from the sternum and ilium, and in 9% there was widespread skeletal dissemination of growth with no diagnostic evidence other than that provided by the marrow biopsy. Evidence of this sort provides the hint that during the period of blood-borne invasion of the systemic organs by tumour cells malignant emboli tend to occur in organs such as those of the reticulo-endothelial system, the hypophysis, and the adrenal cortex, all of which have a sinusoidal capillary circulation. I have a strong impression that the phenomenon of mitotic arrest for long periods is rare in the kidney, lung, brain, and skeletal muscle. On the other hand, there seems to be little doubt that unsuspected microscopic metastases may not infrequently be present in the adrenal cortex, at least in carcinoma of the breast. I have personally seen, or have had reported to me by competent observers during the last 18 months, at least seven instances in which such metastases have been found in surgically excised adrenal glands which are neither enlarged nor deformed. During this period similar latent metastases have been observed in five surgically removed normal-looking ovaries in patients suffering from the same disease.

However suggestive these observations may be, it must be conceded that our present clinical and pathological knowledge of human tumour pathology provides us with no reliable information regarding the environmental factors which induce mitotic arrest of tumour cells in human tissues for periods varying from three to thirty years, and it seems improbable that morphological and histological studies will ever do so. We must inquire, therefore, if fundamental cancer research can or is likely to help our understanding of this remarkable phenomenon.

Experimental Induction of Dormant State

I would suggest that the investigations which have grown out of studies on the low-temperature preservation of animal tumours might supply us with relevant information. The cells of certain normal tissues and those of some malignant tumours show remarkable powers of survival after long periods of freezing to and below -70°C . Michaelis (1905) demonstrated that the cells of the Jensen carcinoma were still viable and would produce tumours in animals after exposure to liquid air for 30 minutes. Shortly afterwards Paul Ehrlich (1907) found that several malignant tumours in animals survived temperatures between -25°C . and -30°C . for 48 hours, and one tumour kept at -8°C . to -10°C . was still viable after two years.

Several observers (Cramer, 1930; Auler, 1932; Klinker, 1939) confirmed these early observations. Barnes and Furth (1937) made the important observation that the survival rate of frozen tumour cells was higher if the tumour was frozen slowly, and Breedis and Furth (1938), using sarco-

mata and the tumour tissue of fowl leukaemia, demonstrated tumour survival for 430 to 448 days at a temperature of -70°C . An animal carcinoma mammae was still viable after 98 days. They were satisfied that their results were due to survival of tumour cells as opposed to a virus or virus-like agent, for after irradiation of the frozen tumour at an intensity which was lethal to its cells but was considered to be non-lethal to any known virus particles their tumours failed to grow on homologous transplantation. The hypothesis that freezing kills tumour cells but spares virus was postulated by Gye (1949) and supported by Mann and Dunn (1949). It was asserted by Craigie that a survival rate, after freezing, of one in a million cells was sufficient to propagate a tumour, while Passey and Dmochowski (1950) demonstrated, by using tissue culture as a test of viability, that transmission was due to cell survival. They also demonstrated that the cell-free supernatant of a centrifugalized thawed-out tumour suspension was incapable of transmitting the tumour, whilst the cellular deposit did so with ease.

The Trauma of Freezing

The therapeutic use, after long periods of preservation in the frozen state, of red blood cells, spermatozoa, bone, skin, and other tissue grafts, as well as large arterial trunks, has called for detailed investigation into the physico-chemical factors involved. Luyet and Gehenio (1940) and Luyet (1951) made the assumption that trauma to cell and nuclear membranes inflicted by ice crystals is a major lethal factor when cells are exposed to low temperatures, possibly because laceration of either membrane would allow nuclear or cytoplasmic material to escape. They decided that cooling and thawing must be carried out so rapidly that the cell contents undergo non-crystalline, amorphous consolidation or vitrification.

The conditions governing this ideal physical transformation are that the rate of cooling must be of the order of several hundred degrees per second to a temperature of -40°C ., the thickness of the material less than 0.1 mm., and its water content less than 70%. These conditions appear to be impossible of practical achievement when dealing with tumour material whose viability must be subsequently determined by grafting. Furthermore, in the case of tumour tissue, a significantly higher survival rate is obtained by slow freezing and rapid thawing than the reverse (Breedis, Barnes, and Furth, 1937; Breedis and Furth, 1938; Craigie, 1949a, 1949b, 1949c, 1954). It is quite conceivable that slow freezing enables tumour cells to effect certain physico-chemical and structural adjustments which seem to be essential if they are to resist anoxia and dehydration. It appears, therefore, that the vitrification hypothesis may not be applicable to tumour cells and the lacerating trauma inflicted on cell membranes by ice crystals is not so lethal a factor as was once thought.

Dehydration at Low Temperatures

It is, in fact, becoming increasingly probable that one of the major factors in the trauma produced by freezing is the inevitable loss of cell water by simple exosmosis, which is clearly due to the development of a hypertonic concentration of inorganic ions in the extracellular fluid as its water crystallizes. For example, a solution of sodium chloride having a concentration of 0.85% at 15°C . reaches a concentration of 30% at -18°C .

Whilst the capacity of a frozen cell to withstand the physico-chemical effects of cold appears to decide whether it shall survive or perish, experimental evidence strongly suggests that a certain sublethal degree of dehydration is an essential prerequisite for survival. It has been shown that extreme degrees of dehydration can be avoided and an optimal degree established by eliminating electrolytes from the fluid in which the tumour cells are suspended, and substituting for them a soluble non-electrolyte. Glycerol has been extensively used for this purpose, and Craigie (1949a, 1949b, 1949c) has successfully employed solutions of glucose

in which the tumour cells are preconditioned before freezing. This, he finds, affords excellent protection for tumour cells against the physical hazards of freezing in slightly hypertonic concentrations of between 6 and 7%. When tumour cells are preconditioned in this way, slowly frozen at -70°C . or lower, stored at this temperature, rapidly thawed, and immediately examined, all the surviving and viable cells are found to be spherical in shape and to have suffered a significant reduction in volume, which rapidly increases to normal as the temperature rises. Using this technique, Craigie (1954) demonstrated that S 37, a highly malignant sarcoma of mice, was still viable and highly malignant after continuous storage at -70°C . for three years and seven months; the Daels guinea-pig tumour, a sarcoma originally induced by radium, survived for two years and five months, and the Walker rat sarcoma for one year and nine months. Neither of these tumours had lost its former capacity to grow rapidly and metastasize widely.

Growth in Peritoneal Cavity

When subjecting malignant tumours to low temperatures it is obviously desirable that they should be first reduced to a homogeneous suspension of single living cells in a suitable fluid. If this is attempted by the usual technique of mincing the solid tumour, a high proportion of its cells are killed and their autolytic products are certainly toxic to the cells which survive. This can be avoided by using the familiar *ascites tumour*, whose cells grow freely in a peritoneal exudate. Craigie (1953) has shown that certain solid mouse sarcomata can be forcibly converted to ascites tumours by intraperitoneal inoculation, and when so converted can be readily propagated as ascites tumours. Tumour cells grown in this way show a high rate of survival when frozen, and this is largely due to eliminating the trauma of "mincing."

Resistance of Normal Cells to Low Temperatures

The ability to survive in a frozen environment without any loss of former reproductive power is a property which the cells of some malignant tumours share with certain normal cells. Breedis and Furth (1938) demonstrated survival of tracheal epithelium after freezing for 327 days; Mider and Morton (1939) successfully grafted rat skin after prolonged freezing at -74°C . Webster (1944) proved that human skin grafts were viable after refrigeration at -72°C . Strumia and Hodge (1945), using autogenous split-thickness grafts of human skin, demonstrated survival after 1 to 61 days at -20°C . to -25°C . They obtained 80% of permanent "takes" with 41 frozen grafts, against 86% in unfrozen controls. Blumenthal and Walsh (1950) showed that thyroid and parathyroid epithelium were viable after freezing at -75°C . Similar results were obtained with ovarian granulosa cells of the rabbit (Smith, 1952), rat ovarian tissue (Parkes and Smith, 1953), and Malpighian cells and epidermal melanoblasts (Billingham and Medawar, 1951). Spermatozoa are highly resistant to freezing. Macrophages and other mesenchymal elements frozen with tumour tissue survive for long periods (Craigie, 1954).

Morphological Changes

Exposure to low temperatures induces a series of remarkably interesting morphological changes in the cells of several malignant tumours which after suspension in 6.5% glucose, followed by slow freezing, are able to survive low temperatures for long periods. These have been studied by Craigie and described in a series of papers (Craigie, 1950, 1952a, 1952b, 1952c, 1954).

The Paramorphic Cell

If tumours which have been maintained at -70°C . for many months are rapidly thawed and quickly examined, cells possessing clearly defined nuclei may be seen which from their appearance show little obvious abnormality and are often partially differentiated. Craigie asserts that these cells are dead, disintegrate at 37°C ., and are incapable of trans-

mitting the tumour on grafting. In addition to these, a very variable number of smaller and highly refractile spherical bodies are seen in which no nuclei are visible, and whose plasma membrane is highly refractile and appears to be thickened (see Plate, Fig. 1). Craigie at first thought that these "spherical bodies" were dead cells, but when suspended in oxygenated Tyrode solution at 37°C . they become hydrated, and by throwing out pseudopodia and becoming mobile declare themselves as living cells. The invisibility of their nuclei may be the result of the inevitable loss of nuclear water during freezing, the nuclear refractive index closely approaching that of the surrounding cytoplasm. The refractile and apparently thickened plasma membrane is revealed by phase-contrast microscopy to be an optical artifact due to the formation of a bright light halo which invests the cell externally and intervenes between the plasma membrane and the cytoplasm internally (Fig. 1).

It is probably highly significant that these cells show intense absorption of ultra-violet light at 2,536 Å, so that when photographed at this wavelength the light absorption is so great that their photographic image is almost black. It is reasonably certain that this phenomenon indicates a heavy intracellular concentration of cytoplasmic pentose-nucleic acid and is possibly due to loss of cell water. It may be that this heavy concentration of large intracellular molecules acts as a biological "anti-freeze" mixture which prevents the crystallization of residual cell water. In any case, it seems safe to assume that an optimal degree of intracellular dehydration is an essential prerequisite for cell survival.

Craigie describes this small spherical refractile cell as "paramorphic," and, whilst the reasons given for its peculiar morphology are to some extent speculative, there is little, if any, doubt that of all the varied cell types present in a malignant tumour the capacity to resist and survive anoxia, dehydration, and metabolic arrest is strictly limited to the paramorphic cells it contains. This belief rests on the demonstration that *the tumour-inducing activity of a thawed-out malignant tumour after a long period of anoxia is directly related to the number of paramorphic cells present in it.*

At 37°C ., under isotonic conditions and access to free oxygen, the paramorphic cell increases in volume, becomes mobile, and its refractility lessens. If frozen in this phase it may again lose volume and assume the characters of the resistant paramorphic cell. If kept at 37°C . refractility disappears, the nucleus becomes visible, mobility is lost, and the degree of absorption of ultra-violet light progressively falls. If frozen in this phase reversion to the paramorphic state is improbable. If kept at 37°C . with ample available oxygen, differentiation of structure begins and the cell becomes capable of mitosis (see Plate, Fig. 2), but if exposed to low temperatures in this phase reversion to the paramorphic state is extremely unlikely and in all probability impossible.

Experimental conditions which govern the induction of the paramorphic state in the cells of malignant tumours, and the biological significance of the morphological changes described, need confirmation and amplification, but within these limitations it may be permissible to make the following generalizations:

1. When defended against the harmful physical effects of cold the cells of some malignant tumours have a remarkable capacity to survive the anoxia and metabolic arrest induced by exposure to low temperatures *in vitro*.
2. After long periods in the frozen state certain cells in these tumours retain their full reproductive vigour and malignant potentialities when grown in tissue culture or transplanted into the tissues of animals of the same species.
3. The capacity of a frozen tumour to survive appears to be strictly limited to its undifferentiated cells.
4. These cells share this attribute with certain cells of normal tissues.
5. Tumour cells capable of surviving undergo rapidly a series of characteristic morphological changes when subjected to low temperatures.

6. Some or all of these morphological changes may be the direct consequence of the operation of purely physical factors, such as dehydration. On the other hand, the ability of a tumour cell to undergo these modifications of structure appears to decide if the cell shall survive or perish in an anoxic environment, and if it survives whether it shall retain its former capacity to multiply and to initiate a race of cells which infiltrate and metastasize in the tissues of an animal of the same species as the original host.

At the beginning of this lecture I described the naturally occurring phenomenon of total mitotic arrest in human malignant tumours. On the evidence provided by our know-

FIG. A.—Tumour mass in mouse six days after grafting cells of a benzpyrene-induced sarcoma which had been kept at -70°C . for nearly five years.

ledge of human tumour pathology this phenomenon seems to fall within the category of unsolved and insoluble biological mysteries. I now invite you to consider whether the experimental induction *in vitro* of a dormant state in tumour cells may be germane to this problem. Your first reaction to this suggestion is likely to be unsympathetic. You may regard the changes which take place in frozen tumour cells as being part of a strictly specific reaction to cold.

You may look upon any comparison between the human cancer cells becoming "dormant" at body temperature and the animal tumour cell becoming "paramorphic" at -70°C . as being so strained and artificial that it hardly merits serious consideration.

May I remind you that tumour cells have been kept alive in a frozen and anoxic state for periods of time which are obviously comparable to those during which tumour cells remain temporarily dormant in human tissues, that this has been achieved by deliberately defending them against the lethal physical effects of cold, and that such cells emerge from the frozen state with no apparent deterioration in their biological or malignant characteristics? I would therefore like to suggest that when the preconditioned tumour cell is very cautiously frozen it suffers no permanent injury, and that the only demonstrable effect of a cold environment upon it is temporarily to inhibit its intracellular enzyme systems without any impairment of their potential efficiency.

If these suppositions prove to be true, and they certainly need confirmation, the experimentally produced paramorphic tumour cells of transplantable animal cancer can be regarded as existing in a biological state closely related to, or possibly identical with, the naturally occurring "dormant cancer cells" of human pathology. By investigating the paramorphic cell we may therefore learn something of the biological state in which these dormant cells survive in human tissues before they begin to proliferate.

The fact that malignant cells can be induced to survive long periods of experimental metabolic arrest would appear to offer an explanation of their *in vivo* ability to withstand separation from their parent growth until a capillary circulation is provided for them. This may well apply to the microscopic foci of isolated cells which form the nucleus of a metastasis, or to groups of cells of comparable size which become completely detached from their parent growth during the process of malignant infiltration. In conclusion,

I would like to claim that the preconditioning techniques employed to obtain maximum survival of cells in the frozen state provide us with a valuable experimental method which, without inflicting any permanent cytological injury, enables us temporarily to suppress the metabolic activities of the cells of malignant tumours.

ADDENDUM

Since this lecture was delivered, another series of malignant tumours, stored at or below -70°C . for a remarkably long period, have been thawed out and tested for viability. I am permitted by Dr. J. Craigie, Director of the Imperial Cancer Research Fund's Laboratories at Mill Hill, to report the following details concerning them:

A Fresh Series

There are six sarcomatous tumours in this series.

They were originally induced by the subcutaneous injection of benzpyrene into mice. All were found to be transmissible by grafting and all were sarcomatous, but each of them differed significantly from the others in cytological structure.

The cells of these tumours were suspended in dextrose and frozen at -70°C . on April 12, 1949. They remained undisturbed at this temperature until February 8, 1954—a period of 4 years 9 months 27 days. They were then thawed out and grafted into mice.

All the tumours grew vigorously (see Fig. A and Plate, Fig. 3) and had suffered no loss of viability while frozen, for they became visible to the naked eye after a substantially shorter period than those produced in animals of a control series.

REFERENCES

- Auler, H. (1932). *Z. Krebsforsch.*, **35**, 103.
 Barnes, W. A., and Furth, J. (1937). *Amer. J. Cancer*, **30**, 75.
 Billingham, R. E., and Medawar, P. B. (1951). *Freezing and Drying*, pp. 55-62. Symposium held June, 1951, Institute of Biology.
 Blumenthal, H. T., and Walsh, L. B. (1950). *Proc. Soc. exp. Biol., N.Y.*, **73**, 62.
 Breedis, C., Barnes, W. A., and Furth, J. (1937). *Ibid.*, **36**, 220.
 — and Furth, J. (1938). *Science*, **88**, 531.
 Clifton, J. A., Philipp, R. J., Ludovic, E., and Fowler, W. M. (1952). *Amer. J. med. Sci.*, **224**, 121.
 Craigie, James (1949a). *Brit. J. Cancer*, **3**, 249.
 — (1949b). *Ibid.*, **3**, 268.
 — (1949c). *British Medical Journal*, **2**, 1485.
 — (1950). 47th Ann. Rep. I.C.R.F., 1949-50, pp. 5-18.
 — (1952a). *J. Path. Bact.*, **64**, 251 (abstract).
 — (1952b). 49th Ann. Rep. I.C.R.F., 1951-2, pp. 5-9.
 — (1952c). *Ann. roy. Coll. Surg. Engl.*, **11**, 287.
 — (1953). 50th Ann. Rep. I.C.R.F., 1952-3.
 — (1954). *Advances in Cancer Research*, vol. 2. In press.
 Cramer, W. (1930). 9th Scientific Rep. I.C.R.F., pp. 21-32.
 Ehrlich, P. (1907). *Z. Krebsforsch.*, **5**, 59.
 Gye, W. E. (1949). *British Medical Journal*, **1**, 511.
 Kettle, E. H. (1912-13). *J. Path. Bact.*, **17**, 40.
 Klinker, J. (1939). *Growth*, **3**, 169.
 Luyet, B. J. (1951). *Freezing and Drying*, pp. 77-98. Symposium held June, 1951, Institute of Biology.
 — and Gehenio, P. M. (1940). *Biodynamica*, **3**, 33.
 Mann, I., and Dunn, W. J. (1949). *British Medical Journal*, **2**, 255.
 Michaelis, L. (1905). *Med. Klin.*, **1**, 202.
 Mider, G. B., and Morton, J. J. (1939). *Amer. J. Cancer*, **35**, 502.
 Parkes, A. S., and Smith, A. U. (1953). *Proc. roy. Soc. B*, **140**, 455.
 Passey, R. D., and Dmochowski, L. (1950). *British Medical Journal*, **2**, 1129.
 Smith, A. U. (1952). *Exp. Cell Res.*, **3**, 574.
 Strumia, M. M., and Hodge, C. C. (1945). *Ann. Surg.*, **121**, 860.
 Webster, J. P. (1944). *Ibid.*, **120**, 431.
 Willis, R. A. (1948). *Pathology of Tumours*. London.

War injuries are far from being the principal cause of amputations, according to reports submitted to the World Health Organization for a conference on prosthetics held in Copenhagen last month. In the United States, for instance, the second world war produced 18,000 amputees in the armed Forces; during the same period, automobile and industrial accidents, disease, and congenital deformity produced 120,000 civilian amputees. In England, since the war, an average of 3,000 to 4,000 new cases of amputation come forward each year to be fitted with new limbs; in Germany, between 7,000 and 8,000. However, for most countries in the world, the size of the problem is unknown. The aim of the conference was to evolve plans for the organization of prosthetic services in countries with growing industrial and transportation facilities.

G. HADFIELD: THE DORMANT CANCER CELL

FIG. 1.—Cell of a benzpyrene-induced mouse sarcoma which survived many months at -70°C . in the paramorphic state. As seen by phase contrast soon after thawing.

FIG. 2.—Appearance by phase contrast of cells of same tumour one hour after thawing. Cells lie at edge of an air bell (B), where a copious oxygen supply has induced rapid differentiation.

FIG. 3.—Section of tumour shown in Fig. A (Text). Tumour is a cellular spindle-cell sarcoma. ($\times 280$.)

H. B. FELL, E. MELLANBY, AND S. R. PELC: EXCESS VITAMIN A ON CHICK ECTODERM

FIG. 1a.—Control culture; 11 days in normal medium. Unstained section, phase contrast. Note cyst of squamous, keratinizing epithelium.

FIG. 1b.—Same section as Fig. 1a by ordinary lighting to show overlying autoradiograph. The blackening of the autoradiograph, due to $^{35}\text{SO}_4$, is considerable over the connective tissue but very scanty over the squamous epithelium.

FIG. 2a.—+A culture; 11 days in medium containing 1,500 i.u. vitamin A per 100 ml. Unstained section, phase contrast. Note cyst of mucous epithelium.

FIG. 2b.—Same section as Fig. 2a by ordinary lighting to show overlying autoradiograph. Note intense blackening over mucus-secreting epithelium due to $^{35}\text{SO}_4$.