INFLUENCE OF EXCESS VITAMIN A ON THE SULPHATE METABOLISM OF CHICK ECTODERM GROWN IN VITRO

BY

H. B. FELL, D.Sc., F.R.S.*Strangeways Research Laboratory, Cambridge***Sir EDWARD MELLANBY, G.B.E., K.C.B., F.R.S.***National Institute for Medical Research, London*

AND

S. R. PELC, D.Phil.*Medical Research Council, Experimental Radiopathology
Unit, Hammersmith Hospital, London*

[WITH SPECIAL PLATE]

In previous experiments (Fell and Mellanby, 1953) it was shown that when the presumptive epidermis from 6–7-day chick embryos is grown in medium containing excess vitamin A (+A medium) it differentiates, not into the squamous, keratinizing epithelium of the skin as in normal medium but into a mucus-secreting, often ciliated, membrane resembling that of the nose. Recently an attempt has been made to investigate some of the biochemical changes which accompany this metaplasia, by studying the uptake of radioactive sulphur given in the form of the sulphate ($\text{Na}_2^{35}\text{SO}_4$) in ectoderm grown in normal and in +A medium.

Singher and Marinelli (1945) found that when labelled sulphate was injected into rats it was incorporated into chondroitin sulphate and related compounds; its incorporation into amino-acids was negligible in animal though considerable in plant tissues (Howard and Pelc, 1951). Dziewiatkowski *et al.* (1949) confirmed these findings in cartilage. Boström *et al.* (1952) demonstrated the uptake of sulphate in cartilage and also in mucous epithelium by means of autoradiographs, though the resolution was not sufficient to allow detailed investigation of the distribution and movement of the sulphate within the tissues. Using a different autoradiographic technique, Glücksmann and Pelc (1954) showed that in the tracheal cartilage sulphate is taken up by the cells and then diffuses into the matrix.

In unpublished experiments, Glücksmann *et al.* (1954) have found that sulphate is taken up readily by mucus-secreting cells, but only to a small extent by squamous, keratinizing epithelium. The object of the present investigation was to see whether cultures of ectoderm in normal and in +A medium, respectively, would show a similar difference in the uptake of sulphate corresponding with their histological development in the two media.

Present Investigation

The method was as follows. Fragments of ectoderm and underlying connective tissue were removed from the trunk and limbs of 7-day chick embryos and cultivated by the hanging-drop method (see Fell and Mellanby, 1953). Explants taken from one side of each chick were grown in normal medium (3 parts plasma and 1 part embryo extract) and those from the opposite side in the same medium to which had been added 1,500 I.U. of vitamin A alcohol per 100 ml. The tissue was transplanted every 48 hours. The results confirmed those previously described. In both the control and the +A series the ectoderm usually formed a closed cyst surrounded by connective tissue, but after seven

to nine days' cultivation the epithelium in normal medium had assumed a squamous, keratinizing structure, while that in +A medium had formed a mucus-secreting membrane which sometimes was ciliated.

After seven days (two pairs of explants) or ten days (12 pairs) the coverslips bearing the cultures were detached from the hollow ground slides, and a drop of Tyrode's solution containing 200 μC . per ml. of $^{35}\text{SO}_4$ was deposited on each explant with a fine pipette. The drop was allowed to stand for about 30 seconds, after which it was sucked off, leaving only a thin layer of fluid over the tissue; the coverslip was replaced and re-sealed, and the preparation was returned to the incubator for about twenty-four hours. The cultures were then fixed for fifteen minutes in acetic alcohol, followed by thirty minutes in formol-saline; they were transferred from the fixative to 70% alcohol and the tissue was removed from the coverslip with a Bard-Parker knife. After being embedded in paraffin wax the cultures were serially sectioned. Autoradiographs were made by the stripping film technique (Doniach and Pelc, 1950).

There was a striking difference between the autoradiographs of the controls and those of the +A cultures. In the former (Plate, Figs. 1a and 1b), although the connective tissue and basement membrane were labelled, there was very little deposit over the epithelium and none over the keratin. On the other hand, in the +A explants (Figs. 2a and 2b) the connective tissue was labelled as before, but the basal cells of the epithelium had a higher content of sulphate than in the controls, and there was a dense autoradiograph over the inner secretory cells.

Summary

These results show that the sulphur metabolism of the presumptive epidermis of the chick converted into mucus-secreting epithelium *in vitro* by the presence of excess vitamin A in the culture medium is profoundly modified. In the controls, little sulphate is taken up by the squamous epithelium, which thus behaves like normal epidermis *in vivo*, whereas in the +A explants the utilization of sulphate is comparable to that of a normal mucous membrane in the body.

REFERENCES

- Boström, H., Odeblad, E., and Friberg, U. (1952). *Arch. Biochem. Biophys.*, **38**, 283.
Doniach, I., and Pelc, S. R. (1950). *Brit. J. Radiol.*, **23**, 184.
Dziewiatkowski, D. D., Benesch, R. E., and Benesch, R. (1949). *J. biol. Chem.*, **178**, 931.
Fell, H. B., and Mellanby, E. (1953). *J. Physiol., Lond.*, **119**, 470.
Glücksmann, A., Howard, A., and Pelc, S. R. (1954). In preparation.
— and Pelc, S. R. (1954). In preparation.
Howard, A., and Pelc, S. R. (1951). *Nature, Lond.*, **167**, 599.
Singher, H. O., and Marinelli, L. (1945). *Science*, **101**, 414.

The Report of the Director of Water Pollution Research (*Water Pollution Research 1953*, H.M.S.O., price 2s. 6d. net) contains an account of an investigation, requested by the Ministry of Housing and Local Government, to ascertain the extent to which radioactive isotopes in water would be removed when the water is treated at waterworks. Radioactive isotopes might contaminate the water supply as a result of enemy action or of discharge of effluents from industry, hospitals, or research establishments. Most of the work was done with radio-strontium, since this, according to the Report, is one of the most dangerous of the possible contaminants. Treatment with coagulants, at any rate in the concentrations normally used in waterworks, was ineffective, not removing more than 5% of the initial activity. Slow sand filters proved efficient at first, removing nearly all the activity, but with continued use they let through an increasing amount of radioactive material until by the fourteenth day their protective effect was negligible. However, softening of the water was much more hopeful. When calcium hydroxide and calcium carbonate were used, up to 75% of the activity was removed; lime, when the water was completely softened, removed 80%; and base-exchange softeners removed all but about 3%.

G. HADFIELD: THE DORMANT CANCER CELL

FIG. 1.—Cell of a benzpyrene-induced mouse sarcoma which survived many months at -70°C . in the paramorphic state. As seen by phase contrast soon after thawing.

FIG. 2.—Appearance by phase contrast of cells of same tumour one hour after thawing. Cells lie at edge of an air bell (B), where a copious oxygen supply has induced rapid differentiation.

FIG. 3.—Section of tumour shown in Fig. A (Text). Tumour is a cellular spindle-cell sarcoma. ($\times 280$.)

H. B. FELL, E. MELLANBY, AND S. R. PELC: EXCESS VITAMIN A ON CHICK ECTODERM

FIG. 1a.—Control culture; 11 days in normal medium. Unstained section, phase contrast. Note cyst of squamous, keratinizing epithelium.

FIG. 1b.—Same section as Fig. 1a by ordinary lighting to show overlying autoradiograph. The blackening of the autoradiograph, due to $^{35}\text{SO}_4$, is considerable over the connective tissue but very scanty over the squamous epithelium.

FIG. 2a.—+A culture; 11 days in medium containing 1,500 i.u. vitamin A per 100 ml. Unstained section, phase contrast. Note cyst of mucous epithelium.

FIG. 2b.—Same section as Fig. 2a by ordinary lighting to show overlying autoradiograph. Note intense blackening over mucus-secreting epithelium due to $^{35}\text{SO}_4$.