In three cases treatment was discontinued owing to confusion and to deterioration of the epilepsy.

I wish to thank Dr. J. M. Crawford, physician-superintendent of Botleys Park Hospital, for his help and encouragement with the above report; Dr. J. Carse, physician-superintendent of Graylingwell Hospital, Chichester; Dr. Parsons-Smith and the staff of the E.E.G. department there for carrying out the necessary E.E.G. investigations; and Miss F. K. FitzPatrick and the nursing staff of Botleys Park Hospital for their invaluable help. I acknowledge with gratitude the financial support given by the Research Subcommittee of the South-west Metropolitan Regional Hospital Board, without which the survey could not have been undertaken.

REFERENCES

- Butter, A. J. M. (1953). *Lancet*, 1, 1024.
Greenstein, L., and Sapirstein, M. R. (1953). *Arch. Neurol. Psychiat.*, Chicago, 70, 469.
Handley, R., and Stewart, A. S. R. (1952). *Lancet*, 1, 742.
Tredgold, A. F. (1952). *A Textbook of Mental Deficiency*, 8th ed. London.
Whitty, C. W. M. (1953). *British Medical Journal*, 2, 540.

USE OF POLYMYXIN B IN THE EXTERNAL EAR

BY

D. A. T. FARRAR, M.B., F.R.C.S.

E.N.T. Registrar, St. Bartholomew's Hospital, London

The polymyxin antibiotics were first described by Benedict and Langlykke (1947) and by Ainsworth *et al.* (1947). A series of antibiotic agents derived from *Bacillus polymyxa* have now been isolated and described; they are called polymyxins, and are differentiated by letters of the alphabet. The action of the polymyxins so far described—A, B, C, D, and E—seems to be bactericidal, and they all have similar antibacterial spectra, without significant differences (Brownlee *et al.*, 1952). *In vitro* experiments (Brownlee *et al.*, 1949; Pulaski *et al.*, 1949) have shown that they are effective against most Gram-negative organisms, in low concentrations. *Proteus vulgaris*, however, is an exception, and, together with the Gram-positive organisms, is sensitive only to high concentrations.

The polymyxins, chiefly B and E (the least toxic), have been used clinically, mainly by parenteral routes, in various infections due to Gram-negative organisms; the chief drawback to their use is their toxicity. A few workers have reported the use of these substances locally with some success (Jackson *et al.*, 1951; Jawetz, 1952). There do not appear to be any reports of a series of aural infections treated locally. There have also been no reports of any sensitization reactions after local use.

Present Series

Infections of the external auditory meatus, and in this group are included infected mastoid cavities, are often due to Gram-negative organisms. These infections are troublesome, since the organisms, particularly *Pseudomonas pyocyanea*, are often resistant to the commonly used antibiotics. A series of cases of otitis externa and infected radical mastoidectomy cavities have therefore been treated with polymyxin B locally. The cases chosen were those in which the infecting organisms were found to be one of the three chief Gram-negative types found in this region, *Ps. pyocyanea*, *Pr. vulgaris*, and *Bact. coli*. Infections caused by mixed organisms were not included.

The preparation used was a solution of 10% polymyxin B sulphate in propylene glycol, acidified with acetic acid. It was supplied to the patients in a bottle, with a dropper, for use at home. They were instructed to fill the ear with the solution, placing the head on one side, with the affected ear

uppermost, morning and evening, and then to insert a plug of cotton-wool. A swab was taken from the ear for laboratory examination before treatment was begun and another after seven days' treatment. In most cases in which organisms were sensitive the ear was found to be dry at the end of this period. The results are shown in the Table. In a

Results

Organisms	Type	No. of Cases	Dry	Failed	% Dry
<i>Pr. vulgaris</i> .. {	Otitis externa	8 (2c)	1	7	10
	Radical cavity	12	1	11	
<i>Bact. coli</i> .. {	Otitis externa	7 (3c)	5	2	65
	Radical cavity	13	8	5	
<i>Ps. pyocyanea</i> .. {	Otitis externa	11 (3c)	10	1	90
	Radical cavity	9	8	1	

c = Cases with bilateral infection in which a control solution was used in one ear.

small number of cases the infection was bilateral, and the infecting organisms were the same on both sides. In these cases the polymyxin solution was instilled into one ear and a control solution, consisting of solvent only, was instilled into the other. At the end of the seven-days period the control solution was stopped, a swab taken for culture, and treatment with the polymyxin solution begun. In no case did the control solution alter the infecting organisms. The number of cases suitable to act as controls is small. No sensitivity or reaction to the solution was noted.

It would seem that the polymyxin B solution is effective in a large proportion of cases against infections of the external ear caused by *Ps. pyocyanea*. To a less extent it may clear up infections caused by *Bact. coli*. It is ineffective against *Pr. vulgaris*. These results conform to the *in vitro* sensitivity of the organisms concerned, except that *Bact. coli* appears to be less effectively controlled in the ear than one might expect from its *in vitro* sensitivity. Since *Ps. pyocyanea* infections are often resistant to other antibiotics and other forms of treatment, it is suggested that polymyxin B sulphate solution has a place in their local treatment.

Summary

A series of cases are reported in which polymyxin B sulphate was used in the treatment of otitis externa and infected mastoid cavities.

Polymyxin B sulphate appears to be effective against infections caused by *Pseudomonas pyocyanea*.

I would like to thank Mr. F. C. W. Capps for his encouragement and help, and Mr. F. Prescott and the Wellcome Research Institute for their generous help and supplies of polymyxin B sulphate.

REFERENCES

- Ainsworth, G. C., Brown, A. M., and Brownlee, G. (1947). *Nature, Lond.*, 160, 263.
Benedict, R. G., and Langlykke, A. F. (1947). *J. Bact.*, 54, 24.
Brownlee, G., Bushby, S. R. M., and Short, E. I. (1949). *Ann. N.Y. Acad. Sci.*, 51, 952.
— (1952). *Brit. J. Pharmacol.*, 7, 170.
Jackson, D. M., Lowbury, E. J. L., and Topley, E. (1951). *Lancet*, 2, 137.
Jawetz, E. (1952). *Arch. intern. Med.*, 89, 90.
Pulaski, E. J., Baker, H. J., Rosenberg, M. L., and Connell, J. F. (1949).

The 1954 *Yearbook* of the British Standards Institution, recently published (12s. 6d.), lists the British Standards current at March 31, 1954, and gives a brief description of each. They cover every field of industry and technology, and the standards laid down may be concerned with design, physical properties, methods of testing, limit tolerances, and so on. Many articles commonly used in hospitals and laboratories are manufactured in accordance with these standards, which are agreed only after discussion with all the parties concerned. The British Medical Association is a member organization and appoints representatives to committees of the B.S.I. when a medical opinion is required. Individuals are also eligible for membership of the B.S.I., and particulars as well as the *Yearbook* may be obtained from it at 2, Park Street, London, W.1.