We are indebted to the Medical Research Council, the Colonial Office, and the Colonial Medical Research Committee, and in particular to Dr. R. Lewthwaite, Director of Colonial Research, for sponsoring these investigations. Our thanks are due to Dr. S. L. A. Manuwa, Inspector-General of Medical Services, Nigeria; to Colonel H. W. Mulligan, Director of the West African Institute for Trypanosomiasis Research; to Dr. A. S. Garrett, Medical Superintendent of the Oji River Leper Settlement; to Dr. Davenport, of the Seventh Day Adventist Mission, Jengre; to Dr. F. J. Cauchi and Dr. E. A. Fejer, of the Colonial Medical Service; to Mr. W. Crewe and Mr. P. J. Moore, of the Helminthiasis Research Scheme, Kumba; and to Mr. F. Henry, of the Mobile Field Unit, North Nigeria, for their co-operation. Our thanks are also due to the United Africa Company for a grant for research in filariasis.

REFERENCES

- Blacklock, D. B. (1926). *Ann. trop. Med. Parasit.*, 20, 1.
Buckley, J. J. C. (1949). *J. Helminth.*, 23, 1.
Mazzotti, L. (1951). *Amer. J. trop. Med.*, 31, 625.

INTRAMUSCULAR BENZATHENE PENICILLIN IN THE PROPHYLAXIS OF STREPTOCOCCAL INFECTION IN RHEUMATIC CHILDREN

BY

C. B. PERRY, M.D., F.R.C.P.
Professor of Medicine

AND

W. A. GILLESPIE, M.D., M.R.C.P.I., D.P.H.
Pathologist

University of Bristol and United Bristol Hospitals

In 1951 Szabo, Edwards, and Bruce described the preparation and chemical properties of a new penicillin compound—N,N'-dibenzylethylenediamine penicillin G (benzathene penicillin). Elias, Price, and Merriam (1951) showed that this substance marked a valuable advance in depot penicillin therapy, and that useful blood levels were maintained for as long as 144 hours after a single dose of 300,000 units, and for 12 days after 2.5 mega units. O'Brien and Smith (1952) reported that two injections at weekly intervals, or even a single injection of 2.3–2.5 mega units, gave excellent results in the treatment of gonorrhoea, 97% of the cases so treated being apparently cured. Further, fourteen days after the single dose appreciable blood levels of penicillin were found. Such a preparation clearly promised to be of value in the prophylaxis of streptococcal infection in patients who had suffered from rheumatic fever, and Stollerman and Rusoff (1952) reported the results of its use in 135 children. They found that 300,000 units maintained adequate blood levels for at least a week, 600,000 units for two weeks, and 1.2 mega units in most cases for four weeks. No patient developed a sore throat during treatment, but there were instances of temporary "bacteriological relapse" but with no evidence of clinical illness. These claims are so important that it appeared desirable to confirm them.

For this purpose a study has been made of children recovering from acute rheumatism or chorea in a hospital school.

Methods

Throat swabs were plated, within three hours of taking, on 5% horse-blood digest agar containing 1:700,000 crystal violet and incubated overnight in an anaerobic jar at 37° C. One colony from every plate with β -haemolytic streptococci was subcultured on blood agar and was grouped by the method of Maxted (1948). All group A cultures were tested for penicillin sensitivity by a ditch-plate method and sent for typing to the Streptococcal Reference Laboratory, Colin-

dale. Serum penicillin levels were measured by the *Sarcina lutea* plate method (Welch, 1948) modified by diluting the suspension of the organism so that its final concentration in the plates was 1 in 2,000 of the original culture.

The study lasted from March 1, 1953, to February 28, 1954. During this period 53 rheumatic fever convalescent children were admitted and 48 were discharged. Their ages were from 6 to 16 years. The duration of their stay in hospital ranged from 83 to 316 days. The children were accommodated in two wards, where there were also some children with orthopaedic complaints. The number of rheumatic children in the wards at a time varied from 10 to 23, and the number of orthopaedic patients varied from 33 to 35 at any one time. Except for benzathene penicillin, no specific treatment was given during the study. Throat swabs were taken at weekly intervals from all the rheumatic patients, and from time to time from the orthopaedic patients. All group A streptococci were typed. From the results of the swabs, estimates were made, as described below, of the load of infection in the community—that is, the numbers who were carrying group A streptococci—and of the numbers of new infections which took place among the rheumatic children. Haemolytic streptococci of groups other than A were isolated occasionally, but were not considered further.

Twenty-two rheumatic children were given the specific treatment which was used for the first time on July 1. At first cases were chosen because they were persistent carriers of group A streptococci; later alternate admissions were treated. Fifteen of the treated children had been persistent carriers for from four to nine weeks before their first injection. The other seven had negative swabs for from two to seventeen weeks before the first injection. The treated children were distributed at random in the two wards. No child was treated until it had been in hospital for at least three weeks. Benzathene penicillin was given intramuscularly, into the buttock, in doses of 1.5 mega units contained in 5 ml. One child received five doses, seven children received four doses, six received three, five received two, and three received only one dose.

Serum penicillin levels were measured 4, 11, 18, and 25 days after the injection. The results are summarized in Table I. Most of the assays at 25 days showed less than

TABLE I.—Serum Penicillin Concentrations

	Days after Injection			
	4	11	18	25
Total No. of samples	66	62	58	56
No. with serum concentrations of				
0.02 u./ml. and over	66	54	33	15
Range (u./ml.)	0.24–0.025	0.14–0	0.10–0	0.07–0
Av. serum concentration (u./ml.)	0.08	0.043	0.029	0.012

0.02 unit per ml., but in many of these there were still detectable traces of penicillin in the serum. Absorption appeared to be irregular, even in the same child. Thus, after some injections serum levels remained above 0.03 unit per ml. for at least 25 days, while after others no penicillin was detected after shorter periods. The Chart shows the levels found in four children and how they varied in the same child from injection to injection. As noted by Stollerman and Rusoff, most of the injections gave rise to local soreness and tenderness for 24 to 48 hours. No urticarial or other skin rashes were seen. However, in 18 out of 67 injections slight fever was noted on the following day. This ranged from 99 to 100.5° F. (37.2 to 38.1° C.). It did not occur after every injection in the same child.

The "control" group of rheumatic children was made up of 43 children who received no treatment and the 22 "treated" children who were observed for from three to seventeen weeks before starting treatment. Seventeen of these were group A carriers on admission and yielded persistently positive swabs, without a change of type, throughout their period of observation, for from four to twenty weeks. Twenty-two had persistently negative swabs through-

out their periods of observation, which varied from four to 21 weeks. The remainder in the control group gave one or more positive swabs intermittently, and often with a

Curves showing the blood levels of penicillin (units per ml.) in four different children at weekly intervals after 1.5 mega units of benzathene penicillin. Arrows indicate the times of injections. Detectable penicillin of less than the minimum assayable value is shown as 0.01 unit per ml.

change of type, presumably as a result of cross-infection. Only one of these infections was accompanied by clinical signs or symptoms—that is, complaint of sore throat with three days' pyrexia.

Effect of Treatment

Group A streptococci were eliminated within four days from the throats of all but one of the 15 persistent carriers in the treated group. In the one exception, group A streptococci of the same type as he started with were isolated from the patient at the second and third swab after his first injection, and then disappeared. As a result of this effect of treatment the load of infection, which had been very high at the beginning of the study, fell after July 1 (Table II).

TABLE II.—Numbers of Swabs which Yielded Group A Streptococci

	Rheumatic Patients				Orthopaedic Patients Untreated	
	Treated		Untreated*		Total	Positive
	Total	Positive	Total	Positive		
March			87	35		
April			99	30		
May			131	69	28	8
June			87	44		
July	43	1	75	37		
August	59	3	42	10		
September ..	51	0	39	5	26	10
October	51	0	53	7	31	6
November ..	27	0	51	15		
December ..	12	0	45	8		
January ..	7	2	49	11	20	7
February ..	9	0	45	10	19	5

* Includes patients subsequently treated.

Hence, in order to compare the rate of occurrence of infections among treated and control patients it was necessary to consider separately the initial period, before any treatment was given, and the period July 1 to February 28. A new infection was shown by the isolation of a group A streptococcus from a throat which was previously negative, or which had previously carried a different type. However, as it was possible that a few of the patients carrying group A streptococci had more than one type in their throats at the same time, and as only one colony was picked for testing from each positive plate, there had to be a gap of at least one negative swab before accepting the isolation of a new type as evidence of infection in a patient who had previously carried Group A streptococci.

In estimating the infection rates, differences in times of exposure to the infective environment were allowed for by

relating the number of infections (as defined above) to the total number of throat-weeks at risk. When a patient yielded a positive swab his throat was not regarded as being again at risk unless it became negative for at least one swab. Thus, for example, if three patients with initially negative swabs had been observed for a period of six weeks, and if two of them had yielded positive swabs at the second and fourth weekly swabbing respectively and had stayed positive subsequently, there would have been two infections in 12 throat-weeks, giving an infection rate of 16.7%. The average infection rate in all the rheumatic children observed between March 1 and June 30, 1953 (before any treatment was given), was 6.6%. Between July 1, 1953, and February 28, 1954, the rates in the control and treated groups were respectively 2.9% and 0.37% (Table III). The fall in the infection rate among the control patients after July 1 was presumably caused by the lessening of the load of infection in the whole community.

Throughout the study the penicillin sensitivity of every culture of group A streptococcus was tested by a filter-paper strip method. In no case was resistance found to be greater than that of the Oxford staphylococcus.

TABLE III.—Group A Streptococcus Infection Rates in Treated and Untreated Rheumatic Fever Convalescent Children

	March to June, 1953			July, 1953, to Feb., 1954		
	Total Swabs	Positive Swabs	%	Total Swabs	Positive Swabs	%
Load of infection in community (including orthopaedic patients)	532	186	35.0	754	137	18.2
	Infection Rate					
	Throat-weeks at Risk	No. of Infections	Rate %	Throat-weeks at Risk	No. of Infections	Rate %
Untreated patients	196	13	6.6	312	9	2.9
Treated ..	0	—	—	270	1	0.37

Conclusions

Monthly intramuscular injections of benzathene penicillin in doses of 1.5 mega units produced prophylactic blood levels of penicillin in the majority of children.

These injections gave rise to some local tenderness for 24 hours, and in some cases slight fever.

These injections eliminated group A haemolytic streptococci from the throats of carriers, and almost completely prevented new infections.

We are indebted to Mr. L. J. Waller for much technical assistance and to Dr. R. E. O. Williams for typing the streptococci. Our thanks are also due to John Wyeth and Brother Ltd. for a generous supply of benzathene penicillin ("penidural"), without which the investigation would not have been possible.

ADDENDUM.—Since this study was completed one child developed a group A streptococcal sore throat 25 days after an injection of benzathene penicillin and subsequently developed a relapse of rheumatism.

REFERENCES

- Elias, W., Price, A. H., and Merion, H. J. (1951). *Antibiot. and Chemother.*, 1, 491.
- Moxed, W. R. (1948). *Lancet*, 2, 255.
- O'Brien, J. F., and Smith, C. A. (1952). *Amer. J. Syph.*, 36, 519.
- Stollerman, G. H., and Rusoff, J. H. (1952). *J. Amer. med. Ass.*, 150, 1571.
- Szabo, J. L., Edwards, C. D., and Bruce, W. F. (1951). *Antibiot. and Chemother.*, 1, 499.
- Welch, H. (1948). *Methods in Medical Research*, 1, 25. Chicago.