HOUSING AND ILLNESS

AN ANALYSIS OF 1,000 MEDICAL CERTIFICATES

BY

M. D. WARREN, M.D., D.P.H., D.I.H.

Deputy Medical Officer of Health, Hampstead;
Assistant Principal Medical Officer, L.C.C.

At the end of 1952 the Metropolitan Borough Council of Hampstead decided to review all its outstanding housing applications. There were 3,867 such applications at that time, of which 3,405 had the necessary residential qualifications. Of these 3,405, 1,249 were considered to have very little or no housing need. Of the remaining 2,156 applications, 706 were supported by one or more medical certificates—that is, 33% of all genuine applications had supporting medical certificates. In addition to these 706, which form the bulk of this analysis, all housing applications supported by medical certificates submitted between April 1, 1953, and March 31, 1954, are included. These are shown separately in the Table, as it was felt that there might be some significant differences between current applications and those on the waiting-list.

The total of 1,028 applications were supported by medical certificates relating to 1,240 persons, made up of 685 females (55%), 291 males (24%), and 264 children (21%). It is not surprising to find that the housewives are the most affected by bad housing, but it is suggested that some of this excess could be regarded as being on a par with "industrial" illness and neurosis. In the same way that adverse environmental conditions in factories and workshops have been shown to affect the health of the employees, so bad working conditions in the home might affect the housewife.

Reasons for Certification

In 861 applications (84%) only one member of the family was certified as suffering from an illness related to their housing conditions; in 131 applications (13%) two members were certified, in 28 applications three members, in seven applications four members, and in one application five members of a family were considered to be affected.

With regard to the number of diseases each person was certified to be suffering from, 1032 persons (83%) were suffering from one illness, 175 persons (14%) from two illnesses, 32 persons from three illnesses, and one person from four illnesses. In calculating these figures an effort has been made to count only different illnesses and not different symptoms arising from one illness. For instance, an old person might be certified as suffering from "bronchitis, hypertension, and arthritis," and this has been counted as three illnesses. On the other hand, a person certified as suffering from "asthma, hay-fever, and bronchitis" is recorded as a single illness—asthma.

A broad grouping of the diagnoses of the total 1,482 "illnesses" is shown in the Table. Twenty-five per cent. of all certificates related to upper or lower respiratory infections excluding tuberculosis, which accounts for 7.5%. The next commonest cause is the group comprising arthritis, fibrositis, and rheumatism, accounting for 15%; while psychological conditions (anxiety states, neurasthenia, depression, etc.) come third at 14% of the

total. Heart disease accounts for 11%, but most of these certificates are requests for ground-floor accommodation.

The differences in illness between the adult males and females are quite marked. The females show an excess of certificates relating to "rheumatic" disorders (20%) and psychological conditions (19%), whilst the males have an excess in the bronchitis-emphysema group (17%) and in

Illness Involved

	Females		Males		Children		Total	%
	Review	1953-4	Review	1953-4	Review	1953-4		
Bronchitis, bronchiectasis, emphysema ..	53	27	42	17	67	20	226	15
Asthma ..	20	9	12	4	25	9	79	5
Tonsillitis, catarrh, colds ..	9	7	3	2	39	18	78	5
Tuberculosis ..	29	10	38	16	8	10	111	7.5
Arthritis, rheumatism, fibrositis ..	119	48	26	12	12	7	224	15
Psychological ..	105	56	22	3	13	9	208	14
V.D.H., unspecified cardiac failure ..	46	17	14	4	1	—	82	11
Coronary disease ..	8	3	10	11	—	—	32	
Hypertension ..	25	15	6	2	—	—	48	
Peptic ulcer ..	15	1	23	9	—	—	48	3
Rheumatic fever ..	3	—	1	—	7	—	11	
Varicose veins ..	18	4	—	2	—	—	24	
Pregnancy ..	22	7	—	—	—	—	29	
Gynaecological ..	9	2	—	—	—	—	11	18
Other ..	119	51	36	23	31	11	271	
Total ..	600	257	233	105	203	84	1,482	

the tuberculosis group (16%). The commonest complaints certified in children were bronchitis (30%), tonsillitis, catarrh, etc. (20%), asthma (12%), and tuberculosis (6%).

There is little difference between the distribution of causes from the two sources of certificates, the only marked difference being between tuberculosis in children, which accounts for 12% of current certificates and 4% of the waiting-list. If it is assumed that there has been little difference in incidence in the last three or four years, then this difference could mean that these cases are tending to be rehoused before other cases.

The Question of Priority

There is no doubt that any badly housed family will be given a better opportunity to be healthy in adequate and satisfactory accommodation, and therefore all families whose health is considered by their doctor to be affected adversely by their living accommodation must be regarded as genuinely in need of rehousing. However, if all such patients were given equal priority on medical grounds, the whole object of awarding priority would be defeated. This difficulty is underlined by the following figures. During 1953, Hampstead Council found accommodation for 143 applicants, involving 443 persons. The number of families on the waiting-list at December 31, 1953, was 4,187, of whom about 1,000 had supporting medical certificates. Owing to the shortage of housing sites, the Council does not believe that it will be able to rehouse more than 900 families in Hampstead during the next five years. Thus the problem presented to the medical officer of health is to allocate priority even among the top priority group. Furthermore, it is the duty of the medical officer of health to prevent illness, and therefore it is necessary so to allocate priority that healthy people who are in danger of falling ill because of their accommodation are given an opportunity of being rehoused before any illness develops. It is not enough for a housing department to rehouse the ill.

Respiratory Disease.—The problem of assessing fairly the priority to be awarded to a family with illness is extremely complex. Unfortunately, little help can be obtained from the "literature." Thus the results of an investigation in Glasgow in 1928-9 (Smith, 1934) into housing conditions and respiratory disease were equivocal: a greater morbidity was found on the new estates, but the two groups

utilized were not strictly comparable, as among the rehoused group there was more unemployment. The fact that 25% of all certificates examined here related to respiratory infection must be contrasted with the observation that these diseases account for about 30% of the general practitioners' consultations (Pemberton, 1951; Logan, 1953).

Rheumatic Disorders.—These accounted for 15% of the certificates. Yet Kellgren *et al.* (1953) could find no evidence in their study of rheumatic complaints in an urban population that damp housing conditions greatly affected the incidence of rheumatic complaints. Furthermore, they showed that 33% of the population stated that they had suffered from a rheumatic complaint during the preceding five years. Similarly the Empire Rheumatism Council (1950) found no satisfactory evidence that home conditions are of aetiological importance in the causation of rheumatoid arthritis. Perry and Roberts (1937) found a significant association between overcrowding and the incidence of rheumatic fever, but Savage (1931) considered that housing factors could be, at the most, only "slightly predisposing factors" in the aetiology of rheumatic fever. More recently Hewitt and Stewart (1952) reported that they could find "no evidence that rheumatism is associated either with bad housing conditions or with poverty, *per se*."

Child Mortality.—The findings in regard to child mortality have been more definite. Thus the Registrar-General (1935) found that mortality from all causes at ages under 5 increased with greater overcrowding, the mortality from bronchitis, measles, and whooping-cough showing this association most definitely. Again, the Chief Medical Officer of the Ministry of Health (1938), in his report for 1937, states, "The density figures show that the level of mortality, especially at 1-2 years, is, without doubt, closely associated with the level of the housing index."

Tuberculosis.—Most investigators have also been able to demonstrate a correlation between adverse housing and tuberculous infection. Benjamin (1953), in a study of the differential incidence and mortality from pulmonary tuberculosis in the London Metropolitan Boroughs, confirmed the findings of Hart and Wright (1939) and Stein (1952) that overcrowding adversely affected tuberculosis mortality. The increased incidence of tuberculosis among contacts (especially child contacts) of known cases has also been demonstrated (Blacklock, 1947; Miller, 1947; Aspin, 1952).

Mental Disease.—Not so much research has been carried out on the causative role of bad housing in mental disease. Hadley *et al.* (1944) found the Army rejection rates for personality disorders one and a half times as great in slum areas as in other areas. Hyde and Kingsley (1944) found the total incidence of mental disorders increased significantly from the best to the poorest communities. But in neither paper is there enough evidence to say that it is the bad housing itself, rather than other associated factors, that causes the increase in mental disorders.

Assessment for Rehousing

Thus the difficulties of fairly allocating priority on medical grounds are great. There are three basic ways in which housing conditions may affect health. These are structural defects and insanitary nuisances, overcrowding, and extraneous factors such as stairs and other tenants.

Structural defect and insanitary nuisances can often be remedied by the intervention of the sanitary inspector; alas, many tenants do not wish the inspector to take action, as they are afraid either that they will be evicted by the landlord or that they may lose points in their housing application.

Overcrowding can be alleviated only by rehousing, and in the Hampstead scheme points are given for overcrowding. In addition to these points, more are given for each additional bedroom required. Unfortunately, in assessing overcrowding a baby under 1 year does not count, and a child between 1 year and 10 years counts as half a unit. In view of the established fact that these children are the most

affected by bad housing, it would seem wiser if they counted at least as one whole unit. However, this difficulty can be overcome to some extent by allotting more medical points to a child with an illness than to an adult with the same condition. Points are also allotted for lack of certain facilities and for separated families and families without separate homes. Additional balancing points are awarded for the period on the waiting-list and for service in the Forces (including the Merchant Navy). These additional points are used to choose between two applications which otherwise have the same total points.

The medical officer of health can award up to a maximum of 20 points for certified ill-health or disablement. The guiding principles in awarding these points are that the illness is related to or adversely affected by housing conditions which can be alleviated only by rehousing, and that it is likely that the patient will materially benefit by rehousing. The medical officer of health, therefore, before allocating points, requires a complete report of the housing conditions, including the structural state of the house and the number, sex, and distribution of the people living there. In addition he needs quite a detailed account of the illness.

Chalke (1954) has elaborated all the information necessary before allocating priority to persons with tuberculosis. This includes the activity of the disease, the infectivity of the sputum, the age of all possible contacts, the standard of patient and contacts in respect of precautionary measures, and the standard of present housing, including whether the patient has to share lavatory and bathroom with other families. After this information has been obtained, Chalke divides his tuberculous applicants into five priority groups. Similar detailed information is required about each applicant, the details varying, of course, according to the disease. A frequent request is for a person to be rehoused on the ground floor because of varicose veins. Immediately one wants to know the severity of the condition, whether an ulcer is present, and why operative treatment is not indicated. A certificate which states that Mrs. A is suffering from a "medical condition" is quite useless.

Provided adequate information is available, the points system works reasonably well. Its only defect is that it cannot cope with a person suffering from an illness aggravated by his present accommodation, which is otherwise satisfactory. For instance, a person with a coronary thrombosis living three stories up may be given the maximum medical points, but this will be insufficient for rehousing unless he has other points, which of course he will not have if his accommodation is otherwise suitable and adequate. Similarly with some psychological situations, such as intolerable hostility between tenants or tenant and landlord. In such cases, although rehousing may be clearly indicated on medical grounds, the award of medical points alone will not be sufficient to obtain overriding priority. In these cases an effort must be made to arrange an exchange; but this is usually difficult to carry out, as the occupiers of ground-floor flats are not willing to leave them for upstairs accommodation, and people are not anxious to move into a house where the landlord or other tenants are known to be querulous.

Summary

An analysis of 1,028 applications for rehousing supported by medical certificates showed that 25% related to respiratory infections, 15% to rheumatic disorders, and 14% to psychological disorders. Of all these certificates, 55% related to women, 24% to men, and 21% to children. A brief survey of the previous papers shows that the fact that bad housing causes or aggravates these conditions is not proved, with the exception of tuberculosis and infections in young children. The medical officer of health who has to award "medical points" for rehousing needs to know much detailed information about the housing conditions and the medical condition before he can allocate points. The former information

he obtains from his inspectors, for the latter he is dependent on the reports of the general practitioner or specialist.

I wish to express my thanks to Dr. D. H. Geffen, Medical Officer of Health, Metropolitan Boroughs of Hampstead and St. Pancras, for his help in preparing this paper and for his permission to publish. The views expressed are entirely my own.

REFERENCES

- Aspin, J. (1952). *Lancet*, 1, 502.
 Benjamin, B. (1953). *Brit. J. Tuberc.*, 47, 4.
 Blacklock, J. W. S. (1947). *British Medical Journal*, 1, 707.
 Chalke, A. H. D. (1954). *Publ. Hlth.*, 67, 66.
 Empire Rheumatism Council (1950). *British Medical Journal*, 1, 799.
 Hadley, E. E., et al. (1944). *Psychiatry*, 7, 379.
 Hart, P. D'Arcy, and Wright, G. P. (1939). *Tuberculosis and Social Conditions in England*. N.A.P.T., London.
 Hewitt, D., and Stewart, A. (1952). *Brit. J. soc. Med.*, 6, 161.
 Hyde, R. W., and Kingsley, L. V. (1944). *New Eng. J. Med.*, 231, 543, 571.
 Kellgren, J. H., Lawrence, J. S., and Aitken-Swan, J. (1953). *Ann. rheum. Dis.*, 12, 5.
 Logan, W. P. D. (1953). Studies on Medical and Population Subjects. No. 7. "General Practitioners' Records" 1951-2. H.M.S.O., London.
 Miller, F. J. W. (1947). *British Medical Journal*, 2, 91.
 Ministry of Health (1938). Annual Report of Chief Medical Officer, 1937. H.M.S.O., London.
 Pemberton, J. (1951). *Lancet*, 1, 224.
 Perry, C. B., and Roberts, J. A. F. (1937). *British Medical Journal*, 2, Suppl. 154.
 Registrar-General (1935). Annual Statistical Review for the Year 1932. Text. H.M.S.O., London.
 Savage, W. G. (1931). *British Medical Journal*, 2, Suppl. 37.
 Smith, C. M. (1934). *Spec. Rep. Ser. med. Res. Coun. Lond.*, No. 192. H.M.S.O., London.
 Stein, L. (1952). *Brit. J. soc. Med.*, 6, 1.

BRITISH DOCTORS IN RUSSIA

EXPERIENCES AND IMPRESSIONS

A party of 17 British doctors and two lay persons connected with the Health Service* recently returned from a three-weeks visit to Russia. The visit was undertaken under the auspices of the Medical Section of the Society for Cultural Relations with the U.S.S.R., and on September 30 six members of the party attended a press conference at the headquarters of the Society and gave some account of their tour. They were Lord Amulree, Dr. Alex Comfort, Dr. L. Crome, Mr. A. W. L. Kessel, Professor Esther Killick, and Dr. N. B. Malleison.

The tourists had spent a week in Moscow, where they met officials of the Ministry of Health, and then had divided into sections, visiting various parts of the Soviet Union, including Leningrad on the Gulf of Finland, and the Republic of Georgia on the shores of the Black Sea. As many contacts as possible were made, from the medical institutes in Moscow to the rural polyclinics. One member of the party, Dr. J. G. Ollerenshaw, a general practitioner of Skipton, spent two days with an "opposite number" in a village about 60 miles from Moscow and also two days on a collective farm in Georgia.

The Status of the Russian Doctor

The party were plied with questions from the press conference on subjects ranging from the professional and economic status of the doctor in Russia, their distribution, and the scope of general practice, to the reliability of public health data and the practice of psychiatry. Much of the information brought back by the party was in a line with the facts given in the paper by Kurashov, the Deputy Minister of Health, printed in the Educational Number of the *British Medical Journal* (August 28, p. 510). The medical service in Russia is free to all, but, except when supplied to hospital in-patients, medicines are charged for at a price proportional to cost. The person who cannot afford to pay is supplied free, and preparations such as insulin are also free. A small amount of private practice still exists, mostly among the more senior members of the profession, and there is no legal restriction on the acceptance

of a fee. But private practice is said to be of very small account.

The medical service is entirely salaried. The new doctor on qualification receives a salary of 670 roubles a month, about the same as any skilled worker. An unskilled labourer earns about 400 roubles a month, a foreman in industry 800, and a miner 2,000. The salary for the ordinary doctor goes up to 1,200 roubles or more, but with the attainment of higher qualifications it rises steeply. A professor may receive 5,000 roubles, and the head of an institute—the institute being a medical faculty separated from the university and under the Ministry of Health—receives 6,000, plus half the salary of a professor, making 8,500 roubles. The great difference between the salary of the clinician and what is termed the "medical scientist" struck the visitors most forcibly.

At age 55, or after 25 years' service, the doctor is eligible for pension amounting to half salary, but apparently he may continue working after reaching pensionable age. Three years after qualification he can claim a subsidized refresher course, and he may do so again after further years for as long as he continues in the service.

Discipline and Appointment

Asked about the disciplining of doctors, the members of the party said that they had not elicited much information. In each district there is a commission, medical in membership, presided over by the head of the local medical organization, which is responsible not only for disciplinary action but for removing misfits and resolving disputes. The doctor's right of appeal against a decision to remove him on the ground of misconduct was not exactly ascertained, but it was understood that he had the right of complaint to his trade union or local party, and that civil action could be taken on his behalf. Vacancies in the service are filled by election by the medical commission after consultation with the various bodies concerned. Each post is re-advertised at five-year intervals, but the incumbent is eligible for re-election if he wishes. When a student qualifies he is invited into a board-room, where he may face some twenty people accustomed to looking over the "new boy." These are representatives of organizations wanting doctors; they state their needs, and with the help of the candidate's teachers an effort is made to place him where he will be most useful and contented.

For three years the young doctor must go where he is sent; but after that he can choose his place of practice. During his last year as an undergraduate he may be taken with others to some centre where he is able to see the service in operation. Often groups of five or six work together, and thus afford a means of bringing medical services into more remote districts. Great prestige attaches to willingness to work in uncongenial places, and this is reinforced by laudatory mention in the papers and the award of medals. In reply to a question Lord Amulree said that he believed some financial bonus also attached to practice in the more remote regions. One Moscow medical school "patronizes" a region 200 miles away, and there is a constant interchange of personnel—doctors, auxiliaries, and students.

Surplus of Doctors

A strong impression was given that, in some areas at any rate, there is a surplus of doctors. At the polyclinics a two-shift system is worked, and the doctor's hours are five or six daily, including the time spent on visiting. One member of the party suspected that the doctors might suffer from lack of clinical material. Another spoke of visiting a medium-sized rayon factory, where they found several doctors at the polyclinic, a separate clinic for young children, another for older children, and yet another doctor, part-time, attached to the athletic activities of the factory. A country hospital which was visited seemed to have a larger staff than could be fully employed. There was said to be, on an average, one doctor to every 660 people, but in some places the ratio was even smaller.

*Their names were given in our news column on September 11 (p. 655).