considered unnecessary. The wound was then closed in layers, positive intrathoracic pressure being induced by the anaesthetist before final closure of the thorax.

Apart from transitory paralytic ileus and a pleural effusion which required aspiration, the patient made an excellent recovery and returned to work in the middle of July.

DISCUSSION

The apparent increase in cases of diaphragmatic hernia, both traumatic and otherwise, in recent years is due partly to better methods of diagnosis and partly to the realization that apparently vague digestive and respiratory symptoms may be due to this condition.

The variety of diseases with which diaphragmatic hernia may be confused led Harrington (1950) to regard this condition as "the masquerader of the upper abdomen." He described two groups of symptoms which may follow traumatic herniation: (1) Cases in which the stomach only is involved, giving rise to predominantly gastric symptoms; and (2) cases in which other viscera, particularly the colon, become herniated, leading to symptoms of intestinal obstruction. In the above case the two groups of symptoms appeared consecutively.

Harrington stressed the need for early surgical intervention because of the risk of strangulation, a risk which has been emphasized by Borrie and Foreman (1948), and by Carter and Giuseffi (1948).

In discussing the aetiology Jenkinson (1925) maintained that trauma played an unimportant part and that a congenital weakness was present in every case, but this view was not shared by Harrington and Kirklin (1938), or by Harrington (1940), who believed that trauma played a predominant part.

In many cases a congenital deficiency can be found either in the central portion or through the sinus of Morgagni. In the case described above the apparent defect was in the region of the oesophagus, and lends support to Jenkinson's view, though the severity of the injury might well have been expected to rupture even the strongest diaphragm.

In Harrington's series of 60 traumatic herniae, automobile accidents accounted for no fewer than 42, and in the majority of cases quoted by Reid (1840) a history of penetrating injury could be obtained.

I wish to thank Mr. D. B. Craig for permission to publish this case; Mr. Geoffrey Wooler, of Leeds, for helpful criticism; and Dr. Mark Burbury for the radiological examination and reports.

G. R. C. PEATFIELD, F.R.C.S.

REFERENCES

- Blackburn, G., and D'Abreu, A. L. (1945). *Brit. J. Surg.*, 33, 152.
 Borrie, J., and Foreman, H. M. (1948). *Lancet*, 2, 12.
 Carter, B. N., and Giuseffi, J. (1948). *Ann. Surg.*, 128, 210.
 Dugan, D. J., and Sampson, P. C. (1948). *J. thorac. Surg.*, 17, 771.
 Dunhill, Sir T. (1948). In Carling and Ross's *British Surgical Practice*, 4, 451.
 Harrington, S. W. (1940). *Amer. J. Surg.*, 50, 381.
 — (1950). *Surg. Clin. N. Amer.*, 30, 961.
 — and Kirklin, B. R. (1938). *Radiology*, 30, 147.
 Jenkinson, E. L. (1925). *Amer. J. Roentgenol.*, 14, 16.
 Lam, C. R. (1950). *Arch. Surg., Chicago*, 60, 421.
 Reid, J. (1840). *Edinb. med. Surg. J.*, 53, 104.
 Robertson, R. W. (1947). *Kentucky med. J.*, 45, 74.

Closed Incomplete Rupture of Pectoralis Major

A case of this condition due to apparently trivial injury is reported because of its rarity.

CASE REPORT

A healthy man aged 30 was admitted to hospital on December 4, 1952, following an injury at work. While stooping forwards he was struck over his right loin and right buttock by a small light wagon as it passed behind him. As a result he fell forwards across a low wooden rail, the force of his fall being taken on outstretched arms and chest. He did not make any complaint of his chest, being more concerned with pain in his back.

He was transferred to the orthopaedic department on December 9. When first seen he had a large fluctuant swelling over the buttock from which 25 oz. (710 ml.) of old blood was aspirated. X-ray examination of the lumbar spine and pelvis was negative. No complaint of pain in the arm or chest was made. It was not until the patient remarked, some three weeks after coming under our care, that a lump had appeared in his left breast that the present condition was diagnosed.

On examination of the left pectoralis major a gap was readily palpable extending upwards from the lower border

FIG. 1.—Arm relaxed.

FIG. 2.—Arm forcibly adducted against trunk.

of the anterior axillary fold for 2½ in. (6.3 cm.) at the junction of the inner third and outer two-thirds of the fold. On tensing the muscle by active compression of the hand-on-hip position both ends of the torn muscle became more obvious. The clavicular head was quite intact.

COMMENT

It is remarkable that the lesion was not noticed earlier, for it must have been present on admission. Doubtless owing to the patient's attention, and ours, being directed to the back, it passed unnoticed for a time.

No surgical intervention was suggested, for the strength of the arm is unimpaired and the patient is unconcerned about the lump.

I wish to thank Mr. J. P. Heron, under whose care the patient was admitted, for permission to publish this report.

W. B. LAW, M.Ch.Orth., F.R.C.S.
 Assistant Orthopaedic Surgeon. Liverpool.